


1005 RECEIVED 20 MAR 2002

FORM PTO-1390 (REV 11-2000) modified		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 2651 USOP
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 10/088768
INTERNATIONAL APPLICATION NO. PCT/JP00/06376	INTERNATIONAL FILING DATE September 19, 2000	PRIORITY DATE CLAIMED September 20, 1999	
TITLE OF INVENTION MCH ANTAGONISTS			
APPLICANT(S) FOR DO/EO/US Kaneyoshi KATO, Masaaki MORI, Nobuhiro SUZUKI, Yukio SHIMOMURA, Shiro TAKEKAWA, Nobuo CHOH			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
<ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input checked="" type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). <ol style="list-style-type: none"> a. <input checked="" type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)). 9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 			
Items 11 to 20 below concern document(s) or information included:			
<ol style="list-style-type: none"> 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. 14. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 20. <input checked="" type="checkbox"/> Other items or information: <div style="display: flex; justify-content: space-between;"> <div> <p>Itemized Return Postcard; Certificate of Express Mailing</p> <p>Forms PCT/IB/301, 304, 308, 332, 338; PCT Request;</p> <p>Cited References (5)</p> </div> <div> <p>Express Mail Label No. EL 916492903 US</p> <p>Date of Deposit March 20, 2002</p> </div> </div> 			

U.S. APPLICATION NO. 10/068768		INTERNATIONAL APPLICATION NO. PCT/JP00/06376		ATTORNEY'S DOCKET NUMBER 2651 USOP	
21. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
				\$ 890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
Total claims	39 - 20 =	19	x \$18.00	\$ 342.00	
Independent claims	10 - 3 =	7	x \$84.00	\$ 588.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable) <u>2</u>				\$ 560.00	
TOTAL OF ABOVE CALCULATIONS =				\$ 2,380.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
SUBTOTAL =				\$ 2,380.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$ 2,380.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	
TOTAL FEES ENCLOSED =				\$ 2,380.00	
				Amount to be refunded:	\$
				charged:	\$
a. <input type="checkbox"/> A check in the amount of \$ _____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>500799</u> in the amount of \$ <u>2,380.00</u> to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>500799</u> . A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Mark Chao, PhD, JD Takeda Pharmaceuticals North America, Inc. Suite 500, 475 Half Day Road Lincolnshire, IL 60069 USA (847)383-3372 fax (847)383-3481					
				 SIGNATURE Mark Chao, PhD, JD NAME 37,293 REGISTRATION NUMBER For Customer No. 23,115	

SPECIFICATION**MCH ANTAGONISTS****Technical Field**

The present invention relates to a melanin-concentrating
5 hormone antagonist containing an aromatic amine derivative, an
agent for the prophylaxis or therapy of obesity or hyperphagia
and an agent for improving emotional disorder or sexual
dysfunction.

Background Art


10 Feeding behavior is an indispensable action for many
organisms including human. An abnormality in feeding behavior
causes deviation from normal life support activities, which in
most cases results in diseases. Along with the recent changes
in feeding environment, obesity is becoming a social problem.
15 It is widely known that obesity is not only a serious risk
factor of life-style related diseases, such as diabetes,
hypertension, arteriosclerosis and the like, but also causes
arthritis and pain due to excessive burden on joints of knee
etc. created by increased body weight. In addition, the
20 dieting boom and the like have increased the potential
population that desires weight loss. There are many reports on
eating disorders, such as hyperorexia and the like, due to
neuropathy and the like, which are genetic or caused by stress.

Consequently, the development and investigation of an
25 agent for the prophylaxis or therapy of obesity or feeding
deterrents started some time ago and mazindol has been on the
market as a centrally acting anorexiant.

Along therewith, a number of appetite-regulating factors
represented by leptin have been found in recent years, and new
30 anti-obesity agents and anorexiants that suppress the activity
of such appetite-regulating factors have been developed. Among
others, a melanin-concentrating hormone (MCH) is known to be a
hormone derived from hypothalamus and promote appetite.


Furthermore, MCH knockout mouse has been reported to show significantly decreased food intake and be lean, as compared to normal mouse, though normal in daily behavior [Nature, vol. 396, p. 670, 1998]. From the foregoing, an MCH antagonist, once
 5 completed, is expected to be a superior anorexiant or anti-obesity agent. However, a compound having an MCH antagonistic action, particularly a non-peptide compound, has not been known as yet.

JP-A-8-253447 discloses a compound having a gonadotropin
 10 releasing hormone antagonistic action, which is represented by the formula




wherein Ar¹ and Ar² are each an aromatic group optionally having substituents,
 15 P and Q are each a divalent aliphatic hydrocarbon group having 2 or more carbon atoms, which optionally contains ether oxygen or ether sulfur in a carbon chain,
 R¹ and R³ are each i) an acyl group represented by -CO-R or -CONH-R (R is hydrocarbon group optionally having substituents
 20 or heterocyclic group optionally having substituents) or ii) a hydrocarbon group optionally having substituents,
 R² and R⁴ are each a hydrogen atom or an alkyl group optionally having substituents,
 R¹ and R² or R³ and R⁴ may form, together with the adjacent
 25 nitrogen atom, a nitrogen-containing heterocyclic group optionally having substituents, and
 j is 0 or 1, or a salt thereof.

JP-A-10-81665 discloses a compound having an MIP-



wherein Ar¹ and Ar² are each an aromatic group optionally having substituents, P and Q are each a divalent aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which optionally has substituents, R¹ and R³ are each (i) a hydrogen atom, (ii) an acyl group or (iii) a hydrocarbon group optionally having substituents, R² and R⁴ are each (i) a hydrogen atom, (ii) an alkyl group optionally having substituents or (iii) an alkylcarbonyl group optionally having substituents, R¹ and R² or R³ and R⁴ optionally form, together with the adjacent nitrogen atom, a monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents, and j is 0 or 1, or a salt thereof, is not influenced by the presence or the kind of cyclic substituent, but unexpectedly has a superior MCH antagonistic action and low toxicity, and is clinically useful as an agent for the prophylaxis or therapy of obesity or hyperphagia, an agent for improving emotional disorder or sexual dysfunction and the like. Based on the finding, the present inventors have further studied and completed the present invention.

Accordingly, the present invention provides
 [1] a melanin-concentrating hormone antagonist containing a compound of the formula



wherein

Ar¹ and Ar² are each an aromatic group optionally having substituents,

5 P and Q are each a divalent aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which optionally has substituents,

10 R¹ and R³ are each (i) a hydrogen atom, (ii) an acyl group or (iii) a hydrocarbon group optionally having substituents,

R² and R⁴ are each (i) a hydrogen atom, (ii) an alkyl group optionally having substituents or (iii) an alkylcarbonyl group optionally having substituents,

15 R¹ and R² or R³ and R⁴ optionally form, together with the adjacent nitrogen atom, a monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents, and

20 j is 0 or 1,

or a salt thereof or a prodrug thereof;

[2] The antagonist of [1], wherein Ar¹ and Ar² are each (i) a C₆₋₁₄ aryl group or (ii) a 5 to 14-membered monocyclic or fused aromatic heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from the group consisting of nitrogen atom, sulfur atom and oxygen atom, which optionally has 1 to 5 substituent(s) selected from the group (group Aa)

consisting of

- (a) a halogen atom,
- (b) a C₁₋₃ alkylenedioxy group,
- (c) a nitro group,
- 5 (d) a cyano group,
- (e) an optionally halogenated C₁₋₆ alkyl group,
- (f) an optionally halogenated C₃₋₆ cycloalkyl group,
- (g) an optionally halogenated C₁₋₆ alkoxy group,
- (h) an optionally halogenated C₁₋₆ alkylthio group,
- 10 (i) a hydroxy group,
- (j) an amino group,
- (k) a mono-C₁₋₆ alkylamino group,
- (l) a di-C₁₋₆ alkylamino group,
- (m) an optionally halogenated C₁₋₆ alkyl-carbonylamino group,
- 15 (n) a formyl group,
- (o) a C₁₋₆ alkyl-carbonyl group optionally substituted by
halogen atom or C₁₋₆ alkoxy-carbonyl group,
- (p) a C₁₋₆ alkyl-carbonyloxy group,
- (q) a carboxyl group,
- 20 (r) a C₁₋₆ alkoxy-carbonyl group,
- (s) a carbamoyl group,
- (t) a mono-C₁₋₆ alkyl-carbamoyl group optionally substituted by
C₁₋₆ alkoxy-carbonyl group,
- (u) a di-C₁₋₆ alkyl-carbamoyl group optionally substituted by
25 C₁₋₆ alkoxy-carbonyl group,
- (v) a sulfo group,
- (w) a C₁₋₆ alkylsulfonyl group,
- (x) a C₁₋₆ alkylsulfinyl group,
- (y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s)
30 selected from the above-mentioned (a) to (x),
- (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4
substituent(s) selected from the above-mentioned (a) to (x),
- (aa) an optionally halogenated C₆₋₁₀ aryl-carbonyl group,

- (ab) an optionally halogenated 5 or 6-membered heterocyclic ring-carbonyl group,
 - (ac) a C₁₋₆ alkoxy-carbonylamino group,
 - (ad) a C₆₋₁₀ aryl-carbonylamino group and
 - 5 (ae) a C₇₋₁₆ aralkyloxy-carbonyl group,
- P and Q are each a divalent C₁₋₆ aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which is optionally substituted by oxo group or thioxo group;
- 10 R¹ and R³ are each (i) hydrogen atom, (ii) acyl group represented by -CO-R^a, -CONR^aR^b, -SO-R^a, -SO₂-R^a, -CONR^aR^b, -COO-R^a, -(C=S)O-R^a, -(C=S)NR^aR^b, -SONR^aR^b, -SO₂NR^aR^b, -SO-O-R^a or -SO₂-O-R^a, wherein R^a is (A) hydrogen atom; (B) carboxyl group; (C) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl
 - 15 group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋₁₆ aralkyl group, which optionally has 1 to 5 substituent(s) selected from the group (group Ba) consisting of
 - (a) a halogen atom,
 - (b) a C₁₋₃ alkylenedioxy group,
 - 20 (c) a nitro group,
 - (d) a cyano group,
 - (e) an optionally halogenated C₁₋₆ alkyl group,
 - (f) an optionally halogenated C₃₋₆ cycloalkyl group,
 - (g) an optionally halogenated C₁₋₆ alkoxy group,
 - 25 (h) an optionally halogenated C₁₋₆ alkylthio group,
 - (i) a hydroxy group,
 - (j) an amino group,
 - (k) a mono-C₁₋₆ alkylamino group,
 - (l) a di-C₁₋₆ alkylamino group,
 - 30 (m) a C₁₋₆ alkyl-carbonylamino group,
 - (n) a formyl group,
 - (o) a C₁₋₆ alkyl-carbonyl group,
 - (p) a C₁₋₆ alkyl-carbonyloxy group,

- (q) a carboxyl group,
- (r) a C₁₋₆ alkoxy-carbonyl group,
- (s) a carbamoyl group,
- (t) a mono-C₁₋₆ alkyl-carbamoyl group,
- 5 (u) a di-C₁₋₆ alkyl-carbamoyl group,
- (v) a sulfo group,
- (w) a C₁₋₆ alkylsulfonyl group,
- (x) a C₁₋₆ alkylsulfinyl group,
- (y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s)
- 10 selected from the aforementioned (a) to (x),
- (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4
- substituent(s) selected from the aforementioned (a) to (x),
- (zz) a 5 to 7-membered heterocyclic group optionally having 1
- to 4 substituent(s) selected from the aforementioned (a) to (x),
- 15 (aa) a di-C₁₋₆ alkyl-carbonylamino group,
- (ab) a sulfamoyl group,
- (ac) a C₁₋₆ alkoxy-carbonylamino group,
- (ad) a C₇₋₁₆ aralkyloxy-carbonylamino group,
- (ae) a C₇₋₁₆ aralkyloxy group,
- 20 (af) a C₆₋₁₀ aryl-carbonyl group,
- (ag) a C₁₋₆ alkyl-carbonyloxy group,
- (ah) a C₆₋₁₀ aryl-carbonylamino group,
- (ai) a C₆₋₁₀ aryl-carbamoyl group,
- (aj) a C₇₋₁₆ aralkylaminocarbonyl group,
- 25 (ak) a C₇₋₁₆ aralkylcarbonylamino group and
- (al) a C₇₋₁₆ aralkyloxy-carbonyloxy group;
- (D) a 5 to 10-membered heterocyclic group containing, besides
- carbon atom, 1 to 4 heteroatom(s) selected from the group
- consisting of nitrogen atom, oxygen atom and sulfur atom, which
- 30 optionally has 1 to 5 substituent(s) selected from the group
- consisting of (a) substituent selected from group Aa,
- (b) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl
- group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋

₁₆ aralkyl group, which optionally has 1 to 5 substituent(s) selected from group Ba,

(c) oxo group and

(d) thioxo group; or

⁵ (E) a C₁₋₆ alkoxy-carbonyl group;

R^b is a hydrogen atom or a C₁₋₆ alkyl group, or

(iii) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋₁₆ aralkyl group optionally having 1 to 5 substituent(s)

¹⁰ selected from group Ba;

R² and R⁴ are each (i) a hydrogen atom, (ii) C₁₋₆ alkyl group optionally having substituents selected from group Ba or (iii) C₁₋₆ alkyl-carbonyl group optionally having substituents selected from group Ba;

¹⁵ R¹ and R² or R³ and R⁴ may form, together with the adjacent nitrogen atom, a group of

(i) the formula



wherein ring A is a 4 to 8-membered ring optionally substituted
²⁰ by hydroxy or oxo, V is a group represented by the formula >O, >C=O, >C(W)-W^a or >N-W (W is (a) hydrogen atom, (b) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋₁₆ aralkyl group, which optionally has 1 to 5 substituent(s) selected from
²⁵ group Ba, or (c) 5 to 10-membered heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from nitrogen, oxygen and sulfur, which optionally has 1 to 5 substituent(s) selected from group Aa, W^a is hydrogen atom, hydroxy group or C₁₋₆ alkyl group),

³⁰ (ii) the formula

- (d) a cyano group,
- (e) an optionally halogenated C₁₋₆ alkyl group,
- (f) an optionally halogenated C₃₋₆ cycloalkyl group,
- (g) an optionally halogenated C₁₋₆ alkoxy group,
- 5 (h) an optionally halogenated C₁₋₆ alkylthio group,
- (i) a hydroxy group,
- (j) an amino group,
- (k) a mono-C₁₋₆ alkylamino group,
- (l) a di-C₁₋₆ alkylamino group,
- 10 (m) a C₁₋₆ alkyl-carbonylamino group,
- (n) a formyl group,
- (o) a C₁₋₆ alkyl-carbonyl group,
- (p) a C₁₋₆ alkyl-carbonyloxy group,
- (q) a carboxyl group,
- 15 (r) a C₁₋₆ alkoxy-carbonyl group,
- (s) a carbamoyl group,
- (t) a mono-C₁₋₆ alkylcarbamoyl group,
- (u) a di-C₁₋₆ alkylcarbamoyl group,
- (v) a sulfo group,
- 20 (w) a C₁₋₆ alkylsulfonyl group,
- (x) a C₁₋₆ alkylsulfinyl group,
- (y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s) selected from the above-mentioned (a) to (x) and
- (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4
- 25 substituent(s) selected from the above-mentioned (a) to (x),
- P and Q are each a C₁₋₆ aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which is optionally substituted by oxo group or thioxo group,
- 30 R¹ and R³ are each (i) hydrogen atom, (ii) an acyl group represented by -CO-R^a, -CONR^aR^b, -SO-R^a, -SO₂-R^a, -CONR^aR^b, -COO-R^a, -(C=S)O-R^a or -(C=S)NR^aR^b wherein R^a is (a) hydrogen atom, (b) carboxyl group,

- (c) a (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋₁₆ aralkyl group, which optionally has 1 to 5 substituent(s) selected from the group (group B) consisting of
- 5 (a) a halogen atom,
 - (b) a C₁₋₃ alkylenedioxy group,
 - (c) a nitro group,
 - (d) a cyano group,
 - (e) an optionally halogenated C₁₋₆ alkyl group,
 - 10 (f) an optionally halogenated C₃₋₆ cycloalkyl group,
 - (g) an optionally halogenated C₁₋₆ alkoxy group,
 - (h) an optionally halogenated C₁₋₆ alkylthio group,
 - (i) a hydroxy group,
 - (j) an amino group,
 - 15 (k) a mono-C₁₋₆ alkylamino group,
 - (l) a di-C₁₋₆ alkylamino group,
 - (m) a C₁₋₆ alkyl-carbonylamino group,
 - (n) a formyl group,
 - (o) a C₁₋₆ alkyl-carbonyl group,
 - 20 (p) a C₁₋₆ alkyl-carboxyloxy group,
 - (q) a carboxyl group,
 - (r) a C₁₋₆ alkoxy-carbonyl group,
 - (s) a carbamoyl group,
 - (t) a mono-C₁₋₆ alkylcarbamoyl group,
 - 25 (u) a di-C₁₋₆ alkylcarbamoyl group,
 - (v) a sulfo group,
 - (w) a C₁₋₆ alkylsulfonyl group,
 - (x) a C₁₋₆ alkylsulfinyl group,
 - (y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s)
 - 30 selected from the aforementioned (a) to (x),
 - (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4 substituent(s) selected from the aforementioned (a) to (x) and
 - (zz) a 5 to 7-membered heterocyclic group optionally having 1


to 4 substituent(s) selected from the aforementioned (a) to (x),
or

- (d) a 5 to 10-membered heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, which optionally has 1 to 5 substituent(s) selected from the group (group C) consisting of
- (a) a halogen atom,
 - (b) a C₁₋₃ alkylenedioxy group,
 - 10 (c) a nitro group,
 - (d) a cyano group,
 - (e) a C₁₋₆ alkyl group optionally having substituents selected from the group consisting of (aa) a halogen atom, (bb) C₁₋₃ alkylenedioxy group, (cc) nitro group, (dd) cyano group, (ee)
 - 15 an optionally halogenated C₁₋₆ alkyl group, (ff) an optionally halogenated C₃₋₆ cycloalkyl group, (gg) an optionally halogenated C₁₋₆ alkoxy group, (hh) an optionally halogenated C₁₋₆ alkylthio group, (ii) a hydroxy group, (jj) amino group, (kk) a mono-C₁₋₆ alkylamino group, (ll) a di-C₁₋₆ alkylamino
 - 20 group, (mm) C₁₋₆ alkyl-carbonylamino group, (nn) a formyl group, (oo) C₁₋₆ alkyl-carbonyl group, (pp) C₁₋₆ alkyl-carboxyloxy group, (qq) carboxyl group, (rr) C₁₋₆ alkoxy-carbonyl group, (ss) carbamoyl group, (tt) a mono-C₁₋₆ alkylcarbamoyl group, (uu) a di-C₁₋₆ alkylcarbamoyl group, (vv) a sulfo group, (ww) C₁₋₆
 - 25 alkylsulfonyl group, (xx) C₁₋₆ alkylsulfinyl group, (yy) C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s) selected from the aforementioned (aa) to (xx), (zz) C₆₋₁₀ aryloxy group optionally having 1 to 4 substituent(s) selected from the aforementioned (aa) to (xx) and (zzz) 5 to 7-membered
 - 30 heterocyclic group optionally having 1 to 4 substituent(s) selected from the aforementioned (aa) to (xx),
 - (f) an optionally halogenated C₃₋₆ cycloalkyl group,
 - (g) an optionally halogenated C₁₋₆ alkoxy group,


- (h) an optionally halogenated C₁₋₆ alkylthio group,
(i) a hydroxy group,
(j) an amino group,
(k) a mono-C₁₋₆ alkylamino group,
5 (l) a di-C₁₋₆ alkylamino group,
(m) an optionally halogenated C₁₋₆ alkyl-carbonylamino group,
(n) a formyl group,
(o) a C₁₋₆ alkyl-carbonyl group,
(p) a C₁₋₆ alkyl-carbonyloxy group,
10 (q) a carboxyl group,
(r) a C₁₋₆ alkoxy-carbonyl group,
(s) a carbamoyl group,
(t) a mono-C₁₋₆ alkylcarbamoyl group,
(u) a di-C₁₋₆ alkylcarbamoyl group,
15 (v) a sulfo group,
(w) a C₁₋₆ alkylsulfonyl group,
(x) a C₁₋₆ alkylsulfinyl group,
(y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s)
selected from the aforementioned (a) to (x) and
20 (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4
substituent(s) selected from the aforementioned (a) to (x), and
R^b is a hydrogen atom or a C₁₋₆ alkyl group) or
(iii) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆
alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or
25 (6) C₇₋₁₆ aralkyl group, which optionally has 1 to 5
substituent(s) selected from group B,
R² and R⁴ are each (i) hydrogen atom, (ii) C₁₋₆ alkyl optionally
having substituents selected from group B or (iii) C₁₋₆ alkyl-
carbonyl group optionally having substituents selected from
30 group B,
R¹ and R² or R³ and R⁴ form, together with the adjacent nitrogen
atom, a nitrogen-containing heterocyclic group represented by
(i) the formula



- wherein ring A is a 4 to 8-membered ring optionally substituted by hydroxy or oxo, V is a group represented by the formula $>O$, $>C=O$, $>C-(W)W^a$ or $>N-W$ (W is (a) hydrogen atom, (b) (1) C_{1-6} alkyl group, (2) C_{2-6} alkenyl group, (3) C_{2-6} alkynyl group, (4) C_{3-6} cycloalkyl group, (5) C_{6-14} aryl group or (6) C_{7-16} aralkyl group, which optionally has 1 to 5 substituent(s) selected from group B, or (c) 5 to 10-membered heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from nitrogen, oxygen and sulfur, which optionally has 1 to 5 substituent(s) selected from group A, W^a is hydrogen atom or hydroxy group),
- (ii) the formula



- wherein ring B is monocyclic or bicyclic 4 to 12-membered ring optionally substituted by oxo group or 1 to 5 C_{1-6} alkyl group(s), ring D is a 4 to 12-membered aromatic ring optionally having 1 to 5 substituent(s) selected from group A or
- (iii) the formula



- wherein ring E is a 5 to 10-membered aromatic ring optionally having 1 to 5 substituent(s) selected from group A
- X is $-CH_2-$, $-CO-$ or $-CH(OH)-$,
- Y is $-CH_2-$, $-O-$ or $-NW^b-$ (W^b is (a) hydrogen atom or (b) C_{1-6} alkyl group optionally having substituents selected from group

B) ;

$k+m$ is an integer of 1 to 4; and

n is an integer of 1 to 3,

[4] the antagonist of [1], wherein Ar¹ and Ar² are each (i) a phenyl group optionally substituted by halogen atom or C₁₋₆ alkoxy group or (ii) a 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom,

[5] the antagonist of [1], wherein P and Q are each a C₁₋₆¹⁰ alkylene group,

[6] the antagonist of [1], wherein j is 0,

[7] the antagonist of [1], wherein

R¹ is (i) C₁₋₆ alkyl group optionally having a 5 or 6-membered nitrogen-containing heterocyclic group, (ii) C₇₋₁₆ aralkyl group optionally having nitro, amino or C₁₋₆ alkoxy-carbonyl or (iii) cyclohexyl group fused with benzene ring optionally having C₁₋₆ alkoxy;

R² is (i) hydrogen atom, (ii) C₁₋₆ alkyl group or (iii) C₇₋₁₆ aralkyl group; or R¹ and R² form, together with the adjacent nitrogen atom, a nitrogen-containing heterocyclic group represented by


(i) the formula




wherein ring A¹ is a 4 to 8-membered ring optionally substituted by hydroxy or oxo, V¹ is a group represented by the formula >O, >C(W¹)-W^{a1} or >N-W¹ (W¹ is (a) hydrogen atom, (b) C₆₋₁₄ aryl group optionally having 1 or 2 substituent(s) selected from the group consisting of a halogen atom, optionally halogenated C₁₋₆ alkyl group and optionally halogenated C₁₋₆ alkoxy group, (c) C₁₋₆ alkyl group optionally having 1 or 2 C₆₋₁₀ aryl group(s) or (d) pyridyl group, W^{a1} is

hydrogen atom, hydroxy group or C₁₋₆ alkyl group),

(ii) the formula




wherein ring B¹ is a monocyclic or bicyclic 5 to 10-membered
 5 ring optionally substituted by oxo group or 1 or 2 C₁₋₆ alkyl
 group(s), ring D¹ is a benzene ring optionally having 1 or 2
 substituent(s) selected from the group consisting of C₁₋₆ alkyl
 group, C₁₋₆ alkoxy group and C₁₋₆ alkyl-carbonyl group or
 (iii) the formula




wherein ring E¹ is a benzene ring optionally having 1 to 3
 substituent(s) selected from the group consisting of C₁₋₃
 alkylenedioxy group, nitro group, C₁₋₆ alkoxy group, amino group,
 C₁₋₆ alkyl-carbonylamino group and C₁₋₆ alkoxy-carbonyl group,
 15 X¹ is -CH₂- or -CO-, and
 Y¹ is -CH₂- or -O-,
 R³ is (i) hydrogen atom,
 (ii) a group represented by the formula -CO-R⁵ (R⁵ is (a)
 hydrogen atom, (b) carboxyl group, (c) C₁₋₆ alkyl group, (d) C₅₋₆
 20 cycloalkyl group optionally having C₁₋₆ alkoxy, and which is
 fused with benzene ring or (e) 5 or 6-membered aromatic
 heterocyclic group containing, besides carbon atom, 1 to 3
 heteroatom(s) selected from the group consisting of nitrogen
 atom, oxygen atom and sulfur atom, which optionally has 1 or 2
 25 substituent(s) selected from the group consisting of a halogen
 atom, C₆₋₁₀ aryl group, C₆₋₁₀ aryl-carbonylamino group),
 (iii) a group represented by the formula -CO-Alk₀-R⁶ [Alk₀ is
 C₁₋₆ alkylene group optionally having hydroxy group, R⁶ is (a)

C₆₋₁₄ aryl group optionally having 1 or 2 substituent(s) selected from the group consisting of a halogen atom, optionally halogenated C₁₋₆ alkyl, nitro, C₁₋₆ alkoxy, C₁₋₃ alkylenedioxy and C₆₋₁₀ aryl group, (b) C₆₋₁₀ aryloxy group, (c) 5 or 6-membered aromatic heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom (d) C₁₋₆ alkyl-carbonyl group, (e) carboxyl group, (f) C₁₋₆ alkoxy-carbonyl group, (g) amino group optionally having 1 or 2 substituent(s) selected from the group consisting of C₁₋₆ alkyl and C₁₋₆ alkyl-carbonyl, (h) 5 to 7-membered heterocyclic group optionally having hydroxy, (i) C₇₋₁₆ aralkyloxy group, (j) C₆₋₁₀ aryl-carbonyl group or (k) C₁₋₆ alkyl-carbonyloxy group],
(iv) a group represented by the formula



wherein Qa is a group represented by the formula $-(CH_2)_s-$ (s is an integer of 1 to 3) or $-(CH_2)_t-CH=$ (t is an integer of 0 to 2) and R⁷ is hydrogen atom or C₁₋₆ alkoxy-carbonyl group,

(v) a group represented by the formula




wherein R⁸ is (a) hydrogen atom, (b) C₁₋₆ alkyl group optionally having substituents selected from the group consisting of C₁₋₆ alkoxy-carbonyl, morpholino and mono- or di-C₁₋₆ alkylamino, (c) C₁₋₆ alkoxy-carbonyl group, (d) a group represented by the formula -CO-R^d (R^d is C₆₋₁₀ aryl group optionally having halogen atom or 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom), (e) a group represented by the formula -CO-(CH₂)^{r¹}-R^e (r¹ is an integer of 1 to 3, R^e is C₁₋₆ alkoxy-carbonyl group or 5 or 6-membered heterocyclic group

containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom) or (f) a group represented by $-\text{CONH}-\text{R}^f$ (R^f is C_{1-6} alkyl group or C_{6-14} aryl group),

5 (vi) a group represented by the formula $-\text{COOR}^9$ (R^9 is optionally halogenated C_{1-6} alkyl group),


(vii) a group represented by the formula



wherein R^{10} is hydrogen atom, C_{1-6} alkoxy-carbonyl group, mono-
10 or di- C_{1-6} alkyl-carbamoyl group, optionally halogenated nicotinoyl group or optionally halogenated isonicotinoyl group,


(viii) a group represented by the formula $-\text{CONR}^{11}-\text{R}^{12}$ (R^{11} is hydrogen atom or C_{1-6} alkyl group, R^{12} is C_{1-6} alkyl group optionally having substituents selected from the group
15 consisting of (a) hydroxy, (b) amino, (c) a mono- or di- C_{1-6} alkyl-amino, (d) C_{1-6} alkyl-carbonyl, (e) C_{1-6} alkoxy-carbonyl, (f) C_{1-6} alkyl-carbonyloxy, (g) sulfamoyl and (h) 5 to 7-membered heterocyclic group optionally substituted by oxo, and (i) C_{6-14} aryl group),

20 (ix) a group represented by the formula



wherein R^{13} is (a) hydrogen atom, (b) C_{1-6} alkyl group optionally having substituents selected from the group consisting of a hydroxy and C_{1-6} alkoxy-carbonyl, (c) C_{7-16}
25 aralkyl group, (d) C_{1-6} alkyl-carbonyl group optionally having substituents selected from the group consisting of a halogen atom and C_{1-6} alkoxy-carbonyl or (e) C_{1-6} alkyl-carbamoyl group optionally having C_{1-6} alkoxy-carbonyl,

(x) a group represented by the formula



(xxi) a group represented by the formula $-\text{CO}-\text{Alk}-\text{NR}^{27}-\text{CO}-\text{Alk}_2-\text{NR}^{32}-\text{CO}-\text{O}-\text{Alk}_3-\text{R}^{31}$

[Alk, R^{27} , Alk_2 , Alk_3 and R^{31} are as defined above; and R^{32} is the same as the aforementioned R^{27}];

5 (xxii) a group represented by the formula $-\text{CO}-\text{Alk}-\text{CO}-\text{NR}^{27}-\text{Alk}_2-\text{R}^{31}$


[Alk, R^{27} , Alk_2 and R^{31} are as defined above]; or

(xxiii) a group represented by the formula $-\text{CO}-\text{Alk}-\text{O}-\text{CO}-\text{O}-\text{Alk}_2-\text{R}^{31}$

10 [Alk, Alk_2 and R^{31} are as defined above];

R^4 is hydrogen atom or C_{1-6} alkyl group;

or R^3 and R^4 may form, together with the adjacent nitrogen atom, a group represented by the formula



15 wherein R^{18} is halogen atom, oxo group, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group,

[8] the antagonist of [1] wherein R^1 is (i) C_{1-6} alkyl group optionally having 5 or 6-membered nitrogen-containing

20 heterocyclic group, (ii) C_{7-16} aralkyl group optionally having nitro, amino or C_{1-6} alkoxy-carbonyl or (iii) cyclohexyl group fused with benzene ring optionally having C_{1-6} alkoxy,

R^2 is (i) hydrogen atom, (ii) C_{1-6} alkyl group or (iii) C_{7-16} aralkyl group, or R^1 and R^2 may form, together with the adjacent nitrogen atom, a nitrogen-containing heterocyclic

25 group represented by


(i) the formula



wherein ring A^1 is a 4 to 8-membered ring optionally substituted by hydroxy or oxo, V^1 is a group represented by the


(iii) a group represented by the formula $-\text{CO}-(\text{CH}_2)_{r^1}-\text{R}^6$ (r^1 is an integer of 1 to 3; R^6 is (a) C_{6-14} aryl group optionally having 1 or 2 substituent(s) selected from the group consisting of halogen atom, optionally halogenated C_{1-6} alkyl, nitro, C_{1-6} alkoxy and C_{1-3} alkylendioxy, (b) C_{6-14} aryloxy group, (c) 5 or 6-membered aromatic heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom, (d) C_{1-6} alkyl-carbonyl group, (e) carboxyl group, (f) C_{1-6} alkoxy-carbonyl group, (g) amino group optionally having 1 or 2 substituent(s) selected from the group consisting of C_{1-6} alkyl and C_{1-6} alkyl-carbonyl or (h) 5 or 6-membered cyclic amino group optionally having hydroxy),

(iv) a group represented by the formula




(Q is a group represented by the formula $-(\text{CH}_2)_s-$ (s is an integer of 1 to 3) or $-(\text{CH}_2)_t-\text{CH}=-$ (t is an integer of 0 to 2), R^7 is hydrogen atom or C_{1-6} alkoxy-carbonyl group),

(v) a group represented by the formula



(R^8 is (a) hydrogen atom, (b) C_{1-6} alkyl group optionally having substituents selected from the group consisting of C_{1-6} alkoxy-carbonyl, morpholino and mono- or di- C_{1-6} alkylamino, (c) C_{1-6} alkoxy-carbonyl group, (d) a group represented by the formula $-\text{CO}-\text{R}^d$ (R^d is C_{6-14} aryl group optionally having halogen atom or 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom),


(e) a group represented by the formula $-\text{CO}-(\text{CH}_2)_{r^1}-\text{R}^e$ (r^1 is an integer of 1 to 3, R^e is C_{1-6} alkoxy-carbonyl group or 5 or 6-membered heterocyclic group containing, besides carbon atom, 1



wherein ring F is 5 to 7-membered cyclic amino group optionally fused with benzene ring, R¹⁵ is hydrogen atom, C₁₋₆ alkoxy-carbonylamino group or optionally halogenated C₁₋₆ alkyl-

carbonylamino group,

(xii) a group represented by the formula



wherein R¹⁶ is (a) C₁₋₆ alkyl group optionally having substituents selected from the group consisting of a hydroxy and C₁₋₆ alkoxy-carbonyl, (b) a formyl group, (c) C₁₋₆ alkoxy-carbonyl group or (d) a 5 or 6-membered heterocyclic ring-carbonyl group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom,


(xiii) a group represented by the formula -SO₂-R¹⁷ (R¹⁷ is (i) C₁₋₆ alkyl group optionally having 5 or 6-membered nitrogen-containing ring group, (ii) C₂₋₆ alkenyl group or (iii) C₆₋₁₄ aryl group optionally having C₁₋₆ alkyl),

(xiv) C₇₋₁₆ aralkyl group optionally having 1 to 3 halogen atom(s), or

(xv) C₁₋₆ alkyl group substituted by 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom,


R⁴ is hydrogen atom or C₁₋₆ alkyl group,

or R³ and R⁴ may form, together with the adjacent nitrogen atom, a group of the formula



wherein R^{18} is halogen atom, oxo group, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group,
 [9] the antagonist of [1] wherein R^1 and R^2 form, together with the adjacent nitrogen atom, a nitrogen-containing heterocyclic
 5 group represented by

(i) the formula




wherein q^1 is hydrogen atom or halogen atom, q^2 is hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or
 10 C_{1-6} alkoxy group, q^3 is hydrogen atom or halogen atom, q^4 is hydrogen atom, halogen atom or C_{1-6} alkoxy group, q^5 is hydrogen atom or C_{1-6} alkyl group optionally having 1 or 2 C_{6-10} aryl group(s),

(ii) the formula




wherein ring B^2 is represented by the formula




wherein ring D^1 is benzene ring optionally having 1 or 2 substituent(s) selected from the group consisting of C_{1-6} alkyl

group, C₁₋₆ alkoxy group and C₁₋₆ alkyl-carbonyl group or
(iii) the formula




wherein ring E¹ is benzene ring optionally having 1 or 2
5 substituent(s) selected from the group consisting of C₁₋₃
alkylenedioxy group, nitro group, C₁₋₆ alkoxy group, amino group,
C₁₋₆ alkyl-carbonylamino group and C₁₋₆ alkoxy-carbonyl group, X¹
is -CH₂- or -CO-, and Y¹ is -CH₂- or -O-),


[10] the antagonist of [1] wherein the compound is represented
10 by the formula




wherein R¹⁹ is (i) hydrogen atom, (ii) carboxyl, (iii) C₁₋₆
alkoxy-carbonyl group, (iv) C₁₋₆ alkyl group optionally having
substituents selected from the group consisting of carboxyl,
15 C₁₋₆ alkyl-carbonyl, C₁₋₆ alkoxy-carbonyl, C₁₋₆ alkoxy-
carbonylamino and C₇₋₁₆ aralkyloxy-carbonylamino, (v) a mono- or
di-C₁₋₆ alkylamino group or (iv) C₆₋₁₄ aryloxy group; P¹ is C₁₋₃
alkylene group; Q¹ is C₁₋₃ alkylene group; X² is CH, C-OH or N;
Y² is hydrogen atom, halogen atom, optionally halogenated C₁₋₆
20 alkyl group or C₁₋₆ alkoxy group; and Z is CO, SO or SO₂,
[11] the antagonist of [1] wherein the compound is represented
by the formula




wherein R^{20} is (i) hydrogen atom or (ii) C_{1-6} alkyl group optionally having substituents selected from the group consisting of C_{1-6} alkoxy-carbonylamino and C_{7-16} aralkyloxy-carbonylamino; P^2 is C_{1-3} alkylene group; X^3 is CH, C-OH or N; Y^3 is hydrogen atom, halogen atom or C_{1-6} alkoxy group, [12] the antagonist of [1] wherein the compound is represented by the formula



wherein R^{21} is a nitrogen-containing heterocyclic group represented by (i) the formula



wherein X^4 is CH or N, Y^4 is hydrogen atom, halogen atom or C_{1-6} alkoxy group or (ii) the formula



15

wherein ring E^2 is benzene ring optionally having 1 to 3 C_{1-6} alkoxy,

R^{22} is (i) hydrogen atom, (ii) C_{7-16} aralkyl group, (iii) formyl group, (iv) C_{1-6} alkyl-carbonyl group, (v) C_{6-14} aryl-carbonyl group optionally having C_{1-6} alkyl or (vi) C_{6-14} aryl-sulfonyl group optionally having 1 to 4 C_{1-6} alkyl; P^3 is C_{1-3} alkylene

group; and Q³ is C₁₋₃ alkylene group,
 [13] the antagonist of [1] wherein the compound is
 1-(5-amino-4,4-diphenylpentyl)-4-phenylpiperidine,
 3,4-dihydro-6-methoxy-1'-(5-amino-4,4-
 5 diphenylpentyl)spiro[naphthalene-2(1H),2'-piperidine] or a salt
 thereof,
 1-[5-amino-4-(4-methoxyphenyl)-4-phenylpentyl]-4-
 phenylpiperidine or a salt thereof,
 1-[5-amino-4,4-bis(4-chlorophenyl)pentyl]-4-(4-
 10 fluorophenyl)piperazine or a salt thereof,
 3,4-dihydro-6-methoxy-1'-(6-amino-4,4-
 diphenylhexyl)spiro[naphthalene-2(1H),2'-piperidine] or a salt
 thereof,
 3,4-dihydro-6,7-dimethoxy-1'-(7-amino-4,4-
 15 diphenylheptyl)spiro[naphthalene-2(1H),2'-piperidine] or a salt
 thereof,
 4,4-diphenyl-5-formylamino-1-(4-phenylpiperidino)pentane or a
 salt thereof (e.g., hydrochloride),
 1-[4-(4-fluorophenyl)piperazin-1-yl]-5-formylamino-4,4-
 20 diphenylpentane or a salt thereof (e.g., dihydrochloride),
 4,4-diphenyl-1-(4-phenylpiperazin-1-yl)-5-(tosylamino)pentane
 or a salt thereof,
 4,4-diphenyl-1-[4-(2-methoxyphenyl)piperazin-1-yl]-5-
 (tosylamino)pentane or a salt thereof (e.g., hydrochloride),
 25 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-
 phenylpiperidino)pentane or a salt thereof (e.g.,
 hydrochloride),
 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-
 phenylpiperazin-1-yl)pentane or a salt thereof (e.g.,
 30 dihydrochloride),
 4-(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-
 formylamino-4-phenylpentane or a salt thereof (e.g.,
 dihydrochloride),

- 4-(4-chlorophenyl)-1-[4-(diphenylmethyl)piperazin-1-yl]-5-formylamino-4-phenylpentane or a salt thereof,
- 5-formylamino-4-(4-methoxyphenyl)-4-phenyl-1-(4-phenylpiperidino)pentane or a salt thereof (e.g., hydrochloride),
- 4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(formylamino)pentane or a salt thereof (e.g., dihydrochloride),
- 1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-5,5-diphenylhexane or a salt thereof (e.g., dihydrochloride),
- 1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-4,4-diphenylhexane or a salt thereof (e.g., dihydrochloride),
- 4,4-diphenyl-1-(4-phenylpiperidino)-6-(tosylamino)hexane or a salt thereof (e.g., hydrochloride),
- 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 5-[4-(4-fluorophenyl)piperazin-1-yl]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., dihydrochloride),
- 1-formylamino-5-(4-hydroxy-4-phenylpiperidino)-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 5-[4-(4-trifluoromethylphenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 5-[4-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 5-[4-(3,5-dichlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 5-[4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridin-1-yl]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 1-formylamino-2,2-diphenyl-5-(4-phenylpiperidino)pentane or a salt thereof,

- 5-[4-(4-chlorophenyl)piperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 7-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-4,4-diphenylheptane or a salt thereof (e.g., hydrochloride),
- 5 5-[4-(4-fluorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 1-formylamino-5-[4-hydroxy-4-(4-methoxyphenyl)piperidino]-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 1-formylamino-5-[4-hydroxy-4-(2-pyridyl)piperidino]-2,2-
- 10 diphenylpentane or a salt thereof (e.g., dihydrochloride),
- 1-acetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 1-acetoacetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane or a salt thereof (e.g., hydrochloride),
- 15 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamate or a salt thereof (e.g., hydrochloride),
- N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid or a salt thereof,
- 20 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-ethylurea or a salt thereof,
- N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]methanesulfonamide or a salt thereof (e.g., hydrochloride),
- 25 phenyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]carbamate or a salt thereof,
- 1-acetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentane or a salt thereof (e.g., dihydrochloride),
- 30 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]oxamate or a salt thereof (e.g., hydrochloride),
- ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]malonamate or a salt thereof (e.g.,


hydrochloride),
 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glutaramate or a salt thereof,
 benzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-
 5 oxoethylcarbamate or a salt thereof (e.g., hydrochloride),
 tert-butyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate or a salt thereof,
 4,4-diphenyl-7-(4-phenylpiperidino)heptylamine or a salt
 10 thereof (e.g., dihydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-4-methylbenzenesulfonamide or a salt thereof (e.g., hydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)acetamide or a
 15 salt thereof (e.g., hydrochloride),
 N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)amine or a salt thereof (e.g., dihydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(3-methoxybenzyl)amine or a salt thereof (e.g., dihydrochloride),
 20 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-methoxybenzyl)amine or a salt thereof (e.g., dihydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-fluorobenzyl)amine or a salt thereof (e.g., dihydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-
 25 thiophenecarboxamide or a salt thereof (e.g., hydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-phenylacetamide or a salt thereof (e.g., hydrochloride),
 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-thienylmethyl)amine or a salt thereof (e.g., dihydrochloride),
 30 or
 N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-methylamine or a salt thereof (e.g., dihydrochloride),
 [14] the antagonist of [1] which is an agent for the

prophylaxis or therapy of a disease caused by melanin-concentrating hormone,

[15] the antagonist of [1] which is an agent for the prophylaxis or therapy of obesity,

5 [16] the antagonist of [1] which is an agent for suppressing food intake,

[17] a compound represented by the formula



wherein R^{23} is C_{1-6} alkyl group having C_{7-16} aralkyloxy-carbonylamino optionally having substituents selected from the group consisting of halogen atom, C_{1-6} alkoxy and C_{1-6} alkyl; P^4 is C_{1-3} alkylene group; X^5 is CH, C-OH or N; Y^5 is hydrogen atom, halogen atom or C_{1-6} alkoxy group; R^{26} is hydrogen atom or C_{1-6} alkyl group; Y^6 and Y^7 are the same or different and each is
 15 hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group, or a salt thereof or a prodrug thereof,

[18] the compound of [17] wherein R^{26} is hydrogen atom,

[19] benzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethylcarbamate (Example 1) or a salt thereof,
 20 4-chlorobenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethylcarbamate (Example 57) or a salt thereof,

3-chlorobenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethylcarbamate (Example 58) or a salt thereof,


25 benzyl 2-(N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-N-methylamino)-2-oxoethylcarbamate (Example 75) or a salt thereof, benzyl 2-((5-(4-(3-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate (Example 76) or a salt thereof,

benzyl 2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate (Example 77) or a salt thereof,

benzyl 2-((5-(4-(2-methoxyphenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate (Example 80) or a salt thereof, or

3-chlorobenzyl 2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate (Example 103) or a salt thereof,

[20] a production method of a compound of [17], which comprises reacting a compound represented by the formula




wherein each symbol is as defined in [17] or a salt thereof with a reactive derivative of an organic acid of the formula

R²³-COOH

wherein R²³ is as defined in [17],

[21] a production method of a compound of [17], which comprises reacting a compound represented by the formula



wherein each symbol is as defined in [17] or a salt thereof with a reactive derivative of the formula

R³²-X

wherein R³² is C₇₋₁₆ aralkyloxy-carbonyl group, and X is a leaving group,

[22] a pharmaceutical composition containing a compound of [17],

[illegible]

¹⁵ halogenated C₁₋₆ alkoxy group, or a salt thereof or a prodrug thereof.

25 substituents selected from the group consisting of nitro, halogen atom, C₁₋₆ alkyl, hydroxy, C₁₋₆ alkoxy and C₆₋₁₀ aryl, [25] the compound of [23] or [24], wherein R²⁹ is indol-2-yl optionally having substituents,

[26] the compound of [23] or [24], wherein R²⁹ is indol-2-yl

[27] N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide (Example 44) or a salt thereof,

5 N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-1-methylindole-2-carboxamide (Example 45) or a salt thereof.

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)indole-2-carboxamide (Example 104) or a salt thereof,

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-
¹⁵ amino)-2-oxoethyl)-5-chloroindole-2-carboxamide (Example 105)
 or a salt thereof,

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-1-methylindole-2-carboxamide (Example 106) or a salt thereof,

20 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-
amino)-2-oxoethyl)-5-fluoroindole-2-carboxamide (Example 107)
or a salt thereof.

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-5-methoxyindole-2-carboxamide (Example 108)

25 or a salt thereof,

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-5-hydroxyindole-2-carboxamide (Example 109) or a salt thereof,

N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)-
30 amino)-2-oxoethyl)indole-2-carboxamide (Example 115) or a salt
thereof,

N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)-amino)-2-oxoethyl)-1-methylindole-2-carboxamide (Example 116)

or a salt thereof,

5-chloro-N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethyl)-1-methylindole-2-carboxamide (Example 117) or a salt thereof,

5 5-chloro-N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethyl)indole-2-carboxamide (Example 118) or a salt thereof,

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-fluoroindole-2-

10 carboxamide (Example 120) or a salt thereof,

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-methoxyindole-2-carboxamide (Example 121) or a salt thereof,

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-

15 piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide (Example 122) or a salt thereof,

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)indole-2-carboxamide (Example 124) or a salt thereof,

20 N-(2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-5-chloroindole-2-carboxamide (Example 125) or a salt thereof,

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide

25 (Example 127) or a salt thereof,

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-chloroindole-2-carboxamide (Example 128) or a salt thereof,

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-fluorophenyl)-


30 piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide (Example 130) or a salt thereof, or

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-chloroindole-2-

THE UNIVERSITY OF CHICAGO

carboxamide (Example 131) or a salt thereof,

[28] a production method of a compound of [23], which comprises reacting a compound represented by the formula




5 wherein each symbol is as defined in [23] or a salt thereof
with a reactive derivative of an organic acid of the formula
R²⁹-Alk₃-COOH

wherein each symbol is as defined in [23],

[29] a pharmaceutical composition containing the compound of

10 [23],

[30] a compound represented by the formula



wherein R²⁶ and R²⁷ are the same or different and each is hydrogen atom or C₁₋₆ alkyl group; R³⁰ is hydrogen atom, C₁₋₆

¹⁵ alkyl group or optionally halogenated C₁₋₆ alkyl-carbonyl group;

Alk is C₁₋₆ alkylene group optionally having substituents; Alk₂

and Alk_3 are the same or different and each is a bond or C_{1-6}

alkylene group optionally having substituents; R³¹ is C₆₋₁₀ aryl

group optionally having substituents; X^2 is CH, C-OH or N; P^5

²⁰ and Q⁵ are the same or different and each is C₁₋₆ alkylene

group; Y^6 , Y^7 and Y^8 are the same or different and each is

hydrogen atom, halogen atom, optionally halogenated C₁₋₆ alkyl

group or optionally halogenated C₁₋₆ alkoxy group, or a salt


thereof or a prodrug thereof,

²⁵ [31] the compound of [30] wherein Alk is C₁₋₆ alkylene group

optionally having substituents selected from the group consisting of halogen atom, hydroxy, amino and C₆₋₁₀ aryl; Alk₂ and Alk₃ are the same or different and each is a bond or C₁₋₆ alkylene group optionally having substituents selected from the group consisting of halogen atom, hydroxy, amino and C₆₋₁₀ aryl; R³¹ is C₆₋₁₀ aryl group optionally having substituents selected from the group consisting of halogen atom, C₁₋₆ alkyl, hydroxy, C₁₋₆ alkoxy and C₆₋₁₀ aryl,

[32] N-(2-((2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)amino)-2-oxoethyl)-2,2,2-trifluoro-N-phenylacetamide (Example 51) or a salt thereof, 2-anilino-N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)acetamide (Example 59) or a salt thereof, or 2-(((benzylamino)carbonyl)amino)-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)acetamide (Example 65) or a salt thereof,

[33] a production method of a compound of [30], which comprises reacting a compound represented by the formula



wherein each symbol is as defined in [30], or a salt thereof, with,

(1) when Alk₂ is C₁₋₆ alkylene group optionally having substituents, a reactive derivative of an organic acid compound of the formula

R³¹-Alk₃-NR³⁰-Alk₂-COOH

wherein each symbol is as defined in [30],

(2) when Alk₂ is a bond, a reactive derivative of the formula R³¹-Alk₃-NR³⁰-CO-X or R³¹-Alk₃-NCO


wherein X is leaving group, and other symbols are as defined in

[30],

[34] a pharmaceutical composition containing a compound of [30],

[35] a method for antagonizing melanin-concentrating hormone,
comprising administering, to a mammal, an effective amount of a

5 compound of the formula



wherein Ar¹ and Ar² are each an aromatic group optionally
having substituents,


P and Q are each a divalent aliphatic hydrocarbon group which
10 optionally contains ether oxygen or ether sulfur in a carbon
chain and which optionally has substituents, ,

R¹ and R³ are each (i) hydrogen atom, (ii) acyl group or (iii)
a hydrocarbon group optionally having substituents,

R² and R⁴ are each (i) hydrogen atom, (ii) an alkyl group
15 optionally having substituents or (iii) an alkylcarbonyl group
optionally having substituents, R¹ and R² or R³ and R⁴

optionally form, together with the adjacent nitrogen atom, a
monocyclic or fused nitrogen-containing heterocyclic group
optionally having substituents, and j is 0 or 1, or a salt
20 thereof or a prodrug thereof,

[36] use of a compound of the formula



wherein Ar¹ and Ar² are each an aromatic group optionally

having substituents,


P and Q are each a divalent aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which optionally has substituents, ,

5 R¹ and R³ are each (i) hydrogen atom, (ii) acyl group or (iii) hydrocarbon group optionally having substituents,

R² and R⁴ are each (i) hydrogen atom, (ii) an alkyl group optionally having substituents or (iii) an alkylcarbonyl group optionally having substituents, R¹ and R² or R³ and R⁴

10 optionally form, together with the adjacent nitrogen atom, a monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents, and j is 0 or 1, or a salt thereof or a prodrug thereof, for production of a melanin-concentrating hormone antagonist and

15 [37] a compound represented by the formula



wherein R²⁷ is hydrogen atom, halogen atom, optionally halogenated C₁₋₆ alkyl group or optionally halogenated C₁₋₆ alkoxy group, or a salt thereof.

20

Best Mode for Embodiment the Invention

As the "aromatic group" represented by Ar¹ and Ar², for example, aromatic hydrocarbon group, aromatic heterocyclic group and the like are used, and aromatic hydrocarbon group is particularly preferable.

25

As the "aromatic hydrocarbon group", for example, monocyclic or fused polycyclic aromatic hydrocarbon group

besides carbon atom, 1 to 3 heteroatoms selected from nitrogen atom, oxygen atom and sulfur atom, such as 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furyl, 3-furyl, 2-quinolyl, pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 2-imidazolyl, 3-
 5 pyridazinyl, 3-isothiazolyl, 3-isooxazolyl, 1-isoquinolyl, 1-indolyl, 2-indolyl, 2-benzothiazolyl and the like, and the like are used. Of these, 5 or 6-membered aromatic heterocyclic group containing, besides carbon atom, 1 to 3 (preferably 1) heteroatoms selected from nitrogen atom, oxygen atom and sulfur
 10 atom, such as 2-thienyl, 3-thienyl, 2-pyridyl, 4-pyridyl and the like, and the like are generally used.

As the substituent that the "aromatic group" represented by Ar¹ and Ar² optionally has, for example, halogen atom (e.g., fluorine, chlorine, bromine, iodine and the like); C₁₋₃
 15 alkylenedioxy group (e.g., methylenedioxy, ethylenedioxy and the like); nitro group; cyano group; optionally halogenated C₁₋₆ alkyl group; optionally halogenated C₃₋₆ cycloalkyl group; optionally halogenated C₁₋₆ alkoxy group; optionally halogenated C₁₋₆ alkylthio group; hydroxy group; amino group; mono-C₁₋₆
 20 alkylamino group (e.g., methylamino, ethylamino and the like); di-C₁₋₆ alkylamino group (e.g., dimethylamino, diethylamino and the like); optionally halogenated C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like); formyl group; C₁₋₆ alkyl-carbonyl group (e.g., acetyl, propionyl,
 25 butyryl and the like) optionally substituted by halogen atom or C₁₋₆ alkoxy-carbonyl group (e.g., methoxycarbonyl, ethoxycarbonyl and the like); C₁₋₆ alkyl-carboxyloxy group (e.g., acetyloxy, propionyloxy, butyryloxy and the like); carboxyl group; C₁₋₆ alkoxy-carbonyl group (e.g., methoxycarbonyl,
 30 ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and the like); carbamoyl group; mono-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl, ethylcarbamoyl and the like) optionally substituted by C₁₋₆ alkoxy-carbonyl group; di-C₁₋₆ alkyl-

The aforementioned C₆₋₁₀ aryl group and C₆₋₁₀ aryloxy group optionally have 1 to 4 substituents selected from halogen atom, C₁₋₃ alkylenedioxy, nitro, cyano, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₃₋₆ cycloalkyl, optionally halogenated C₁₋₆ alkoxy, optionally halogenated C₁₋₆ alkylthio, hydroxy, amino, mono-C₁₋₆ alkylamino, di-C₁₋₆ alkylamino, C₁₋₆ alkyl-carbonylamino, formyl, C₁₋₆ alkyl-carbonyl, C₁₋₆ alkyl-carbonyloxy, carboxyl, C₁₋₆ alkoxy-carbonyl, carbamoyl, mono-C₁₋₆ alkyl-carbamoyl, di-C₁₋₆ alkyl-carbamoyl, sulfo, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl and the like.

45

As the "optionally halogenated C₁₋₆ alkyl group" used in the present specification, for example, C₁₋₆ alkyl group optionally having 1 to 5 halogen atom(s) (e.g., fluorine, chlorine, bromine, iodine and the like), such as methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromomethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, propyl, 3,3,3-trifluoropropyl, isopropyl, butyl, 4,4,4-trifluorobutyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl, 6,6,6-trifluorohexyl and the like, and the like are used.

As the "optionally halogenated C₃₋₆ cycloalkyl group" used in the present specification, for example, C₃₋₆ cycloalkyl group optionally having 1 to 4 halogen atom(s) (e.g., fluorine, chlorine, bromine, iodine and the like), such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 2,2,3,3-tetrafluorocyclopentyl, 4-chlorocyclohexyl and the like, and the like are used.

As the "optionally halogenated C₁₋₆ alkoxy group" used in the present specification, for example, C₁₋₆ alkoxy group optionally having 1 to 3 halogen atom(s) (e.g., fluorine, chlorine, bromine, iodine and the like), such as methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-trifluoroethoxy, propoxy, isopropoxy, butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy and the like, and the like are used.

As the "optionally halogenated C₁₋₆ alkylthio group" used in the present specification, for example, C₁₋₆ alkylthio group optionally having 1 to 3 halogen atom(s) (e.g., fluorine, chlorine, bromine, iodine and the like), such as methylthio, difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio, hexylthio and the like, and the like are used.

As the "hydrocarbon group" represented by R¹ and R³, for

the like), C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like), formyl group, C₁₋₆ alkyl-carbonyl group (e.g., acetyl, propionyl, butyryl and the like), C₁₋₆ alkyl-carbonyloxy group (e.g., acetyloxy, propionyloxy, butyryloxy and the like), carboxyl group, C₁₋₆ alkoxy-carbonyl group (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and the like), carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl, ethylcarbamoyl and the like), di-C₁₋₆ alkyl-carbamoyl group (e.g., dimethylcarbamoyl, diethylcarbamoyl and the like), sulfo group, C₁₋₆ alkylsulfonyl group (e.g., methylsulfonyl, ethylsulfonyl and the like), C₁₋₆ alkylsulfinyl group (e.g., methylsulfinyl, ethylsulfinyl and the like), C₆₋₁₀ aryl group (e.g., phenyl, naphthyl and the like), C₆₋₁₀ aryloxy group (e.g., phenyloxy, naphthyloxy and the like), 5 to 7-membered heterocyclic ring [e.g., 5 to 7-membered heterocyclic ring containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom (e.g., 1-, 2- or 3-pyrrolidinyl, 2- or 4-imidazolidinyl, 2-, 3- or 4-pyrazolidinyl, 1-, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl, morpholinyl, 2-thienyl, 3-thienyl, 2-pyridyl, 4-pyridyl, 2-furyl, 3-furyl, pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 2-imidazolyl, 3-pyridazinyl, 3-isothiazolyl, 3-isooxazolyl and the like) or fused ring group thereof (e.g., fused ring group with benzene ring and the like)], di-C₁₋₆ alkyl-carbonylamino group, sulfamoyl group, C₁₋₆ alkoxy-carbonylamino group (e.g., methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, butoxycarbonylamino, tert-butoxycarbonylamino and the like), C₇₋₁₆ aralkyloxy-carbonylamino group (e.g., benzyloxycarbonylamino and the like), C₇₋₁₆ aralkyloxy group (e.g., benzyloxy and the like), C₆₋₁₀ aryl-carbonyl group (e.g., benzoyl and the like), C₁₋₆ alkyl-carbonyloxy group (e.g., acetoxy and the like), C₆₋₁₀ aryl-

carbonylamino group (e.g., benzoylamino and the like), C₆₋₁₀ aryl-carbamoyl group (e.g., phenylcarbamoyl and the like) and the like are used.


The above-mentioned C₆₋₁₀ aryl group, C₆₋₁₀ aryloxy group
 5 and 5 to 7-membered heterocyclic ring may have 1 to 4
 substituents selected from halogen atom, C₁₋₃ alkylenedioxy,
 nitro, cyano, optionally halogenated C₁₋₆ alkyl, optionally
 halogenated C₃₋₆ cycloalkyl, optionally halogenated C₁₋₆ alkoxy,
 optionally halogenated C₁₋₆ alkylthio, hydroxy, amino, mono-C₁₋₆
 10 alkylamino, di-C₁₋₆ alkylamino, C₁₋₆ alkyl-carbonylamino, formyl,
 C₁₋₆ alkyl-carbonyl, C₁₋₆ alkyl-carbonyloxy, carboxyl, C₁₋₆
 alkoxy-carbonyl, carbamoyl, mono-C₁₋₆ alkyl-carbamoyl, di-C₁₋₆
 alkyl-carbamoyl, sulfo, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl
 and the like.

15 The "hydrocarbon group" represented by R¹ and R³ may have,
 for example, 1 to 5, preferably 1 to 3, suitable substituent(s)
 selected from the above-mentioned substituents at substitutable
 position(s) of the hydrocarbon group. When the number of the
 substituent is two or more, these substituents may be the same
 20 or different.


As the "acyl group" represented by R¹ and R³, for example,
 a group represented by -CO-R^a, -CONR^aR^b, -SO-R^a, -SO₂-R^a, -
 CONR^aR^b, -COO-R^a, -(C=S)O-R^a, -(C=S)NR^aR^b, -SONR^aR^b, -SO₂NR^aR^b, -
 SO-O-R^a, -SO₂-O-R^a (R^a is hydrogen atom, carboxyl group,
 25 hydrocarbon group optionally having substituents, heterocyclic
 group optionally having substituents or C₁₋₆ alkoxy-carbonyl
 group, and R^b is hydrogen atom or C₁₋₆ alkyl group) and the like
 is used. Particularly, -CO-R^a, -CONH-R^a and the like are
 preferable.

30 As the "acyl group" represented by R³,
 (xvi) a group represented by the formula


(xvii) a group represented by the formula



(xviii) a group represented by the formula



(xix) a group represented by the formula



50

$$-\text{CO}-\text{Alk}-\text{NR}^{27}-\text{CO}-\text{Alk}_2-\text{NR}^{30}-\text{Alk}_3-\text{R}^{31}$$

¹⁰ [Alk, R²⁷, Alk₂, Alk₃ are as defined above; R³⁰ is hydrogen atom, C₁₋₆ alkyl group (e.g., methyl) or optionally halogenated C₁₋₆ alkyl-carbonyl group (e.g., trifluoromethylcarbonyl); R³¹ is C₆₋₁₀ aryl group (e.g., phenyl) optionally having substituents (e.g., halogen atom (e.g., fluorine, chlorine), C₁₋₆ alkyl group

¹⁵ (e.g., methyl), hydroxy group, C₁₋₆ alkoxy group, C₆₋₁₀ aryl group (e.g., phenyl) and the like)];

$$-\text{CO}-\text{Alk}-\text{NR}^{27}-\text{CO}-\text{Alk}_2-\text{NR}^{32}-\text{CO}-\text{O}-\text{Alk}_3-\text{R}^{31}$$

[Alk, R^{27} , Alk₂, Alk₃ and R^{31} are as defined above; and R^{32} is
²⁰ the same as the aforementioned R^{27}];

$$-\text{CO}-\text{Alk}-\text{CO}-\text{NR}^{27}-\text{Alk}_2-\text{R}^{31}$$

[Alk, R^{27} , Alk₂ and R^{31} are as defined above]:

$$^{25} \quad -\text{CO}-\text{Alk}-\text{O}-\text{CO}-\text{O}-\text{Alk}_2-\text{R}^{31}$$

[Alk, Alk₂ and R³¹ are as defined above] and the like are mentioned.

As the "hydrocarbon group optionally having substituents
"represented by the aforementioned R^a, those similar to the
30 aforementioned "hydrocarbon group optionally having
substituents "represented by R¹ and R³ are used.

As the "heterocyclic group" represented by R^a , for example, 5 to 10-membered (monocyclic or bicyclic) heterocyclic

As the substituent of the "alkylcarbonyl group", for example, those similar to the substituent that the aforementioned "hydrocarbon group" represented by R^1 and R^3 optionally has and the like are used.

As the "divalent aliphatic hydrocarbon group" of the
15 "divalent aliphatic hydrocarbon group which optionally contains
ether oxygen or ether sulfur in a carbon chain" represented by
P and Q, for example, a divalent group obtained by respectively
removing one hydrogen atom bonded to the same or different
carbon atom of saturated or unsaturated aliphatic hydrocarbon
20 and the like are exemplified, with preference given to those
having not more than 6 carbon atoms. Examples thereof include
(i) alkylene group (e.g., $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-CH_2-$,
 $CH(CH_3)-CH_2-$, $-(CH_2)_5-$, $-CH_2-CH(CH_3)-CH_2-CH_2-$, $-(CH_2)_6-$ and the
like),

(iii) alkynylene group (e.g., $-C\equiv C-$, $-CH_2-C\equiv C-$, $-CH_2-C\equiv C-CH_2-$ and the like) and the like. Preferably, C_{1-6} alkylene group (e.g., methylene, ethylene, propylene, trimethylene,

53

group is used.

The "divalent aliphatic hydrocarbon group" may contain ether oxygen or ether sulfur in a carbon chain, which is optionally substituted by oxo group or thioxo group.

To be specific, for example, $-\text{CH}_2-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{CH}_2-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{O}-\text{CH}_2-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{S}-\text{CH}_2-$, $-\text{CH}_2-\text{CH}_2-\text{S}-\text{CH}_2-$, $-\text{CH}_2-\text{S}-\text{CH}_2-\text{S}-\text{CH}_2-$ and the like are used.

As the "monocyclic or fused nitrogen-containing heterocyclic group" of the "monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents" formed by R^1 and R^2 , and R^3 and R^4 , together with the adjacent nitrogen atom, for example, monocyclic or fused 3 to 9-membered, preferably 5 to 7-membered, nitrogen-containing heterocyclic group optionally containing, besides the nitrogen atom of the bond moiety, 1 or 2 kinds of preferably 1 to 3 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom (e.g., pyrrolidyl, piperidyl, piperazyl and the like) and the like are used.

As the substituent that the "monocyclic or fused nitrogen-containing heterocyclic group" optionally has, for example, those similar to the substituent that the aforementioned Ar^1 and Ar^2 optionally have and the like are used.


As the "monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents", for example (i) a group represented by the formula




wherein ring A is a 4 to 8-membered ring optionally substituted by 1 or 2 hydroxy group(s) or oxo group(s); V is a group represented by the formula $>\text{O}$, $>\text{C}=\text{O}$, $>\text{C}(\text{W})-\text{W}^a$ or $>\text{N}-\text{W}$ (W is hydrogen atom, hydrocarbon group optionally having substituents

or heterocyclic group optionally having substituents, W^a is hydrogen atom, hydroxy group or C₁₋₆ alkyl group),

(ii) a group represented by the formula




5 wherein ring B is a monocyclic or bicyclic 4 to 12-membered ring optionally substituted by 1 or 2 oxo group(s) or 1 to 5 C₁₋₆ alkyl group(s), ring D is a 4 to 12-membered aromatic ring optionally having substituents, preferably a group represented by the formula




wherein ring B is monocyclic or bicyclic 4 to 12-membered ring optionally substituted by 1 or 2 oxo group(s) or 1 to 5 C₁₋₆ alkyl group(s), D^a is a halogen atom (e.g., fluorine, chlorine, bromine, iodine and the like), a C₁₋₆ alkyl group (e.g., methyl, ethyl, propyl, isopropyl and the like), a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, propoxy, isopropoxy and the like), a C₁₋₃ alkylenedioxy group (e.g., methylenedioxy, ethylenedioxy and the like), a nitro group, an amino group or a C₁₋₆ alkyl-carbonyl group (e.g., acetyl, propionyl and the like),

²⁰ (iii) a group represented by the formula




wherein ring E is a 4 to 12-membered aromatic ring optionally having substituents, X is $-CH_2-$, $-CO-$ or $-CH(OH)-$, Y is $-CH_2-$, $-O-$ or $-NW^b-$ (W^b is hydrogen atom or C_{1-6} alkyl group optionally having substituents), k and m are each an integer of 0 to 4,

As the "4 to 8-membered ring optionally substituted by 1 or 2 hydroxy group(s) or oxo group(s)" represented by A, for example, a group represented by the formula



(V is as defined above), preferably the formula



(V is as defined above, G is halogen atom (e.g., fluorine, chlorine and the like), C₁₋₆ alkyl group (e.g., methyl, ethyl, propyl, isopropyl and the like), optionally halogenated C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, propoxy, isopropoxy, trifluoromethyl and the like), hydrogen atom, cyano group and the like) and the like are generally used.

G is preferably halogen atom such as fluorine, chlorine
²⁰ and the like; C₁₋₆ alkyl group such as methyl, ethyl, propyl,

isopropyl and the like; C₁₋₆ alkoxy group such as methoxy, ethoxy, propoxy, isopropoxy and the like, and the like.

As the "hydrocarbon group optionally having substituents" represented by W, for example, those similar to the

5 "hydrocarbon group optionally having substituents" represented by the aforementioned R^1 and R^3 , and the like are used, and C_{6-14} aryl group (e.g., phenyl and the like), C_{7-16} aralkyl group (e.g., benzyl and the like) and the like are particularly preferable.

10 As the substituent that this hydrocarbon group optionally has, for example, those similar to the substituents that the aforementioned "hydrocarbon group" represented by R^1 and R^3 optionally have and the like are used.

As the "heterocyclic group" represented by W, for example,
a 5 to 10-membered (monocyclic or bicyclic) heterocyclic group
containing, besides carbon atom, 1 or 2 kinds of preferably 1
to 4 heteroatom(s) selected from nitrogen atom, oxygen atom and
sulfur atom and the like are used. Specific examples include
1-, 2- or 3-pyrrolidinyl, 2- or 4-imidazolidinyl, 2-, 3- or 4-
pyrazolidinyl, 1-, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl,
morpholinyl, 2-thienyl, 3-thienyl, 2-pyridyl, 4-pyridyl, 2-
furyl, 3-furyl, 4-quinolyl, 8-quinolyl, 4-isoquinolyl,
pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 2-imidazolyl, 3-
pyridazinyl, 3-isothiazolyl, 3-isooxazolyl, 1-indolyl, 2-
isoindolyl and the like, with preference given to aromatic ones.
Particularly, for example, a 5 or 6-membered aromatic
heterocyclic group containing, besides carbon atom, 1 to 3
heteroatom(s) selected from nitrogen atom, oxygen atom and
sulfur atom (e.g., 2-thienyl, 3-thienyl, 2-pyridyl, 4-pyridyl
and the like) and the like are preferable.

As the substituent that the "heterocyclic group" optionally has, for example, those similar to the substituents that the aforementioned "an aromatic group optionally having

[illegible]

5

alkyl group(s)" represented by B, for example



and the like are used.

15

⁵ benzene ring is preferable.


¹⁰ in similar number.

tert-butyl, pentyl, hexyl and the like are used.

15 has, for example, 1 to 3 halogen atom(s), nitro group, cyano
group, optionally halogenated C₃₋₆ cycloalkyl group, optionally
halogenated C₁₋₆ alkoxy group, optionally halogenated C₁₋₆
alkylthio group, hydroxy group, amino group, mono-C₁₋₆
alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkyl-
20 carbonylamino group (e.g., acetylamino, propionylamino,
butyrylamino and the like), carboxyl group, C₁₋₆ alkoxy-carbonyl
group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆
alkyl-carbamoyl group, sulfo group, C₁₋₆ alkylsulfonyl group,
C₁₋₆ alkylsulfinyl group, C₆₋₁₀ aryl group, C₆₋₁₀ aryloxy group

and 5 or 6-membered heterocyclic group (e.g., thienyl, furyl, pyridyl and the like) and the like are used.


As the "nitrogen-containing aromatic heterocyclic group", for example, 5 to 10-membered (monocyclic or bicyclic) aromatic heterocyclic group containing one nitrogen atom besides carbon atom, and further, optionally containing 1 or 2 kinds of preferably 1 to 3 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom and the like are used. Specific examples include



and the like. When a counter ion is necessary, for example, halogen ion (e.g., chlorine ion, bromine ion, iodine ion and the like) and the like are used.


As the substituent that the "nitrogen-containing aromatic heterocyclic group" optionally has, for example, those similar to the substituents that the aforementioned "aromatic group" represented by Ar^1 and Ar^2 optionally has and the like are used.

Of the aforementioned examples, a monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents formed by R^1 and R^2 or R^3 and R^4 , together with the adjacent nitrogen atom, for example, (i) a group represented by the formula



wherein ring E^a is a benzene ring optionally having substituents, which is preferably, for example, benzene ring optionally having 1 to 4 substituents selected from the group consisting of halogen atom, C_{1-3} alkylendioxy group, nitro

(ii) a group represented by the formula



61

substituent(s) selected from the group consisting of halogen atom, C₁₋₃ alkylendioxy group, nitro group, cyano group, optionally halogenated C₁₋₆ alkyl group, optionally halogenated C₃₋₆ cycloalkyl group, optionally halogenated C₁₋₆ alkoxy group, 5 optionally halogenated C₁₋₆ alkylthio group, hydroxy group, amino group, mono-C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkyl-carbonylamino group, formyl group, C₁₋₆ alkyl-carbonyl group, C₁₋₆ alkyl-carbonyloxy group, carboxyl group, C₁₋₆ alkoxy-carbonyl group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, 10 di-C₁₋₆ alkyl-carbamoyl group, sulfo group, C₁₋₆ alkylsulfonyl group, C₁₋₆ alkylsulfinyl group, C₆₋₁₀ aryl group and C₆₋₁₀ aryloxy group; W^a is hydrogen atom, hydroxy group or C₁₋₆ alkyl group) and the like are preferable.

As W, for example, C₆₋₁₄ aryl group and C₇₋₁₆ aralkyl group, 15 each of which optionally has 1 or 2 substituent(s) selected from the group consisting of halogen atom, cyano group, optionally halogenated C₁₋₆ alkyl group, optionally halogenated C₁₋₆ alkoxy group and the like, are preferable. Particularly, phenyl group optionally substituted by halogen atom or 20 optionally halogenated C₁₋₆ alkoxy group, and the like are preferable.

Of the aforementioned examples, as the aromatic group represented by Ar¹ and Ar², C₆₋₁₀ aryl group (e.g., phenyl group and the like), 5 to 10-membered (monocyclic or bicyclic) 25 aromatic heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom (particularly, thienyl group) and the like are preferable, with particular preference given to C₆₋₁₀ aryl group (e.g., phenyl group and the like).

30 As Ar¹ and Ar², for example, (i) C₆₋₁₄ aryl group (particularly, phenyl group) or (ii) 5 to 10-membered (monocyclic or bicyclic) aromatic heterocyclic group (particularly thienyl group) containing, besides carbon atom, 1

to 3 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom, which optionally has 1 to 3 substituent(s) selected from the group consisting of halogen atom, methylenedioxy group, nitro group, cyano group, optionally
5 halogenated C₁₋₆ alkyl group, C₃₋₆ cycloalkyl group, optionally halogenated C₁₋₆ alkoxy group, optionally halogenated C₁₋₆ alkylthio group, hydroxy group, amino group, mono-C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, formyl group, C₁₋₆ alkyl-carbonyl group, C₁₋₆ alkyl-carbonyloxy group, carboxyl
10 group, C₁₋₆ alkoxy-carbonyl group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆ alkyl-carbamoyl group, sulfo group, C₁₋₆ alkylsulfonyl group, phenyl group and phenyloxy group and the like are preferable. Of these, (i) phenyl group optionally having substituents selected from halogen atom, C₁₋₆
15 alkoxy group and optionally halogenated C₁₋₆ alkyl group or (ii) 5 or 6-membered aromatic heterocyclic group (particularly, thienyl group) containing, besides carbon atom, 1 to 3 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom are preferable.

20 Ar¹ and Ar² are more preferably phenyl group optionally having substituents selected from halogen atom, C₁₋₆ alkoxy group and optionally halogenated C₁₋₆ alkyl group, particularly preferably phenyl group optionally substituted by halogen atom or C₁₋₆ alkoxy group.

25 As P and Q, C₁₋₆ alkylene group or C₂₋₆ alkenylene group, which optionally contain ether oxygen or ether sulfur in a carbon chain, and the like are preferable. Of these, C₁₋₆ alkylene group or C₂₋₆ alkenylene group is preferable and particularly, C₁₋₆ alkylene group (e.g., methylene, ethylene,
30 trimethylene, tetramethylene and the like) is generally used.

As P, C₃₋₅ alkylene group (e.g., trimethylene, tetramethylene and the like) and the like are preferable, and particularly, trimethylene and tetramethylene are preferable.

As Q, C₁₋₃ alkylene group (e.g., methylene, ethylene, trimethylene) and the like are preferable, and particularly, methylene is preferable.

As the acyl group represented by R¹, a group represented
 5 by -CO-R^a or -CONH-R^a (R^a is as defined above), and the like are preferable.

As the "hydrocarbon group optionally having substituents" represented by R¹ and R^a, for example, (i) C₁₋₆ alkyl group, (ii) C₂₋₆ alkenyl group, (iii) C₂₋₆ alkynyl group, (iv) C₃₋₆
 10 cycloalkyl group optionally fused with benzene ring, (v) C₆₋₁₄ aryl group or (vi) C₇₋₁₆ aralkyl group and the like, each of which optionally has 1 to 3 substituent(s) selected from halogen atom, C₁₋₃ alkylenedioxy group, nitro group, cyano group, optionally halogenated C₁₋₆ alkyl group, optionally halogenated
 15 C₃₋₆ cycloalkyl group, optionally halogenated C₁₋₆ alkoxy group, optionally halogenated C₁₋₆ alkylthio group, hydroxy group, amino group, mono-C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like), formyl group, C₁₋₆
 20 alkyl-carbonyl group, C₁₋₆ alkyl-carbonyloxy group, carboxyl group, C₁₋₆ alkoxy-carbonyl group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆ alkyl-carbamoyl group, sulfo group, C₁₋₆ alkylsulfonyl group, C₁₋₆ alkylsulfinyl group, C₆₋₁₀ aryl group, C₆₋₁₀ aryloxy group and 5 to 7-membered heterocyclic
 25 ring (e.g., thienyl, furyl, pyridyl and the like) is preferable.

Of the above-mentioned examples, as R¹, (i) C₁₋₆ alkyl group optionally having 5 or 6-membered nitrogen-containing heterocyclic group (e.g., pyridyl group), (ii) C₇₋₁₆ aralkyl group optionally having nitro, amino or C₁₋₆ alkoxy-carbonyl
 30 (particularly benzyl group), (iii) cyclohexyl group fused with benzene ring optionally having C₁₋₆ alkoxy and the like are preferable.

As the "alkyl group optionally having substituents"


As the "monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents" formed by R¹ and R² together with the adjacent nitrogen, for example,

(i) a group represented by the formula




wherein ring A¹ is 4 to 8-membered ring optionally substituted by hydroxy or oxo, V¹ is a group represented by the formula >O,
 5 >C(W¹)-W^{a1} or >N-W¹ (W¹ is (a) hydrogen atom, (b) C₆₋₁₄ aryl group optionally having 1 or 2 substituent(s) selected from the group consisting of halogen atom, optionally halogenated C₁₋₆ alkyl group and optionally halogenated C₁₋₆ alkoxy group, (c) C₁₋₆ alkyl group optionally having 1 or 2 C₆₋₁₀ aryl group(s) or (d)
 10 pyridyl group, W^{a1} is hydrogen atom, hydroxy group or C₁₋₆ alkyl group),

(ii) a group represented by the formula



wherein ring B¹ is monocyclic or bicyclic 5 to 10-membered ring
 15 optionally substituted by oxo group or 1 or 2 C₁₋₆ alkyl group(s), ring D¹ is benzene ring optionally having 1 or 2 substituent(s) selected from the group consisting of C₁₋₆ alkyl group, C₁₋₆ alkoxy group and C₁₋₆ alkyl-carbonyl group, and
 (iii) a group represented by the formula



wherein ring E¹ is benzene ring optionally having 1 to 3 substituent(s) selected from C₁₋₃ alkylenedioxy group, nitro group, C₁₋₆ alkoxy group, amino group, C₁₋₆ alkyl-carbonylamino group and C₁₋₆ alkoxy-carbonyl group, X¹ is -CH₂- or -CO-, Y¹ is
 25 -CH₂- or -O-) and the like are preferable.

As the acyl group represented by R³, a group represented

As the "hydrocarbon group optionally having substituents" represented by R^a or R^3 , for example, (i) C_{1-6} alkyl group, (ii) C_{2-6} alkenyl group, (iii) C_{2-6} alkynyl group, (iv) C_{3-6} cycloalkyl group optionally fused with benzene ring, (v) C_{6-14} aryl group or (vi) C_{7-16} aralkyl group, each optionally having 1 to 3 substituent(s) selected from, for example, halogen atom, C_{1-3} alkylenedioxy group, nitro group, cyano group, optionally halogenated C_{1-6} alkyl group, optionally halogenated C_{3-6} cycloalkyl group, optionally halogenated C_{1-6} alkoxy group, optionally halogenated C_{1-6} alkylthio group, hydroxy group, amino group, mono- C_{1-6} alkylamino group, di- C_{1-6} alkylamino group, C_{1-6} alkyl-carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like), formyl group, C_{1-6} alkyl-carbonyl group, C_{1-6} alkyl-carbonyloxy group, carboxyl group, C_{1-6} alkoxy-carbonyl group, carbamoyl group, mono- C_{1-6} alkyl-carbamoyl group, di- C_{1-6} alkyl-carbamoyl group, sulfo group, C_{1-6} alkylsulfonyl group, C_{1-6} alkylsulfinyl group, C_{6-10} aryl group, C_{6-10} aryloxy group and 5 to 7-membered heterocyclic ring (e.g., thienyl, furyl, pyridyl and the like) and the like are preferable.

As the acyl group represented by R^3 , for example, $-\text{CO}-R^a$ (R^a is as defined above) and the like are preferable, and particularly, $-\text{CO}-R^c$ (R^c is (1) C_{1-6} alkyl group, (2) C_{2-6} alkenyl group, (3) C_{2-6} alkynyl group, (4) C_{3-6} cycloalkyl group, (5) C_{6-14} aryl group or (6) C_{7-16} aralkyl group, each optionally having 1 to 5 substituent(s) selected from the group consisting of halogen atom, C_{1-3} alkylenedioxy group, nitro group, cyano group, optionally halogenated C_{1-6} alkyl group, optionally halogenated C_{3-6} cycloalkyl group, optionally halogenated C_{1-6} alkoxy group, optionally halogenated C_{1-6} alkylthio group, hydroxy group, amino group, mono- C_{1-6} alkylamino group, di- C_{1-6}

alkylamino group, C₁₋₆ alkyl-carbonylamino group, formyl group, C₁₋₆ alkyl-carbonyl group, C₁₋₆ alkyl-carbonyloxy group, carboxyl group, C₁₋₆ alkoxy-carbonyl group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆ alkyl-carbamoyl group, sulfo
 5 group, C₁₋₆ alkylsulfonyl group, C₁₋₆ alkylsulfinyl group, C₆₋₁₀ aryl group, C₆₋₁₀ aryloxy group and 5 to 7-membered heterocyclic ring (e.g., thienyl, furyl, pyridyl and the like) is preferable.


As R^c, for example, C₇₋₁₆ aralkyl group optionally having 1 to 3 substituent(s) selected from halogen atom, C₁₋₃
 10 alkylenedioxy group, nitro group, cyano group, optionally halogenated C₁₋₆ alkyl group, optionally halogenated C₃₋₆ cycloalkyl group, optionally halogenated C₁₋₆ alkoxy group, optionally halogenated C₁₋₆ alkylthio group, hydroxy group, amino group, mono-C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group,
 15 C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like), formyl group, C₁₋₆ alkyl-carbonyl group, C₁₋₆ alkyl-carbonyloxy group, carboxyl group, C₁₋₆ alkoxy-carbonyl group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆ alkyl-carbamoyl group, sulfo
 20 group, C₁₋₆ alkylsulfonyl group, C₁₋₆ alkylsulfinyl group, C₆₋₁₀ aryl group and C₆₋₁₀ aryloxy group, and the like are preferable. Particularly, C₇₋₁₆ aralkyl group optionally substituted by 1 to 3 halogen atom(s) or C₁₋₆ alkoxy group is preferable.

As R³,

- 25 (i) hydrogen atom;
 (ii) a group represented by the formula -CO-R⁵ (R⁵ is (a) hydrogen atom, (b) carboxyl group, (c) C₁₋₆ alkyl group (particularly, C₁₋₃ alkyl group such as methyl, ethyl and the like), (d) C₅₋₆ cycloalkyl group (e.g., cyclopentyl and
 30 cyclohexyl) optionally having C₁₋₆ alkoxy (e.g., methoxy) and fused with benzene ring, (e) 5 or 6-membered aromatic heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen

(iii) a group represented by the formula $\text{-CO-Alk}_0\text{-R}^6$ [Alk_0 is C_{1-6} alkylene group optionally having hydroxy group, preferably a group represented by the formula $(\text{CH}_2)_{r^1}$ (r^1 is an integer of 1 to 3); R^6 is (a) C_{6-10} aryl group (e.g., phenyl) optionally having 1 or 2 substituent(s) selected from the group consisting of halogen atom, optionally halogenated C_{1-6} alkyl (e.g., trifluoromethyl), nitro, C_{1-6} alkoxy (e.g., methoxy, ethoxy), C_{1-3} alkylendioxy (e.g., methylenedioxy) and C_{6-10} aryl group (e.g., phenyl), (b) C_{6-10} aryloxy group (e.g., phenyloxy), (c) 5 or 6-membered aromatic heterocyclic group (e.g., pyridyl) containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom, (d) C_{1-6} alkyl-carbonyl group (e.g., acetyl), (e) carboxyl group, (f) C_{1-6} alkoxy-carbonyl group (e.g., C_{1-3} alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like), (g) amino optionally having 1 or 2 substituent(s) selected from the group consisting of C_{1-6} alkyl (e.g., C_{1-3} alkyl such as methyl, ethyl and the like) and C_{1-6} alkyl-carbonyl (e.g., acetyl), (h) 5 to 7-membered heterocyclic ring optionally having hydroxy, (i) C_{7-16} aralkyloxy group (e.g., benzyloxy), (j) C_{6-10} aryl-carbonyl group (e.g., benzoyl), (k) C_{1-6} alkyl-carbonyloxy group (e.g., acetoxy)];

(iv) a group represented by the formula




wherein Qa is a group represented by the formula $-(CH_2)_s-$ (s is



an integer of 1 to 3) or $-(CH_2)_t-CH=$ (t is an integer of 0 to 2), R^7 is hydrogen atom or C_{1-6} alkoxy-carbonyl group (e.g., C_{1-3} alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl and the like);

⁵ (v) a group represented the formula




wherein R⁸ is (a) hydrogen atom, (b) C₁₋₆ alkyl group (e.g., C₁₋₃ alkyl group such as methyl, ethyl, propyl and the like) optionally having substituents selected from the group consisting of C₁₋₆ alkoxy-carbonyl (e.g., C₁₋₃ alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like), morpholino and mono- or di-C₁₋₆ alkylamino (e.g., methylamino, ethylamino, dimethylamino, diethylamino), (c) C₁₋₆ alkoxy-carbonyl group (e.g., C₁₋₃ alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl and the like), (d) a group represented by the formula -CO-R^d (R^d is C₆₋₁₀ aryl group (e.g., phenyl, naphthyl) optionally having halogen atom (e.g., chlorine) or 5 or 6-membered heterocyclic group (e.g., pyridyl) containing, besides carbon atom, 1 or 2 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom), (e) a group represented by the formula -CO-(CH₂)_{r¹}-R^e (r¹ is an integer of 1 to 3, R^e is C₁₋₆ alkoxy-carbonyl group (e.g., C₁₋₃ alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl and the like) or 5 or 6-membered heterocyclic group (e.g., pyridyl and the like) containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom) or (f) a group represented by -CONH-R^f (R^f is C₁₋₆ alkyl group (e.g., C₁₋₃ alkyl group such as methyl, ethyl and the like) or C₆₋₁₄ aryl group (e.g., phenyl, naphthyl and the like));

(vi) a group represented by the formula $-\text{COOR}^9$ (R^9 is optionally halogenated C_{1-6} alkyl group (e.g., methyl, ethyl,

trifluoromethyl));

(vii) a group represented by the formula




wherein R^{10} is hydrogen atom, C_{1-6} alkoxy-carbonyl group (e.g.,

⁵ C_{1-3} alkoxy-carbonyl group such as methoxycarbonyl, ethoxycarbonyl and the like), mono- or di- C_{1-6} alkyl-carbamoyl group (e.g., methylcarbamoyl, ethylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl), optionally halogenated nicotinoyl group or optionally halogenated isonicotinoyl group;

¹⁰ (viii) a group represented by the formula $-\text{CONR}^{11}-\text{R}^{12}$ (R^{11} is hydrogen atom or C_{1-6} alkyl group (e.g., C_{1-3} alkyl group such as methyl, ethyl and the like), R^{12} is C_{1-6} alkyl group (e.g., C_{1-3} alkyl group such as methyl, ethyl, propyl and the like) optionally having substituents selected from the group

¹⁵ consisting of (a) hydroxy, (b) amino, (c) mono- or di- C_{1-6} alkyl-amino (e.g., methylamino, ethylamino, dimethylamino, diethylamino), (d) C_{1-6} alkyl-carbonyl (e.g., C_{1-3} alkyl-carbonyl such as acetyl, ethylcarbonyl and the like), (e) C_{1-6} alkoxy-carbonyl (e.g., C_{1-3} alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like), (f) C_{1-6} alkyl-carbonyloxy (e.g., C_{1-3} alkyl-carbonyloxy such as acetyloxy, ethylcarbonyloxy and the like), (g) sulfamoyl, (h) 5 to 7-membered heterocyclic group optionally substituted by oxo and (i) C_{6-14} aryl (e.g., phenyl));


²⁵ (ix) a group represented by the formula



wherein R^{13} is (a) hydrogen atom, (b) C_{1-6} alkyl group (e.g., C_{1-3} alkyl group such as methyl, ethyl, propyl and the like) optionally having substituents selected from the group


³⁰ consisting of a hydroxy and C_{1-6} alkoxy-carbonyl (e.g., C_{1-3} alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the

like), (c) C₇₋₁₆ aralkyl group (e.g., benzyl), (d) C₁₋₆ alkyl-carbonyl group (e.g., C₁₋₃ alkyl-carbonyl group such as acetyl, ethylcarbonyl and the like) optionally having substituents selected from the group consisting of a halogen atom (e.g.,
 5 fluorine, chlorine) and C₁₋₆ alkoxy-carbonyl (e.g., C₁₋₃ alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like) or (e) C₁₋₆ alkyl-carbamoyl group (e.g., C₁₋₃ alkyl-carbamoyl group such as methylcarbamoyl, ethylcarbamoyl and the like) optionally having C₁₋₆ alkoxy-carbonyl (e.g., C₁₋₃ alkoxy-
 10 carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like);
 (x) a group represented by the formula




wherein R¹⁴ is C₁₋₆ alkyl group (e.g., C₁₋₃ alkyl group such as methyl, ethyl, propyl and the like) or C₇₋₁₆ aralkyl group (e.g.,
 15 benzyl);

(xi) a group represented by the formula



wherein ring F is 5 to 7-membered non-aromatic heterocyclic group (particularly piperidyl) optionally fused with benzene
 20 ring, R¹⁵ is hydrogen atom, C₁₋₆ alkoxy-carbonylamino group (e.g., C₁₋₃ alkoxy-carbonylamino group such as methoxycarbonylamino, ethoxycarbonylamino and the like) or optionally halogenated C₁₋₆ alkyl-carbonylamino group (e.g., optionally halogenated C₁₋₃ alkoxy-carbonylamino group such as
 25 methylcarbonylamino, ethylcarbonylamino, trifluoromethylcarbonylamino and the like);


(xii) a group represented by the formula



wherein R¹⁶ is (a) C₁₋₆ alkyl group (e.g., C₁₋₃ alkyl group such


30

(xvi) a group represented by the formula




wherein the symbols are as defined above;

(xvii) a group represented by the formula




wherein the symbols are as defined above;

5 (xviii) a group represented by the formula




wherein the symbols are as defined above;

(xix) a group represented by the formula




10 wherein the symbols are as defined above;

(xx) a group represented by the formula




wherein the symbols are as defined above;

(xxi) a group represented by the formula




wherein the symbols are as defined above;

(xxii) a group represented by the formula



wherein the symbols are as defined above; and

20 (xxiii) a group represented by the formula



wherein the symbols are as defined above.


As the "alkyl group optionally having substituents" represented by R⁴, for example, C₁₋₆ alkyl group optionally
 25 having 1 to 3 substituent(s) selected from halogen atom, nitro group, cyano group, optionally halogenated C₃₋₆ cycloalkyl group, optionally halogenated C₁₋₆ alkoxy group, optionally halogenated C₁₋₆ alkylthio group, hydroxy group, amino group, mono-C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkyl-
 30 carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like), carboxyl group, C₁₋₆ alkoxy-carbonyl

group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆ alkyl-carbamoyl group, sulfo group, C₁₋₆ alkylsulfonyl group, C₁₋₆ alkylsulfinyl group and C₆₋₁₀ aryl group, and the like are preferable.

5 As the "alkylcarbonyl group optionally having substituents" represented by R⁴, for example, C₁₋₆ alkyl-carbonyl group (e.g., acetyl, ethylcarbonyl, propylcarbonyl and the like) optionally having 1 to 3 substituent(s) selected from halogen atom, nitro group, cyano group, optionally halogenated
10 C₃₋₆ cycloalkyl group, optionally halogenated C₁₋₆ alkoxy group, optionally halogenated C₁₋₆ alkylthio group, hydroxy group, amino group, mono-C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino, propionylamino, butyrylamino and the like), carboxyl group, C₁₋₆
15 alkoxy-carbonyl group, carbamoyl group, mono-C₁₋₆ alkyl-carbamoyl group, di-C₁₋₆ alkyl-carbamoyl group, sulfo group, C₁₋₆ alkylsulfonyl group, C₁₋₆ alkylsulfinyl group and C₆₋₁₀ aryl group, and the like are preferable.

As R⁴, hydrogen atom, C₁₋₆ alkyl group (e.g., C₁₋₃ alkyl
20 group such as methyl, ethyl and the like) and the like are preferable. Of these, hydrogen atom and methyl group are preferable, and hydrogen atom is particularly preferable.

The monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents, which is formed by R³ and
25 R⁴ together with the adjacent nitrogen atom is, for example, a group represented by the formula




wherein R¹⁸ is halogen atom, oxo group, optionally halogenated C₁₋₆ alkyl group or optionally halogenated C₁₋₆ alkoxy group, and
30 the like are preferable.

As j, 0 is preferable.

Furthermore, a compound used for the preparation of the present invention, a compound wherein preferable groups represented by the aforementioned symbols are optionally combined and the like are preferably used. Specific examples
5 include the following compounds and the like.


(1) A compound represented by the formula



wherein R¹⁹ is (i) hydrogen atom, (ii) carboxyl, (iii) C₁₋₆ alkoxy-carbonyl group (e.g., C₁₋₃ alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like), (iv) C₁₋₆ alkyl group (e.g., C₁₋₃ alkyl group such as methyl, ethyl, propyl and the like) optionally having substituents selected from the group consisting of carboxyl, C₁₋₆ alkyl-carbonyl (e.g., C₁₋₃ alkyl-carbonyl such as acetyl, ethylcarbonyl and the like), C₁₋₆ alkoxy-carbonyl (e.g., C₁₋₃ alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl and the like), C₁₋₆ alkoxy-carbonylamino (e.g., methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, butoxycarbonylamino, t-butoxycarbonylamino) and C₇₋₁₆ aralkyloxy-carbonylamino (e.g., benzyloxycarbonylamino), (v) a mono- or di-C₁₋₆ alkylamino group (e.g., mono- or di-C₁₋₃ alkylamino such as methylamino, ethylamino, dimethylamino, diethylamino and the like) or (iv) C₆₋₁₄ aryloxy group (e.g., phenyloxy), P¹ is C₁₋₃ alkylene group, Q¹ is C₁₋₃ alkylene group, X² is CH, C-OH or N, Y² is hydrogen atom, halogen atom (e.g., fluorine, chlorine), optionally halogenated C₁₋₆ alkyl group (e.g., optionally halogenated C₁₋₃ alkyl group such as methyl, ethyl, propyl, trifluoromethyl and the like) or C₁₋₆ alkoxy group (e.g., C₁₋₃ alkoxy group such as methoxy, ethoxy and the like), and Z is CO, SO or SO₂.


(preferably CO)], or a salt thereof.

(2) A compound represented by the formula




wherein R²⁰ is (i) hydrogen atom or (ii) C₁₋₆ alkyl group (e.g.,
 5 C₁₋₃ alkyl group such as methyl, ethyl, propyl and the like)
 optionally having substituents selected from the group
 consisting of C₁₋₆ alkoxy-carbonylamino (e.g.,
 methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino,
 butoxycarbonylamino, t-butoxycarbonylamino) and C₇₋₁₆
 10 aralkyloxy-carbonylamino (e.g., benzyloxycarbonylamino), P² is
 C₁₋₃ alkylene group (e.g., methylene, ethylene and trimethylene,
 preferably trimethylene), X³ is CH, C-OH or N (preferably CH),
 Y³ is hydrogen atom, halogen atom (e.g., fluorine, chlorine) or
 C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, propoxy), or a salt
 15 thereof.


(3) A compound represented by the formula



wherein R²¹ is nitrogen-containing heterocyclic group
 represented by (i) the formula



wherein X⁴ is CH or N, and Y⁴ is hydrogen atom, halogen atom or
 C₁₋₆ alkoxy group, or (ii) the formula



wherein the symbols in the formula are as defined above, or a salt thereof.

More preferable compounds are exemplified by, but not limited to, the following and the like.

Reference Example IA-1: 1-(5-amino-4,4-diphenylpentyl)-4-phenylpiperidine

Reference Example IA-2: 3,4-dihydro-6-methoxy-1'-(5-amino-4,4-diphenylpentyl)spiro[naphthalene-2(1H),2'-piperidine]

Reference Example IA-3: 1-[5-amino-4-(4-methoxyphenyl)-4-phenylpentyl]-4-phenylpiperidine

Reference Example IA-4: 1-[5-amino-4,4-bis(4-chlorophenyl)-pentyl]-4-(4-fluorophenyl)piperazine

Reference Example IA-5: 3,4-dihydro-6-methoxy-1'-(6-amino-4,4-diphenylhexyl)spiro[naphthalene-2(1H),2'-piperidine]

Reference Example IA-6: 3,4-dihydro-6,7-dimethoxy-1'-(7-amino-4,4-diphenylheptyl)spiro[naphthalene-2(1H),2'-piperidine]

Reference Example IIA-1: 1-(N,N-dimethylamino)-4,4-diphenyl-5-(formylamino)pentane

Reference Example IIA-2: 1-(N-benzyl-N-methylamino)-4,4-diphenyl-5-(formylamino)pentane hydrochloride

Reference Example IIA-3: 4,4-diphenyl-5-formylamino-1-(morpholino)pentane hydrochloride

Reference Example IIA-4: 4,4-diphenyl-5-formylamino-1-(2,3,4,5-tetrahydro-1H-3-benzazepin-3-yl)pentane hydrochloride

Reference Example IIA-5: 4,4-diphenyl-5-formylamino-1-(4-phenylpiperidino)pentane hydrochloride

Reference Example IIA-6: 1-[4-(4-fluorophenyl)piperazin-1-yl]-5-formylamino-4,4-diphenylpentane dihydrochloride

Reference Example IIA-7: 3,4-dihydro-6-methoxy-1'-(5-formylamino-4,4-diphenylpentyl) spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IIA-8: 1-benzylamino-4,4-diphenyl-5-(tosylamino)pentane hydrochloride

Reference Example IIA-9: 1-(N-benzyl-N-methylamino)-4,4-diphenyl-5-(tosylamino)pentane hydrochloride

Reference Example IIA-10: 4,4-diphenyl-1-(3-nitrobenzylamino)-5-(tosylamino)pentane hydrochloride

¹⁰ Reference Example IIA-11: 1-(3-aminobenzylamino)-4,4-diphenyl-5-(tosylamino)pentane

Reference Example IIA-12: 4,4-diphenyl-1-[3-(methoxycarbonyl)-benzylamino]-5-(tosylamino)pentane hydrochloride

Reference Example IIA-13: 4,4-diphenyl-1-(2-picolylamino)-5-
15 (tosylamino)pentane dihydrochloride

Reference Example IIA-14: 4,4-diphenyl-1-(1-hexamethyleneimino)-5-(tosylamino)pentane hydrochloride

Reference Example IIA-15: 4,4-diphenyl-1-(4-phenylpiperazin-1-yl)-5-(tosylamino)pentane

20 Reference Example IIA-16: 4,4-diphenyl-1-[4-(2-methoxyphenyl)-
piperazin-1-yl]-5-(tosylamino)pentane hydrochloride

Reference Example IIA-17: 4,4-diphenyl-5-mesylamino-1-(4-phenylpiperidino)pentane hydrochloride

Reference Example IIA-18: 5-benzenesulfonylamino-4,4-diphenyl-
25 1-(4-phenylpiperidino)pentane

Reference Example IIA-19: 4,4-diphenyl-1-(4-phenylpiperidino)-5-(2,4,6-trimethylbenzenesulfonylamino)pentane

Reference Example IIA-20: 4,4-diphenyl-1-(4-phenylpiperidino)-5-(2,4,6-triisopropylbenzenesulfonylamino)pentane

30 Reference Example IIA-21: 4,4-diphenyl-5-(1-naphthylsulfonylamino)-1-(4-phenylpiperidino)pentane

Reference Example IIA-22: 4,4-diphenyl-5-(2-naphthylsulfonylamino)-1-(4-phenylpiperidino)pentane

Reference Example IIA-23: 3,4-dihydro-6-methoxy-1'-(5-acetylamino-4,4-diphenylpentyl) spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IIA-24: 3,4-dihydro-6-methoxy-1'-(5-tosylamino-4,4-diphenylpentyl) spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IIA-25: 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-phenylpiperidino)pentane hydrochloride

Reference Example IIA-26: 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-phenylpiperazin-1-yl)pentane dihydrochloride

Reference Example IIA-27: 4-(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-formylamino-4-phenylpentane dihydrochloride

Reference Example IIA-28: 4-(4-chlorophenyl)-1-[4-(diphenylmethyl)piperazin-1-yl]-5-formylamino-4-phenylpentane

Reference Example IIA-29: 5-formylamino-4-(4-methoxyphenyl)-4-phenyl-1-(4-phenylpiperidino)pentane hydrochloride

Reference Example IIA-30: 4-(4-methoxyphenyl)-5-(1-naphthylsulfonylamino)-4-phenyl-1-(4-phenylpiperidino)pentane hydrochloride

Reference Example IIA-31: 4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(formylamino)pentane dihydrochloride

Reference Example IIA-32: 4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(mesylamino)pentane dihydrochloride

Reference Example IIA-33: 4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(tosylamino)pentane dihydrochloride

Reference Example IIA-34: 1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-5,5-diphenylhexane dihydrochloride

Reference Example IIA-35: 1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-4,4-diphenylhexane dihydrochloride

Reference Example IIA-37: 3,4-dihydro-6-methoxy-1'-(6-acetylamino-4,4-diphenylhexyl) spiro[naphthalene-2(1H),2'-

Reference Example IIA-38: 3,4-dihydro-6-methoxy-1'-(6-tosylamino-4,4-diphenylhexyl) spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example 1B-1: 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride

Reference Example 1B-3: 1-formylamino-5-(4-hydroxy-4-phenylpiperidino)-2,2-diphenylpentane hydrochloride

Reference Example 1B-5: 5-[4-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride

Reference Example 1B-7: 5-[4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridin-1-yl]-1-formylamino-2,2-diphenylpentane

Reference Example 1B-8: 1-formylamino-2,2-diphenyl-5-(4-phenylpiperidino)pentane

formylamino-2,2-diphenylpentane hydrochloride

Reference Example 1B-10: 7-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-4,4-diphenylheptane hydrochloride

5 Reference Example 2B-1: 5-[4-(4-fluorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride

Reference Example 2B-2: 1-formylamino-5-[4-hydroxy-4-(4-methoxyphenyl)piperidino]-2,2-diphenylpentane hydrochloride

10 Reference Example 2B-3: 1-formylamino-5-[4-hydroxy-4-(2-pyridyl)piperidino]-2,2-diphenylpentane dihydrochloride

Reference Example 3B-1: 1-acetylamino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 3B-2: 1-acetoacetylamino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 3B-3: ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamate hydrochloride

Reference Example 3B-4: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid

Reference Example 3B-5: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-ethylurea

Reference Example 3B-6: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]methanesulfonamide

25 hydrochloride

Reference Example 3B-7: phenyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]carbamate

Reference Example 3B-8: 1-acetylamino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentane


30 dihydrochloride

Reference Example 3B-9: ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]oxamate hydrochloride

Reference Example 3B-10: ethyl N-[5-[4-(4-chlorophenyl)-4-

- hydroxypiperidino]-2,2-diphenylpentyl]malonamate hydrochloride
Reference Example 3B-11: ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glutarmate
Example 1: benzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-
5 pentyl)amino)-2-oxoethylcarbamate hydrochloride
Example 2: tert-butyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethylcarbamate
Example 3: 4,4-diphenyl-7-(4-phenylpiperidino)heptylamine dihydrochloride
10 Example 4: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-4-methylbenzenesulfonamide hydrochloride
Example 5: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-acetamide hydrochloride
Example 6: N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)-
15 heptyl)amine dihydrochloride
Example 7: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(3-methoxybenzyl)amine dihydrochloride
Example 8: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-methoxybenzyl)amine dihydrochloride
20 Example 9: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-fluorobenzyl)amine dihydrochloride
Example 10: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-thiophenecarboxamide hydrochloride
Example 11: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-
25 phenylacetamide hydrochloride
Example 12: N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-thienylmethyl)amine dihydrochloride
Example 13: N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)-heptyl)-N-methylamine dihydrochloride
30 compounds of Example 14 - Example 131.

Of the above-mentioned compounds (I), a compound represented by the formula



wherein R^{23} is C_{1-6} alkyl group (e.g., methyl, ethyl, propyl and the like) having C_{7-16} aralkyloxy-carbonylamino (e.g., benzyloxycarbonylamino) optionally having substituents selected from the group consisting of halogen atom (e.g., fluorine atom, chlorine atom, iodine atom), C_{1-6} alkoxy (e.g., methoxy, ethoxy) and C_{1-6} alkyl (e.g., methyl, ethyl), P^4 is C_{1-3} alkylene group (e.g., methylene, ethylene, trimethylene), X^5 is CH, C-OH or N, Y^5 is hydrogen atom, halogen atom (e.g., fluorine atom, chlorine atom, iodine atom) or C_{1-6} alkoxy (e.g., methoxy, ethoxy, propoxy); R^{26} is hydrogen atom or C_{1-6} alkyl group; Y^6 and Y^7 are the same or different and each is hydrogen atom, halogen atom (e.g., fluorine atom, chlorine atom, iodine atom), optionally halogenated C_{1-6} alkyl group (e.g., methyl, ethyl, trifluoromethyl) or optionally halogenated C_{1-6} alkoxy group (methoxy, ethoxy, trifluoromethoxy) and a salt thereof are novel compounds.

As R^{23} , C_{1-3} alkyl group (e.g., methyl, ethyl, propyl) having benzyloxycarbonylamino and the like, and the like are preferable.

As P^4 , trimethylene is preferable.

As Y^5 , hydrogen atom, fluorine atom and methoxy are preferable.

As Y^6 and Y^7 , hydrogen atom is preferable.

As R^{26} , hydrogen atom is preferable.

As specific examples of compound (Ia), the compounds of Examples 1, 57, 58, 75, 76, 77, 80 and 103 and the like are preferable.

A compound represented by the aforementioned formula (Ib)

or (Ic) and a salt thereof are also novel compounds.

As specific examples of compound (Ib), the compounds of Examples 44, 45, 47, 104, 105, 106, 107, 108, 109, 115, 116, 117, 118, 120, 121, 122, 124, 125, 127, 128, 130 and 131 and
 5 the like are preferable.

As specific examples of compound (Ic), the compounds of Examples 51, 59 and 65 and the like are preferable.

The prodrug of the compound (Ia) of the present invention may be a compound that is converted to compound (Ia) by a
 10 reaction with an enzyme, gastric acid and the like in the body under physiological conditions. In other words, it may be a compound that undergoes enzymatic oxidation, reduction, hydrolysis and the like into compound (Ia), or a compound that undergoes hydrolysis and the like due to gastric acid and the
 15 like into compound (Ia).

Examples of the prodrug of compound (Ia) includes compound (Ia) wherein amino group is acylated, alkylated or phosphorylated (e.g., compound (Ia) wherein amino group is eicosanoylated, alanylated, pentylaminocarbonylated, (5-methyl-
 20 2-oxo-1,3-dioxolen-4-yl)methoxycarbonylated, tetrahydrofuranylated, pyrrolidylmethylated, pivaloyloxymethylated, tert-butylated and the like); compound (Ia) wherein hydroxyl group is acylated, alkylated, phosphorylated or borated (e.g., a compound (Ia) wherein
 25 hydroxyl group is acetylated, palmitoylated, propanoylated, pivaloylated, succinylated, fumarylated, alanylated, dimethylaminomethylcarbonylated and the like); compound (Ia) wherein carboxyl group is esterified or amidated (e.g., compound (Ia) wherein carboxyl group is ethyl esterified,
 30 phenyl esterified, carboxymethyl esterified, dimethylaminomethyl esterified, pivaloyloxymethyl esterified, ethoxycarbonyloxyethyl esterified, phthalidyl esterified, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl esterified,

cyclohexyloxycarbonylethyl esterified, methyl amidated and the like); and the like. These compounds can be produced from compound (Ia) by a method known *per se*.

The prodrug of compound (Ia) may be a compound that is converted to compound (Ia) under the physiological conditions described in Development of Pharmaceutical Products, vol. 7, Molecule Design, pp. 163-198, Hirokawa Shoten (1990).

The compounds (Ib) and (Ic) may be used as a prodrug, and as the prodrug, those similar to the prodrugs of the aforementioned compound (Ia) are mentioned.

When compound (I) forms a salt and the salt is used as a pharmaceutical product, it is preferably a pharmaceutically acceptable salt.

Examples of the pharmaceutically acceptable salt are, but not limited to, salts with inorganic acid salts such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromate and nitrate; salts with organic acid such as acetate, malate, maleate, fumarate, tartrate, succinate, citrate, lactate, methanesulfonate, p-toluenesulfonate, palmitin acid, salicylate and stearate.

As the pharmaceutically acceptable salt, salts with inorganic base, salts with organic base and the like are also mentioned.

Preferable examples of salts with inorganic base include alkali metal salts such as sodium salt, potassium salt and the like; alkaline earth metal salts such as calcium salt, magnesium salt and the like; aluminum salt, ammonium salt and the like.

Preferable examples of salts with organic base include salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N-dibenzylethylenediamine and the like.


The compound (I) or a salt thereof may be labeled with

an isotope (e.g., ^3H , ^{14}C , ^{35}S , ^{125}I and the like).

The compound (I) or a salt thereof may be anhydrate or hydrate.

The compound (I) or a salt thereof to be used for the
 5 agent of the present invention can be produced according to a method known *per se*, such as a method described in, for example, JP-A-8-253447, JP-A-10-81665, JP-A-11-71350 and the like or a similar method.


For example, compound (Ia) or a salt thereof can be
 10 produced by reacting a compound represented by the formula



wherein each symbol is as defined above, or a salt thereof, with a reactive derivative of an organic acid represented by the formula



wherein R^{23} is as defined above, for acylation, or reacting a compound represented by the formula




wherein each symbol is as defined above, or a salt thereof,
 20 with a reactive derivative represented by the formula




wherein R^{32} is C_{7-16} aralkyloxy-carbonyl group and X is a leaving group;


the compound (Ib) or a salt thereof can be produced by reacting
 25 a compound represented by the formula



wherein each symbol is as defined above, or a salt thereof, with a reactive derivative of an organic acid represented by the formula




wherein each symbol is as defined above, for acylation; and the compound (Ic) or a salt thereof can be produced by reacting a compound represented by the formula




10 wherein each symbol is as defined above, or a salt thereof, with,

(1) when Alk_2 is C_{1-6} alkylene group optionally having substituents, a reactive derivative of an organic acid represented by the formula



wherein each symbol is as defined above, for acylation, or

(2) when Alk_2 is a bond, by reacting with a reactive derivative represented by the formula



20 wherein X is a leaving group and other symbols are as defined above.

The compounds (Ia-1), (Ia-3), (Ib-1) and (Ic-1) and salts thereof can be produced according to a method known *per se*, such as a method described in, for example, JP-A-8-253447, JP-
25 A-10-81665, JP-A-11-71350 and the like or a similar method.

⁵ with preference given to active ester.

¹⁰ aralkyl phenyl carbonate, aralkyl p-nitrophenyl carbonate, N-((aralkyloxy)carbonyloxy)succinimide, 1-

²⁰ (e.g., phenyl carbamate, p-nitrophenyl carbamate, N-((amino)carbonyloxy)succinimide, 1-((amino)carbonyl)imidazole and the like), isocyanate and the like are used. Of these, isocyanate is preferable.

²⁵ according to a known method [e.g., method described in ORGANIC FUNCTIONAL GROUP PREPARATIONS 2nd ed., ACADEMIC PRESS, INC.].

30 reactive derivative represented by (Ia-4) or (Ic-3) and compound (Ia-1), (Ia-3), (Ib-1), (Ic-1) or a salt thereof are reacted in an inert solvent at a reaction temperature of from about -20°C to about 50°C (preferably from about 0°C to at room


temperature), for a reaction time of about 5 min to about 100 h.

As the inert solvent, for example, ether solvent, halogen solvent, aromatic solvent, acetonitrile, N,N-dimethylformamide (DMF), acetone, methyl ethyl ketone, dimethyl sulfoxide (DMSO),
 5 water and the like can be used alone or in combination. Of these, acetonitrile, tetrahydrofuran, dichloromethane, chloroform and the like are preferable. In addition, by co-existence of 1 to 10 equivalents, preferably 1 to 3 equivalents, of a base may afford more smooth progress of the reaction.

10 As the base, both inorganic base and organic base are effective. Examples of the inorganic base include hydroxide, hydride, carbonate, hydrogencarbonate, salt with organic acid and the like of alkali metal and alkaline earth metal. Of these, potassium carbonate, sodium carbonate, sodium hydroxide,
 15 potassium hydroxide, sodium hydrogencarbonate and potassium hydrogencarbonate are preferable. As the organic base, pyridine, 2,6-lutidine, triethylamine, N,N-diisopropylethylamine and the like are mentioned. Of these, tertiary amines such as triethylamine, N,N-
 20 diisopropylethylamine and the like are preferable.

In the case of acylation of carboxylic acid with active ester, moreover, it can be produced by conducting reaction in the presence of 1 to 1.5 equivalents of carboxylic acid and a dehydrative condensation agent (1 to 1.5 equivalents) such as
 25 dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC) and the like, in an inert solvent (e.g., halogen solvent, acetonitrile, tetrahydrofuran) at about 0°C to at room temperature, for about 0.5 to about 24 h. In this case, the co-presence of 1 to 1.5
 30 equivalents of an activation agent of carboxylic acid, such as N-hydroxysuccinimide (HOSu), 1-hydroxybenztriazole (HOBt), N-hydroxy-5-norbornene-2,3-dicarboxyimide (HONB) and the like, may afford more smooth progress of the reaction.

A compound represented by the formula



wherein R^{27} is hydrogen atom, halogen atom (e.g., fluorine atom, chlorine atom, bromine atom and the like), optionally
 halogenated C_{1-6} alkyl group (e.g., methyl, ethyl, propyl m
 trifluoromethyl and the like) or optionally halogenated C_{1-6}
 alkoxy group (e.g., methoxy, ethoxy, propoxy, trifluoromethoxy
 and the like) or a salt thereof is a novel synthetic
 intermediate for the production of compound (Ib) or a salt
 thereof. This synthetic intermediate can be produced according
 to a method known *per se*, such as a method described in, for
 example, JP-A-8-253447, JP-A-10-81665, JP-A-11-71350 and the
 like or a similar method.

Inasmuch as the compound (I) or a pharmaceutically
 acceptable salt thereof has a superior MCH receptor
 antagonistic action, it is useful as an agent for the
 prophylaxis or therapy of diseases caused by MCH. In addition,
 the compound of the present invention shows low toxicity, and
 superior oral absorption performance and transfer into the
 brain.

Accordingly, a melanin-concentrating hormone antagonist
 (hereinafter sometimes to be briefly referred to as MCH
 antagonist) containing the compound (I) or a pharmaceutically
 acceptable salt thereof is safely administered as an agent for
 the prophylaxis or therapy of the diseases caused by MCH to
 mammals (e.g., rat, mouse, guinea pig, rabbit, sheep, horse,

pig, cow, monkey, human and the like).

The diseases caused by MCH include, for example, obesity [e.g., malignant mastocytosis, exogenous obesity, hyperinsular obesity, hyperplasmic obesity, hypophyseal
5 adiposity, hypoplasmic obesity, hypothyroid obesity, hypothalamic obesity, symptomatic obesity, infantile obesity, upper body obesity, alimentary obesity, hypogonadal obesity, systemic mastocytosis, simple obesity, central obesity and the like], hyperphagia, emotional disorder, sexual dysfunction and
10 the like.

The compound (I) or a pharmaceutically acceptable salt thereof is useful as an agent for the prophylaxis or treatment of life-style related diseases such as diabetes, diabetic complications (e.g., diabetic retinopathy, diabetic neuropathy,
15 diabetic nephropathy and the like), arteriosclerosis, gonarthrititis and the like.

Furthermore, the compound or a pharmaceutically acceptable salt thereof is useful as an agent for suppressing food intake.

20 The MCH antagonist and a pharmaceutical composition of the present invention can be concurrently used with diet therapy (e.g., diet therapy for diabetes and the like), or an exercise therapy.

The MCH antagonist and a pharmaceutical composition of
25 the present invention can be produced by formulating compound (I), (Ia), (Ib), (Ic) or a pharmaceutically acceptable salt thereof as it is or along with a pharmacologically acceptable carrier according to a method known *per se*.

As the pharmacologically acceptable carrier, various
30 organic or inorganic carrier substance conventionally used as a material for preparation, such as excipient, lubricant, binder, disintegrant for solid preparation; solvent, dissolution aids, suspending agent, isotonicity agent, buffer, soothing agent and

10

15

20

25

30

94

Examples of the isotonicity agent include glucose, D-sorbitol, sodium chloride, glycerine, D-mannitol and the like.

Examples of the buffer include buffers such as phosphate, acetate, carbonate, citrate and the like and the like.

5 Examples of the soothing agent include benzyl alcohol and the like.

Examples of the preservative include p-oxybenzoic acid esters, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid and the like.

10 Examples of the antioxidant include sulfite, ascorbic acid and the like.

Examples of the dosage form of the MCH antagonist and a pharmaceutical composition of the present invention include oral preparations such as tablet (inclusive of sugar-coated
15 tablet and firm coated tablet), powder, granule, capsule (inclusive of soft capsule), liquid and the like; parenteral preparations such as injection (e.g., subcutaneous injection, intravenous injection, intramuscular injection, intraperitoneal injection and the like), external preparation (e.g., transnasal
20 administration preparation, percutaneous preparation, ointment and the like), suppository (e.g., rectal suppository, pessary and the like), sustained-release preparation (e.g., sustained release microcapsule and the like), pellet, drops and the like; and the like, and can be safely administered orally or
25 parenterally (e.g., topical, rectal, intravenous administration and the like).

The content of compound (I) or a pharmaceutically acceptable salt thereof in the MCH antagonist of the present invention, and the content of compound (Ia), (Ib), (Ic) or a
30 pharmaceutically acceptable salt thereof in the pharmaceutical composition of the present invention is, for example, about 0.1 to 100 wt% of the whole amount of the MCH antagonist or the pharmaceutical composition, respectively.

GI-262570, MCC-555, YM-440, DRF-2593, BM-13-1258, KRP-297, R-119702, CS-011 and the like are mentioned.

As the insulin secretagogue, for example, sulfonylurea agent is mentioned. Examples of the sulfonylurea agent include
 5 tolbutamide, chlorpropamide, tolazamide, acetohexamide, glyclopyramide and ammonium salt thereof, glibenclamide, gliclazide, glimepiride and the like.

In addition to the above, insulin secretagogue includes, for example, repaglinide, nateglinide, mitiglinide (KAD-1229),
 10 JTT-608 and the like.

As the biguanide agent, for example, metformin, buformin, phenformin and the like are mentioned.

As the insulin, for example, animal insulin extracted from pancreas of cow and swine; semi-synthetic human insulin
 15 enzymatically synthesized from insulin extracted from pancreas of swine; human insulin genetically synthesized using Escherichia coli or yeast; and the like are mentioned. As insulin, insulin zinc containing 0.45 to 0.9 (w/w)% of zinc; protamine insulin zinc produced from zinc chloride, protamine
 20 sulfate and insulin, and the like can be used. Moreover, insulin can be a fragment or derivative thereof (e.g., INS-1 and the like).

While insulin includes various types such as very rapid acting type, short-acting type, biphasic type, intermediate-
 25 acting type, extended type and the like, which can be determined depending on the disease state of patients.

As the α -glucosidase inhibitor, for example, acarbose, voglibose, miglitol, emiglitate and the like are mentioned.

As the β 3 adrenergic receptor agonist, for example, AJ-
 30 9677, BMS-196085, SB-226552, AZ40140 and the like are mentioned.

In addition to the above, the "antidiabetic agent" includes, for example, ergoset, pramlintide, leptin, BAY-27-9955 and the like.

As the above-mentioned "diabetes complication treatment agent", for example, aldose reductase inhibitor, glycation inhibitor, protein kinase C inhibitor and the like are mentioned.

5 As the aldose reductase inhibitor, for example, tolrestat; epalrestat; imirestat; zenarestat; SNK-860; zopolrestat; ARI-509; AS-3201 and the like are mentioned.

As the glycation inhibitor, for example, pimagedine and the like are mentioned.

10 As the protein kinase C inhibitor, for example, NGF, LY-333531 and the like are mentioned.

In addition to the above, the "diabetes complication treatment agent" includes, for example, alprostadil, tiapride hydrochloride, cilostazol, mexiletine hydrochloride, ethyl
15 icosapentate, memantine, pimagedline (ALT-711) and the like.

As the above-mentioned "anti-obesity agent other than MCH antagonist", for example, lipase inhibitor, anorexiant and the like are mentioned.

As the lipase inhibitor, for example, orlistat and the
20 like are mentioned.

As the anorexiant, for example, mazindol, dexfenfluramine, fluoxetine, sibutramine, biamine and the like are mentioned.

In addition to the above, the "anti-obesity agent other than MCH antagonist" includes, for example, lipstatin and the
25 like.

As the above-mentioned "hypertension treatment agent", for example, angiotensin converting enzyme inhibitor, calcium antagonist, potassium channel opener, angiotensin II antagonist and the like are mentioned.

30 As the angiotensin converting enzyme inhibitor, for example, captoril, enalapril, alacepril, delapril (HCl), lisinopril, imidapril, benazepril, cilazapril, temocapril, trandolapril, manidipine (HCl) and the like are mentioned.

As the calcium antagonist, for example, nifedipine, amlodipine, efonidipine, nicardipine and the like are mentioned.

As the potassium channel opener, for example, levcromakalim, L-27152, AL 0671, NIP-121 and the like are⁵ mentioned.

As the angiotensin II antagonist, for example, losartan, candesartan cilexetil, valsartan, irbesartan, CS-866, E4177 and the like are mentioned.

As the above-mentioned "hyperlipidemia treatment agent",
 10 for example, HMG-CoA reductase inhibitor, fibrate compound and
 the like are mentioned.

As the HMG-CoA reductase inhibitor, for example, pravastatin, simvastatin, lovastatin, atorvastatin, fluvastatin, lipantil, cerivastatin, itavastatin, ZD-4522 or salts thereof (e.g., sodium salt and the like) and the like are mentioned.

As the fibrate compound, for example, bezafibrate, cinnofibrate, clofibrate, simfibrate and the like are mentioned.

As the above-mentioned "arthritis treatment agent", for example, ibuprofen and the like are mentioned.

20 As the above-mentioned "anti-anxiety", for example, chlórdiazepoxide, diazepam, oxazolam, medazepam, cloxazolam, bromazepam, lorazepam, alprazolam, fludiazepam and the like are mentioned.

As the above-mentioned "antidepressant", for example,
25 fluoxetine, fluvoxamine, imipramine, paroxetine, sertraline and
the like are mentioned.

The time of administration of the aforementioned combination drug is not limited. The MCH antagonist or a pharmaceutical composition and a combination drug may be simultaneously administered to an administration subject or administered in a staggered manner. The dose of the combination drug can be determined according to the dose clinically employed, and can be determined as appropriate

depending on the administration subject, administration route, disease, combination and the like.

The mode of administration of the combination drug is not particularly limited, and may be any as long as the MCH antagonist or pharmaceutical composition and combination drug are combined on administration. Such administration mode is exemplified by (1) administration of a single preparation obtained by simultaneously formulating the MCH antagonist or pharmaceutical composition and combination drug(s), (2) simultaneous administration by the same administration route of two kinds of preparations obtained by separately formulating the MCH antagonist or pharmaceutical composition and combination drug(s), (3) staggered administration by the same administration route of two kinds of preparations obtained by separately formulating the MCH antagonist or pharmaceutical composition and combination drug, (4) simultaneous administration by different administration routes of two kinds of preparations obtained by separately formulating the MCH antagonist or pharmaceutical composition and combination drug, (5) staggered administration by different administration routes of two kinds of preparations obtained by separately formulating the MCH antagonist or pharmaceutical composition and combination drug (e.g., administration of MCH antagonist or pharmaceutical composition and combination drug in this order, and administration in the reversed order) and the like.

The admixing ratio of the MCH antagonist or pharmaceutical composition and combination drug can be appropriately determined depending on the administration subject, administration route, disease and the like.

The present invention is explained in detail in the following by referring to Reference Examples, Examples, Formulation Examples and Experimental Examples. The present invention is not limited by these examples, and may be modified

as long as it does not deviate from the range of the present invention.

In the following Reference Example and Example, "room temperature" means 0 to 30°C and other definitions are as

⁵ follows.

s : singlet

d : doublet

t : triplet

q : quartet

10 m : multiplet

br : broad

```
brs      : broad singlet
```

ABq : AB quartet

dd : double doublet

¹⁵ J : coupling constant

Hz : Hertz

CDCl_3 : deuterated chloroform

THF : tetrahydrofuran

DMF : N,N-dimethylformamide

²⁰ DMSO : dimethyl sulfoxide

¹H-NMR : proton nuclear magnetic resonance (free compound was used for measurement)

In the present specification and drawings, when base, amino acid and the like are shown using abbreviations, they are
25 based on abbreviations according to IUPAC-IUB Commission on Biochemical Nomenclature and conventional abbreviations employed in this field. Examples thereof are given in the following. When optical isomer is present due to amino acid, it is an L form unless particularly indicated.

30 DNA : deoxyribonucleic acid

cDNA : complementary deoxyribonucleic acid

A : adenine

T : thymine

	G	: guanine
	C	: cytosine
	RNA	: ribonucleic acid
	mRNA	: messenger ribonucleic acid
5	dATP	: deoxyadenosine triphosphate
	dTTP	: deoxythymidine triphosphate
	dGTP	: deoxyguanosine triphosphate
	dCTP	: deoxycytidine triphosphate
	ATP	: adenosine triphosphate
10	EDTA	: ethylenediamine tetraacetic acid
	SDS	: sodium dodecylsulfate
	EIA	: enzyme immunoassay
	Gly	: glycine
	Ala	: alanine
15	Val	: valine
	Leu	: leucine
	Ile	: isoleucine
	Ser	: serine
	Thr	: threonine
20	Cys	: cysteine
	Met	: methionine
	Glu	: glutamic acid
	Asp	: aspartic acid
	Lys	: lysin
25	Arg	: arginine
	His	: histidine
	Phe	: phenylalanine
	Tyr	: thyrosin
	Trp	: tryptophan
30	Pro	: proline
	Asn	: asparagine
	Gln	: glutamine
	pGlu	: pyroglutamic acid

	Me	: methyl group
	Et	: ethyl group
	Bu	: butyl group
	Ph	: phenyl group
5	TC	: thiazolidine-4(R)-carboxamide group

The substituent, protecting group and reagent frequently appear in the present specification are shown using the following symbols.

	Tos	: p-toluenesulfonyl
10	CHO	: formyl
	Bzl	: benzyl
	Cl ₂ Bzl	: 2,6-dichlorobenzyl
	Bom	: benzyloxymethyl
	Z	: benzyloxycarbonyl
15	Cl-Z	: 2-chlorobenzyloxycarbonyl
	Br-Z	: 2-bromobenzyloxycarbonyl
	Boc	: t-butoxycarbonyl
	DNP	: dinitrophenol
	Trt	: trityl
20	Bom	: t-butoxymethyl
	Fmoc	: N-9-fluorenylmethoxycarbonyl
	HOBt	: 1-hydroxybenzotriazole
	HOObt	: 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine
	HONB	: N-hydroxy-5-norbornene-2,3-dicarboxyimide
25	DCC	: N,N'-dicyclohexylcarbodiimide
	WSC	: 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride
	DMAP	: 4-dimethylaminopyridine
	IPE	: diisopropyl ether
30	THF	: tetrahydrofuran
	DMF	: N,N-dimethylformamide

The sequence numbers in the Sequence Listing in the present specification show the following sequences.

- [SEQ ID NO:1]
Synthetic DNA used for screening cDNA encoding rat SLC-1.
- [SEQ ID NO:2]
Synthetic DNA used for screening cDNA encoding rat SLC-1.
- 5 [SEQ ID NO:3]
Full length amino acid sequence of rat SLC-1.
- [SEQ ID NO:4]
Full length base sequence of rat SLC-1 cDNA comprising Sal I
recognition sequence added on the 5' side and Spe I recognition
10 sequence added on the 3' side.
- [SEQ ID NO:5]
Riboprobe used for the determination of expression amount of
SLC-1 mRNA in each clone of rat SLC-1 expression CHO cells.
- [SEQ ID NO:6]
- 15 [SEQ ID NO:7]
Synthetic DNA used for obtaining cDNA encoding human SLC-1.
- [SEQ ID NO:8]
Primer used to make cDNA encoding human SLC-1 double-stranded.
- [SEQ ID NO:9]
Full length base sequence of cDNA encoding human SLC-1.
- 20 [SEQ ID NO:10]
Full length amino acid sequence of human SLC-1.
- [SEQ ID NO:11]
Synthetic DNA used for screening cDNA encoding human SLC-1(S).
- [SEQ ID NO:12]
25 Synthetic DNA used for screening cDNA encoding human SLC-1(S).
- [SEQ ID NO:13]
Synthetic DNA used for screening cDNA encoding human SLC-1(L).
- [SEQ ID NO:14]
30 Synthetic DNA used for screening cDNA encoding human SLC-1(L).
- [SEQ ID NO:15]
Full length base sequence of human SLC-1(S) cDNA comprising Sal
I recognition sequence added on the 5' side and Spe I
recognition sequence added on the 3' side.

[SEQ ID NO:15]

Full length base sequence of human SLC-1(L) cDNA comprising Sal I recognition sequence added on the 5' side and Spe I recognition sequence added on the 3' side.

5 [SEQ ID NO:16]

Riboprobe used for the determination of expression amount of SLC-1 mRNA in each clone of human SLC-1(S) expression CHO cells and human SLC-1(L) expression CHO cells.

The transformant *Escherichia coli* DH10B/phSLC1L8 with a
10 plasmid containing DNA encoding the base sequence obtained in Reference Example 6D, which is depicted in SEQ:No. 9 has been deposited at the National Institute of Bioscience and Human-Technology (NIBH) since February 1,1999 under deposit No. FERM BP-6632 and at the Institute for Fermentation, Osaka (IFO)
15 under deposit No. IFO 16254 since January 21, 1999.

Examples

The following Reference Examples IA-IVA can be produced according to JP-A-8-253447.

Reference Example IA-1: 1-(5-amino-4,4-diphenylpentyl)-4-
20 phenylpiperidine 1-(5-formamino-4,4-diphenylpentyl)-4-phenylpiperidine

Reference Example IA-2: 3,4-dihydro-6-methoxy-1'-(5-amino-4,4-diphenylpentyl) spiro[naphthalene-2(1H),2'-piperidine]

Reference Example IA-3: 1-[5-amino-4-(4-methoxyphenyl)-4-
25 phenylpentyl]-4-phenylpiperidine

Reference Example IA-4: 1-[5-amino-4,4-bis(4-chlorophenyl)-pentyl]-4-(4-fluorophenyl)piperazine

Reference Example IA-5: 3,4-dihydro-6-methoxy-1'-(6-amino-4,4-diphenylhexyl) spiro[naphthalene-2(1H),2'-piperidine]

30 Reference Example IA-6: 3,4-dihydro-6,7-dimethoxy-1'-(7-amino-4,4-diphenylheptyl) spiro[naphthalene-2(1H),2'-piperidine]

Reference Example IIA-1: 1-(N,N-dimethylamino)-4,4-diphenyl-5-(formylamino)pentane

- Reference Example IIA-18: 5-benzenesulfonylamino-4,4-diphenyl-1-(4-phenylpiperidino)pentane
- Reference Example IIA-19: 4,4-diphenyl-1-(4-phenylpiperidino)-5-(2,4,6-trimethylbenzenesulfonylamino)pentane
- 5 Reference Example IIA-20: 4,4-diphenyl-1-(4-phenylpiperidino)-5-(2,4,6-triisopropylbenzenesulfonylamino)pentane
- Reference Example IIA-21: 4,4-diphenyl-5-(1-naphthylsulfonylamino)-1-(4-phenylpiperidino)pentane
- Reference Example IIA-22: 4,4-diphenyl-5-(2-naphthylsulfonylamino)-1-(4-phenylpiperidino)pentane
- 10 Reference Example IIA-23: 3,4-dihydro-6-methoxy-1'-(5-acetylamino-4,4-diphenylpentyl)spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride
- Reference Example IIA-24: 3,4-dihydro-6-methoxy-1'-(5-tosylamino-4,4-diphenylpentyl)spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride
- 15 Reference Example IIA-25: 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-phenylpiperidino)pentane hydrochloride
- Reference Example IIA-26: 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-phenylpiperazin-1-yl)pentane dihydrochloride
- 20 Reference Example IIA-27: 4-(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-formylamino-4-phenylpentane 2 hydrochloride
- Reference Example IIA-28: 4-(4-chlorophenyl)-1-[4-(diphenylmethyl)piperazin-1-yl]-5-formylamino-4-phenylpentane
- 25 Reference Example IIA-29: 5-formylamino-4-(4-methoxyphenyl)-4-phenyl-1-(4-phenylpiperidino)pentane hydrochloride
- Reference Example IIA-30: 4-(4-methoxyphenyl)-5-(1-naphthylsulfonylamino)-4-phenyl-1-(4-phenylpiperidino)pentane
- 30 hydrochloride
- Reference Example IIA-31: 4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(formylamino)pentane dihydrochloride

Reference Example IIA-32: 4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(mesylamino)pentane dihydrochloride

Reference Example IIA-33: 4,4-bis(4-chlorophenyl)-1-[4-(4-
5 fluorophenyl)piperazin-1-yl]-5-(tosylamino)pentane
dihydrochloride

Reference Example IIA-34: 1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-5,5-diphenylhexane dihydrochloride

Reference Example IIA-35: 1-[4-(4-fluorophenyl)piperazin-1-yl]-
10 6-formylamino-4,4-diphenylhexane dihydrochloride

Reference Example IIA-36: 4,4-diphenyl-1-(4-phenylpiperidino)-6-(tosylamino)hexane hydrochloride

Reference Example IIA-37: 3,4-dihydro-6-methoxy-1'-(6-acetylamino-4,4-diphenylhexyl) spiro[naphthalene-2(1H),2'-
15 piperidine] dihydrochloride

Reference Example IIA-38: 3,4-dihydro-6-methoxy-1'-(6-tosylamino-4,4-diphenylhexyl)spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IIA-39: 3,4-dihydro-6-methoxy-1'-(6-
20 benzylamino-4,4-diphenylhexyl) spiro[naphthalene-2(1H),2'-
piperidine] dihydrochloride

Reference Example IIA-40: 7-acetylamino-4,4-diphenyl-1-[3-(methoxycarbonyl)benzylamino]heptane dihydrochloride

Reference Example IIA-41: 7-acetylamino-4,4-diphenyl-1-(β -phenethylamino)heptane dihydrochloride

Reference Example IIA-42: 7-acetylamino-1-[2-(6,7-dimethoxy-1,2,3,4-tetrahydronaphthylamino)]-4,4-diphenylheptane dihydrochloride

Reference Example IIA-43: 7-acetylamino-1-{N-benzyl-N-[2-(6,7-
30 dimethoxy-1,2,3,4-tetrahydronaphthyl)]amino}-4,4-
diphenylheptane dihydrochloride

Reference Example IIA-44: 1'-(7-acetylamino-4,4-diphenylheptyl)-3,4-dihydro-8-methoxyspiro[naphthalene-

2(1H),2'-piperidine] dihydrochloride

Reference Example IIA-45: 1'-(7-acetylamino-4,4-diphenylheptyl)-3,4-dihydro-6-methoxyspiro[naphthalene-

2(1H),2'-piperidine] dihydrochloride

- 5 Reference Example IIA-46: 1'-(7-acetylamino-4,4-diphenylheptyl)-3,4-dihydro-6,7-dimethoxyspiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IIA-47: 1'-[7-(cyclohexylacetyl)amino-4,4-diphenylheptyl]-3,4-dihydro-6,7-dimethoxyspiro[naphthalene-

- 10 2(1H),2'-piperidine] hydrochloride

Reference Example IIA-48: 3,4-dihydro-6,7-dimethoxy-1'-[4,4-diphenyl-7-(phenylacetyl)amino]heptyl]spiro[naphthalene-

2(1H),2'-piperidine] hydrochloride

- Reference Example IIA-49: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(2-fluorophenylacetyl)amino]heptyl)-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

- 15 Reference Example IIA-50: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(4-fluorophenylacetyl)amino]heptyl)-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

- 20 Reference Example IIA-51: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(4-chlorophenylacetyl)amino]heptyl)-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IIA-52: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(3-nitrophenylacetyl)amino]heptyl)-

- 25 spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IIA-53: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(4-nitrophenylacetyl)amino]heptyl)-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

- Reference Example IIA-54: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(4-methylphenylacetyl)amino]heptyl)-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

- 30 Reference Example IIA-55: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(4-trifluoromethylphenylacetyl)amino]heptyl)-

- spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
Reference Example IIA-56: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(2-methoxyphenylacetyl) amino]heptyl)-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
5 Reference Example IIA-57: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(3-methoxyphenylacetyl) amino]heptyl)-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
Reference Example IIA-58: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(4-methoxyphenylacetyl) amino]heptyl)-
10 spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
Reference Example IIA-59: 3,4-dihydro-6,7-dimethoxy-1'-(7-[(3,4-dimethoxyphenylacetyl) amino]-4,4-diphenylheptyl)-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
Reference Example IIA-60: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-
15 diphenyl-7-[(3,4-methylenedioxyphenylacetyl) amino]heptyl)-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
Reference Example IIA-61: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(phenoxyacetyl) amino]heptyl)spiro[naphthalene-
2(1H),2'-piperidine] hydrochloride
20 Reference Example IIA-62: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(2-thienylacetyl) amino]heptyl)spiro[naphthalene-
2(1H),2'-piperidine] hydrochloride
Reference Example IIA-63: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(3-thienylacetyl) amino]heptyl)spiro[naphthalene-
25 2(1H),2'-piperidine] hydrochloride
Reference Example IIA-64: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(3-phenylpropionyl) amino]heptyl)spiro[naphthalene-
2(1H),2'-piperidine] hydrochloride
Reference Example IIA-65: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-
30 diphenyl-7-[[3-(4-methoxyphenyl)propionyl] amino]heptyl)-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride
Reference Example IIA-66: 3,4-dihydro-6,7-dimethoxy-1'-(4,4-diphenyl-7-[(1-naphthylacetyl) amino]heptyl)spiro[naphthalene-

2 (1H), 2'-piperidine] hydrochloride

Reference Example IIA-67: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-[(2-naphthylacetyl) amino]heptyl}spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

5 Reference Example IIA-68: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-bis(4-fluorophenyl)-7-[(4-methoxyphenylacetyl)amino]heptyl}-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IIIA-1:

(1) 4,4-diphenyl-5-hydroxy-6-heptenenitrile

10 (2) 7-(6,7-dimethoxy-1,2,3,4-tetrahydronaphthalene-2-spiro-
2'-piperidin-1'-yl)-4,4-diphenyl-5-heptenenitrile
hydrochloride

(3) 1-(7-amino-4,4-diphenyl-2-heptenyl)-6',7'-dimethoxy-1',2',3',4'-tetrahydronaphthalene-2'-spiro-2-piperidine

15 Reference Example IIIA-2: N-(7-(6,7-dimethoxy-1,2,3,4-
tetrahydronaphthalene-2-spiro-2'-piperidine-1'-yl)-4,4-
diphenyl-5-heptenyl)-3-(4-methoxyphenyl)propionamide
hydrochloride

Reference Example IVA-1: 4,4-diphenyl-1-[(6-methoxy-1,2,3,4-tetrahydro-2-naphthyl)amino]-7-[[3-(4-methoxyphenyl)-propionyl]amino]heptane hydrochloride

Reference Example IVA-2: 4,4-diphenyl-1-[3-(4-methoxyphenyl)piperidino]-7-{[3-(4-methoxyphenyl)-propionyl]amino}heptane hydrochloride

25 Reference Example IVA-3: 4,4-diphenyl-1-(4-phenylpiperidino)-7-
[(3-phenylpropionyl)amino]heptane hydrochloride

Reference Example IVA-4: 4,4-diphenyl-1-[4-(3-methoxyphenyl)-piperidino]-7-[(3-phenylpropionyl)amino]heptane hydrochloride

Reference Example IVA-5: 4,4-diphenyl-1-[4-(4-

30 methoxyphenyl)piperidino]-7-{[3-(4-methoxyphenyl)propionyl]-
amino}heptane hydrochloride

Reference Example IVA-6: 4,4-diphenyl-7-([3-(4-methoxyphenyl)-propionyl]amino)-1-[2,3,4,5-tetrahydro-3(1H)-benzazepin-3-

yl]heptane hydrochloride

Reference Example IVA-7: 1-[7-acetyl-2,3,4,5-tetrahydro-3(1H)-benzazepin-3-yl]-4,4-diphenyl-7-([3-(4-methoxyphenyl)-propionyl]amino)heptane hydrochloride

5 Reference Example IVA-8: 4,4-diphenyl-1-(7,8-dimethoxy-2,3,4,5-tetrahydro-3(1H)-benzazepin-3-yl)-7-([3-(4-methoxyphenyl)-propionyl]amino)heptane hydrochloride

Reference Example IVA-9: 1-(8,9-dimethoxy-6,6-dimethyl-1,2,3,4,5,6-hexahydro-3-benzazocin-3-yl)-4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptane hydrochloride

Reference Example IVA-10: 4,4-diphenyl-7-([3-(4-methoxyphenyl)-propionyl]amino)-1-(cis-1,2,3,4,4a,9,10,10a-octahydrobenzo[f]quinolin-1-yl)heptane hydrochloride

Reference Example IVA-11: 1-(3-aza-6-methyl-1,1a,2,3,4,4a-hexahydro-9-fluorenon-3-yl)-4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptane hydrochloride

Reference Example IVA-12: 3,4-dihydro-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}spiro[naphthalene-2(1H),2'-pyrrolidine] hydrochloride

20 Reference Example IVA-13: 3,4-dihydro-6-methoxy-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IVA-14: 6-ethoxy-3,4-dihydro-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-15: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-([3-(4-dimethylaminophenyl)propionyl]amino)heptyl}-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-16: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-([3-(4-fluorophenyl)propionyl]amino)heptyl}-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-17: 3,4-dihydro-6,7-dimethoxy-1'-{7-([3-(4-chlorophenyl)propionyl]amino)-4,4-diphenylheptyl}-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-18: 3,4-dihydro-6,7-dimethoxy-1'-{7-([3-(3,5-difluorophenyl)propionyl]amino)-4,4-diphenylheptyl}-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

5 Reference Example IVA-19: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-([3-(4-pyridyl)propionyl]amino)heptyl}-
spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IVA-20: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-([2-(5-methoxyindane)carbonyl]amino)heptyl}-

10 spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-21: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-([3-(3,4-methylenedioxyphenyl)propionyl]amino)-heptyl}spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-22: 3,4-dihydro-6,7-dimethoxy-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-
15 spiro[naphthalene-2(1H),2'-piperidine]-1-one hydrochloride

Reference Example IVA-23: 3,4-dihydro-6-methoxy-5-nitro-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

20 Reference Example IVA-24: 3,4-dihydro-6-methoxy-7-nitro-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-25: 7-amino-3,4-dihydro-6-methoxy-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-

25 spiro[naphthalene-2(1H),2'-piperidine] dihydrochloride

Reference Example IVA-26: 7-acetylamino-3,4-dihydro-6-methoxy-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)-heptyl}spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-27: 7-acetyl-3,4-dihydro-6-methoxy-1'-
30 {4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-28: 3,4-dihydro-6,7-methylenedioxy-1'-{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-29: 6,7-diethoxy-3,4-dihydro-1'-(4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl)-spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

5 Reference Example IVA-30: 3,4-dihydro-1'-(4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl)spiro[naphthalene-2(1H),2'-hexamethylenimine] hydrochloride

Reference Example IVA-31: (+)-3,4-dihydro-6-methoxy-1'-(4,4-diphenyl-7-[(4-methoxyphenylacetyl)amino]heptyl)-

10 spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-32: (-)-3,4-dihydro-6-methoxy-1'-(4,4-diphenyl-7-[(4-methoxyphenylacetyl)amino]heptyl)-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-33: (-)-3,4-dihydro-6,7-dimethoxy-1'-

15 (4,4-diphenyl-7-[(4-methoxyphenylacetyl)amino]heptyl)-

spiro[naphthalene-2(1H),2'-piperidine]-1-one hydrochloride

Reference Example IVA-34: (-)-3,4-dihydro-6,7-dimethoxy-1'-

(4,4-diphenyl-7-[(4-fluorophenylacetyl)amino]heptyl)-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

20 Reference Example IVA-35: (+)-3,4-dihydro-6,7-dimethoxy-1'-

(4,4-diphenyl-7-[(4-fluorophenylacetyl)amino]heptyl)-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-36: (-)-3,4-dihydro-6,7-dimethoxy-1'-

(4,4-diphenyl-7-([3-(4-fluorophenyl)propionyl]amino)heptyl)-

25 spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-37: (+)-3,4-dihydro-6,7-dimethoxy-1'-

(4,4-diphenyl-7-([3-(4-fluorophenyl)propionyl]amino)heptyl)-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-38: (+)-3,4-dihydro-6,7-dimethoxy-1'-(7-

30 ([3-(4-chlorophenyl)propionyl]amino)-4,4-diphenylheptyl)-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-39: (-)-3,4-dihydro-6,7-dimethoxy-1'-(7-

([3-(4-chlorophenyl)propionyl]amino)-4,4-diphenylheptyl)-

spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-40: (-)-3,4-dihydro-6,7-dimethoxy-1'-
{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

5 Reference Example IVA-41: (+)-3,4-dihydro-6,7-dimethoxy-1'-
{4,4-diphenyl-7-([3-(4-methoxyphenyl)propionyl]amino)heptyl}-
spiro[naphthalene-2(1H),2'-piperidine] hydrochloride

Reference Example IVA-42: 3,4-dihydro-4'-{4,4-diphenyl-7-([3-
(4-methoxyphenyl)propionyl]amino)heptyl}spiro[naphthalene-
10 2(1H),3'-morpholine] hydrochloride

Reference Example IVA-43: 3,4-dihydro-7-methoxy-4'-{4,4-
diphenyl-7-[3-(4-methoxyphenyl)propionylamino]heptyl}-
spiro[naphthalene-2(1H),3'-morpholine] hydrochloride

Reference Example IVA-44: 3,4-dihydro-6,7-dimethoxy-4'-{4,4-
15 diphenyl-7-[3-(4-methoxyphenyl)propionylamino]heptyl}-
spiro[naphthalene-2(1H),3'-morpholine] hydrochloride

Reference Example IVA-45: 3,4-dihydro-6,7-dimethoxy-4'-methyl-
1'-{4,4-diphenyl-7-[3-(4-methoxyphenyl)propionylamino]heptyl}-
spiro[naphthalene-2(1H),2'-piperazine] dihydrochloride

20 Reference Examples 1B-40B can be produced according to
JP-A-10-81665.

Reference Example 1B-1: 5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-1-formylamino-2,2-diphenylpentane
hydrochloride

25 Reference Example 1B-2: 5-[4-(4-fluorophenyl)piperazin-1-yl]-1-
formylamino-2,2-diphenylpentane dihydrochloride

Reference Example 1B-3: 1-formylamino-5-(4-hydroxy-4-
phenylpiperidino)-2,2-diphenylpentane hydrochloride

Reference Example 1B-4: 5-[4-(4-trifluoromethylphenyl)-4-
30 hydroxypiperidino]-1-formylamino-2,2-diphenylpentane
hydrochloride

Reference Example 1B-5: 5-[4-[3,5-bis(trifluoromethyl)phenyl]-
4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane

hydrochloride

Reference Example 1B-6: 5-[4-(3,5-dichlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride

5 Reference Example 1B-7: 5-[4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridin-1-yl]-1-formylamino-2,2-diphenylpentane hydrochloride

Reference Example 1B-8: 1-formylamino-2,2-diphenyl-5-(4-phenylpiperidino)pentane

10 Reference Example 1B-9: 5-[4-(4-chlorophenyl)piperidino]-1-formylamino-2,2-diphenylpentane hydrochloride

Reference Example 1B-10: 7-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-4,4-diphenylheptane hydrochloride

15 Reference Example 2B-1: 5-[4-(4-fluorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride

Reference Example 2B-2: 1-formylamino-5-[4-hydroxy-4-(4-methoxyphenyl)piperidino]-2,2-diphenylpentane hydrochloride

20 Reference Example 2B-3: 1-formylamino-5-[4-hydroxy-4-(2-pyridyl)piperidino]-2,2-diphenylpentane dihydrochloride

Reference Example 3B-1: 1-acetylamino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 3B-2: 1-acetoacetylamino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 3B-3: ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamate hydrochloride

Reference Example 3B-4: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid

Reference Example 3B-5: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-ethylurea

Reference Example 3B-6: N-[5-[4-(4-chlorophenyl)-4-

hydroxypiperidino]-2,2-diphenylpentyl]methanesulfonamide
hydrochloride

Reference Example 3B-7: phenyl N-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]carbamate

5 Reference Example 3B-8: 1-acetylamino-5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentane
dihydrochloride

Reference Example 3B-9: ethyl N-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]oxamidate hydrochloride

10 Reference Example 3B-10: ethyl N-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]malonamate hydrochloride

Reference Example 3B-11: ethyl N-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]glutarmate

Reference Example 3B-12: ethyl N-[5-[4-(4-chlorophenyl)-4-
15 hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentyl]succinamate
dihydrochloride

Reference Example 4B-1: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-pentamethyleneurea
hydrochloride

20 Reference Example 4B-2: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-hydroxypropyl)urea
hydrochloride

Reference Example 4B-3: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(4-hydroxybutyl)urea

25 hydrochloride

Reference Example 4B-4: ethyl 3-[3-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]ureido]propionate

Reference Example 4B-5: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(2-

30 dimethylaminoethyl)urea

Reference Example 4B-6: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-
diethylaminopropyl)urea

Reference Example 4B-7: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[3-(2-pyrrolidone-1-yl)propyl]urea

Reference Example 4B-8: 1-[5-[4-(4-chlorophenyl)-4-
5 hydroxypiperidino]-2,2-diphenylpentyl]-3-(2-
piperidinoethyl)urea

Reference Example 4B-9: 2-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]ethanesulfonamide hydrochloride

10 Reference Example 4B-10: 2-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]ethanesulfonic acid

Reference Example 5B-1: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid

15 Reference Example 5B-2: N-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]oxamic acid

Reference Example 5B-3: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]malonic acid

Reference Example 5B-4: N-[5-[4-(4-chlorophenyl)-4-
20 hydroxypiperidino]-2,2-diphenylpentyl]glutaramic acid

Reference Example 5B-5: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentyl]succinamic acid

Reference Example 6B-1: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glycine ethyl ester
25 dihydrochloride

Reference Example 6B-2: ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-aminobutyrate dihydrochloride

Reference Example 7B-1: N-[5-[4-(4-chlorophenyl)-4-
30 hydroxypiperidino]-2,2-diphenylpentyl]glycine

Reference Example 7B-2: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-aminobutyric acid

Reference Example 8B-1: N-[5-[4-(4-chlorophenyl)-4-

- hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-hydroxypyrrolidin-1-yl)propanamide
- Reference Example 8B-2: 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-1-(3-pyrrolidin-1-yl-propionylamino)pentane
- Reference Example 8B-3: 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-[3-(dimethylamino)propionylamino]-2,2-diphenylpentane
- Reference Example 9B: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(t-butoxycarbonyl)-aminopropanamide
- Reference Example 10B: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-aminopropanamide dihydrochloride
- Reference Example 11B: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(acetylamino)-propanamide
- Reference Example 12B: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(propionylamino)propanamide
- Reference Example 13B: 1-[4,4-diphenyl-5-(phenyloxycarbonylamino)pentanoyl]-4-(4-chlorophenyl)-4-hydroxypiperidine
- Reference Example 14B: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]-3-[3-(hydroxy)propyl]urea
- Reference Example 15B: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]-3-[3-(dimethylamino)ethyl]urea
- Reference Example 16B: 1-(5-acetylamino-4,4-diphenylpentanoyl)-4-(4-chlorophenyl)-4-hydroxypiperidine
- Reference Example 17B: ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]succinamate

- Reference Example 18B: N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]succinamic acid
- Reference Example 19B: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]-3-[3-(2-oxo-1-pyrrolidino)propyl]urea
- Reference Example 20B: 5-[3-(4-chlorophenyl)-3-hydroxypyrrolidin-1-yl]-2,2-diphenyl-1-formylpentanamine
- Reference Example 21B: 1-[5-[4-(4-chlorophenyl)-3-hydroxypiperidino]-2,2-diphenylpentyl]-3-[3-(hydroxy)propyl]urea
- Reference Example 22B: 1-formylamino-[5-[4-hydroxy-4-(4-chlorophenyl)hexamethylenimin-1-yl]-2,2-diphenylpentane hydrochloride
- Reference Example 23B: 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-2-phenyl-2-(2-thienyl)pentane hydrochloride
- Reference Example 24B: 2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylaminopentane hydrochloride
- Reference Example 25B: ethyl N-[2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]]pentylsuccinamate hydrochloride
- Reference Example 26B: N-[2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]]pentylsuccinamic acid
- Reference Example 27B-1: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[(1-ethoxycarbonyl)piperidin-4-yl]urea
- Reference Example 27B-2: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[2-(1-pyrrolidino)ethyl]urea
- Reference Example 27B-3: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[2-(diethylamino)ethyl]urea

5 hydroxypiperidino]-2,2-diphenylpentyl]-3-(5-hydroxypentyl) urea

Reference Example 27B-6: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[2-(dimethylamino)-ethyl]-3-methylurea

Reference Example 27B-7: 1-[5-[4-(4-chlorophenyl)-4-
10 hydroxypiperidino]-2,2-diphenylpentyl]-3-[2-
(methylamino)ethyl]-3-methylurea

Reference Example 27B-8: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(2-hydroxyethyl)-3-methylurea

15 Reference Example 27B-9: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[2-(acetylamino)-ethyl]urea

Reference Example 27B-10: ethyl 4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureidobutyrate

20 Reference Example 27B-11: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-hydroxypropyl)urea

Reference Example 27B-12: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-benzylpiperidin-4-yl)urea

25 Reference Example 27B-13: N-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-4-methylpiperazine-1-
carboxamide

Reference Example 27B-14: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-benzylpiperazine-1-carboxamide

Reference Example 27B-15: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-1,2,4,5-tetrahydro-3-benzazepine-3-carboxamide

- Reference Example 27B-16: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(trifluoroacetyl amino)pyrrolidine-1-carboxamide
- Reference Example 27B-17: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-(t-butoxycarboxamide)piperidine-1-carboxamide
- Reference Example 27B-18: ethyl [4-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]piperidino]acetate
- Reference Example 27B-19: 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[1-(trifluoroacetyl)-piperidin-4-yl]urea
- Reference Example 27B-20: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-formyl-1-piperazinecarboxamide
- Reference Example 27B-21: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-(3-hydroxypropyl)-1-piperazinecarboxamide
- Reference Example 27B-22: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-(ethoxycarbonyl)-1-piperazinecarboxamide
- Reference Example 27B-23: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-(morpholinocarbonylmethyl)-1-piperazinecarboxamide
- Reference Example 28B-1: 3-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]propionic acid
- Reference Example 28B-2: 4-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]butyric acid
- Reference Example 29B: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]vinylsulfonamide
- Reference Example 30B: 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-2-(pyrrolidino)-ethylsulfonamide
- Reference Example 31B: 1-[5-[4-(4-chlorophenyl)-4-

hydroxypiperidino]-2,2-diphenylpentyl]-3-[3-(carbamoyloxy)-
propyl]urea

Reference Example 32B: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(piperidin-4-yl)urea

5 Reference Example 33B-1: ethyl 4-[4-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]-
piperidino-4-oxobutyrate

Reference Example 33B-2: N-ethyl-4-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-
10 piperidinecarboxamide

Reference Example 33B-3: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-acetylpiperidin-4-
yl)urea

Reference Example 33B-4: N-ethoxycarbonylmethyl-4-[5-[4-(4-
15 chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-
aminocarbonylamino-1-piperidinecarboxamide

Reference Example 33B-5: ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]-
piperidino-3-oxopropionate

20 Reference Example 34B-1: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-ethylpiperidin-4-
yl)urea

Reference Example 34B-2: 1-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]-3-[1-(2-hydroxyethyl)-
25 piperidin-4-yl]urea

Reference Example 34B-3: ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-
hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-
piperidino]propionate

Reference Example 34B-4: 1-[5-[4-(4-chlorophenyl)-4-
30 hydroxypiperidino]-2,2-diphenylpentyl]-3-[1-(3-hydroxypropyl)-
piperidin-4-yl]urea

Reference Example 35B: 1-[(piperidin-4-yl)carboxamide]-5-[4-(4-
chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane

dihydrochloride

Reference Example 36B-1: 1-[(N-ethylpiperidin-4-yl)carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

5 Reference Example 36B-2: 1-[[N-(ethoxycarbonylmethyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

Reference Example 36B-3: 1-[[N-(2-morpholinoethyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-
10 diphenylpentane 3 hydrochloride

Reference Example 36B-4: 1-[[N-(2-dimethylaminoethyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane 3 hydrochloride

Reference Example 37B-1: 1-[[N-ethylcarbamoyl]piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-
15 diphenylpentane hydrochloride

Reference Example 37B-2: 1-[[N-methylcarbamoyl]piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

20 Reference Example 37B-3: 1-[[N-phenylcarbamoyl]piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 37B-4: 1-[[N-(4-chlorobenzoyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-
25 diphenylpentane hydrochloride

Reference Example 37B-5: 1-[[N-(ethoxycarbonylacetyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 37B-6: 1-[[N-(3-methoxycarbonylpropionyl)-piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-
30 hydroxypiperidino]-2,2-diphenylpentane hydrochloride

Reference Example 37B-7: 1-[[N-(nicotinoyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-

diphenylpentane dihydrochloride

Reference Example 37B-8: 1-[[N-(4-dimethylaminobutyryl)-piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

5 Reference Example 38B: 1-[(N-propylpiperidin-4-yl)carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

Reference Example 39B: 1-[[N-3-pyridylacetyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

10 Reference Example 40B: 1-[[N-ethylcarbamoyl)piperidin-4-yl]carboxamide]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

Reference Examples 1C-15C can be produced according to
15 JP-A-11-71350.

Reference Example 1C: 1-(tert-butoxycarbonyl)piperidin-4-yl N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl) carbamate

Reference Example 2C: piperidin-4-yl N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl) carbamate

Reference Example 3C: 1-(N-ethylcarbamoyl)piperidin-4-yl N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-carbamate

Reference Example 4C: 1-(nicotinoyl)piperidin-4-yl N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl) carbamate

Reference Example 5C-1: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(2-chlorethoxy carbonyl)piperidin-4-yl)urea

Reference Example 5C-2: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(acetoxycetyl)-piperidin-4-yl)urea

Reference Example 5C-3: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(nicotinoyl)-

piperidin-4-yl) urea

Reference Example 5C-4: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(isonicotinoyl)-piperidin-4-yl)urea

5 Reference Example 5C-5: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(benzoyl)piperidin-4-yl)urea

Reference Example 6C: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(2-hydroxyacetyl)-piperidin-4-yl)urea

Reference Example 7C: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-3-(1-(2-pyrrolidin-1-yl)ethyloxycarbonyl)piperidin-4-yl)urea

Reference Example 8C-1: N-(5-(4-(4-chlorophenyl)-4-
15 hydroxypiperidino)-2,2-diphenylpentyl)nicotinamide
dihydrochloride

Reference Example 8C-2: 2-chloroethyl (5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentylamino) carbamate

Reference Example 9C: 1-(5-(4-(4-chlorophenyl)-4-
20 hydroxypiperidino)-2,2-diphenylpentyl)-4,5-dihydro-2-oxazolone

Reference Example 10C-1: 2-(1-(t-butoxycarbonyl)piperidin-4-yl)-N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)acetamide

Reference Example 10C-2: 2-(1-(t-butoxycarbonyl)piperidin-4-ylidene)-N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)acetamide

Reference Example 11C-1: N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-diphenylpentyl)-2-(piperidin-4-yl)acetamide dihydrochloride

30 Reference Example 11C-2: N-(5-(4-(4-chlorophenyl)-4-
hydroxypiperidino)-2,2-diphenylpentyl)-2-(piperidine-4-
ylidene)acetamide dihydrochloride

Reference Example 12C-1: N-(5-(4-(4-chlorophenyl)-4-

hydroxypiperidino)-2-phenyl-2-(2-pyridyl)pentyl)-1-(ethoxycarbonyl)piperidine-4-carboxamide dihydrochloride

Reference Example 12C-2: N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2-phenyl-2-(2-pyridyl)pentyl)-1-(isonicotinoyl)piperidine-4-carboxamide 3 hydrochloride

Reference Example 12C-3: N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-bis(4-fluorophenyl)pentyl)-1-(isonicotinoyl)piperidine-4-carboxamide dihydrochloride

Reference Example 12C-4: N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2-(4-fluorophenyl)-2-phenylpentyl)-1-(isonicotinoyl)piperidine-4-carboxamide dihydrochloride

Reference Example 13C-1: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2-phenyl-2-(2-pyridyl)pentyl)-3-(3-hydroxypropyl)urea dihydrochloride

Reference Example 13C-2: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2-phenyl-2-(2-pyridyl)pentyl)-3-(1-(nicotinoyl)piperidin-4-yl)urea 3 hydrochloride

Reference Example 13C-3: 1-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2-phenyl-2-(2-pyridyl)pentyl)-3-(1-(isonicotinoyl)piperidin-4-yl)urea 3 hydrochloride

Reference Example 13C-4: 1-(5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-bis(4-fluorophenyl)pentyl)-3-(1-(isonicotinoyl)piperidin-4-yl)urea dihydrochloride

Reference Example 13C-5: 1-(5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-bis(4-fluorophenyl)pentyl)-3-(3-hydroxypropyl)urea

Reference Example 13C-6: 1-(5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-bis(4-fluorophenyl)pentyl)-3-(1-(nicotinoyl)piperidin-4-yl)urea

Reference Example 13C-7: 1-(5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-(4-fluorophenyl)-2-phenylpentyl)-3-(1-(isonicotinoyl)piperidin-4-yl)urea dihydrochloride

Reference Example 13C-8: 1-(5-[4-(4-chlorophenyl)-4-

hydroxypiperidino]-2-(4-fluorophenyl)-2-phenylpentyl)-3-(3-hydroxypropyl)urea hydrochloride

Reference Example 13C-9: 1-(5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-(4-fluorophenyl)-2-phenylpentyl)-3-(1-


5 (nicotinoyl)piperidin-4-yl)urea dihydrochloride

Reference Example 14C: N-(5-(4-(4-chlorophenyl)-4-hydroxypiperidino)-2,2-bis(4-fluorophenyl)pentyl)acetamide hydrochloride

Reference Example 15C: N-(5-[4-(4-chlorophenyl)-4-
10 hydroxypiperidino]-2-(4-fluorophenyl)-2-phenylpentyl)acetamide
hydrochloride

Reference Example 1D

benzyl 2-((5-hydroxy-2,2-diphenylpentyl) amino)-2-oxoethylcarbamate




15

To a solution of 5-amino-4,4-diphenylpentanol (15.8 g) in acetonitrile (100 ml) were added 2-(((benzyloxy)carbonyl)-amino)acetic acid (13 g) and WSC (14 g). The mixture was stirred at room temperature overnight. The reaction mixture was concentrated and ethyl acetate and water were added to the residue. The organic layer was separated, washed with saturated aqueous sodium hydrogen carbonate, dried and concentrated. The residue was recrystallized from IPE/ethyl acetate to give the title compound (21 g).
melting point: 122-123°C.

25 melting point: 122-123°C.

Reference Example 2D

tert-butyl 2-((5-hydroxy-2,2-diphenylpentyl) amino)-2-oxoethylcarbamate




To a solution of 5-amino-4,4-diphenylpentanol (4 g) in acetonitrile (30 ml) were added 2-(((tert-butoxy)carbonyl)-amino)acetic acid (3.5 g), WSC(4 g) and triethylamine(5 ml).
 5 The mixture was stirred at room temperature overnight. The reaction mixture was concentrated and ethyl acetate and water were added to the residue. The organic layer was separated, washed with saturated aqueous sodium hydrogen carbonate, dried and concentrated. The residue was purified by silica gel
 10 column chromatography (eluent: ethyl acetate) to give the title compound (4 g).

oil:

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.2-1.6 (2H, m), 1.43 (9H, s), 1.802.3 (2H, m), 3.4-3.6 (2H, m), 3.69 (2H, d), 4.04 (2H, d), 5.0 (1H, br), 5.70
 15 (1H, br), 7.1-7.4 (10H, m).

Reference Example 3D

2,2-diphenyl-5-(4-phenylpiperidino)pentanenitrile




To a solution of 5-bromo-2,2-diphenylpentanenitrile (9.5
 20 g) in acetonitrile (100 ml) were added potassium carbonate (6 g) and 4-phenylpiperidine (4.8 g). The reaction mixture was stirred at 60°C overnight and concentrated. The residue was recrystallized from IPE/ethyl acetate to give the title

compound (11 g).

melting point: 88-89°C.

Reference Example 4D

7-oxo-4,4-diphenyl-7-(4-phenylpiperidino)heptanenitrile



5


To a solution of 6-cyano-4,4-diphenylhexanoic acid (5.87 g) in dichloromethane (60 ml) was added thionyl chloride (3.57 g) under ice-cooling and the mixture was stirred at room temperature for 1 h. To the reaction mixture was added a solution of phenylpiperidine (4.8 g) and triethylamine (5 g) in dichloromethane (20 ml) by small portions and the mixture was stirred at room temperature for another 1h. The reaction mixture was washed with 1N hydrochloric acid and saturated brine and dried over anhydrous sodium sulfate. Concentration under reduced pressure gave the residue, which was subject to silica gel column chromatography with elution of hexane-ethyl acetate (4:1-1:1) to give the title compound (7.5 g).

oil:

¹H-NMR(CDCl₃)δ: 1.19-1.95 (4H, m), 1.96-2.08 (4H, m), 2.20-2.79 (6H, m), 2.91 (1H, dt, J=2.6, 18.0 Hz), 3.38-3.52 (1H, m), 4.66-4.80 (1H, m), 7.10-7.38 (15H, m).

Reference Example 5D

2,2-diphenyl-5-(4-phenylpiperidino)pentylamine



2,2-diphenyl-5-(4-phenylpiperidino)pentanenitrile (10.0 g) produced in Reference Example 3D was added to a suspension of lithium aluminum hydride (4.8 g) and aluminum chloride (16.9 g) in THF (350 ml) under ice-cooling and the mixture was stirred at room temperature for 3 h. To the reaction mixture was added 1N aqueous sodium hydroxide solution (400 ml) by small portions under ice-cooling and the mixture was stirred for 5 min. Ether (500 ml) was added and the mixture was filtered through Celite. The organic layer of the filtrate was washed with saturated brine and dried over anhydrous sodium sulfate. Concentration under reduced pressure gave the residue, which was subject to silica gel column chromatography with elution of ethyl acetate-methanol (5:1)-ethyl acetate-methanol-saturated aqueous ammonia (50:10:1) to give the title compound (8.1 g).

oil:


$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.25-1.36 (4H, m), 1.69-2.03 (6H, m), 2.10-2.18 (2H, m), 2.28-2.48 (3H, m), 2.90 (2H, d, $J=11.4$ Hz), 3.33 (2H, s), 7.12-7.32 (15H, m).

Reference Example 6D

benzyl 2-((5-(4-methylbenzenesulfonyloxy)-2,2-diphenylpentyl)-amino)-2-oxoethylcarbamate

Reference Example 7D

132




In the same manner as in Reference Example 6D, the compound was synthesized from the compound obtained in Reference Example 8D.

5 oil.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.34-1.42 (2H, m), 2.01-2.09 (2H, m), 2.43 (3H, s), 3.73 (2H, d, $J=5.6$ Hz), 3.90-3.96 (4H, m), 5.08 (2H, s), 5.29 (1H, s), 5.48 (1H, s), 7.09 (4H, d, $J=7.0$ Hz), 7.19-7.36 (13H, m), 7.73 (2H, d, $J=8.2$ Hz).

10 Reference Example 10D

benzyl 2-((4-iodo-2,2-diphenylbutyl)amino)-2-oxoethylcarbamate



In the same manner as in Reference Example 7D, the compound was synthesized from the compound obtained in


15 Reference Example 9D.

amorphous powder.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.52-2.70 (2H, m), 3.37-3.59 (2H, m), 3.92-4.10 (4H, m), 4.18 (2H, s), 5.13 (2H, s), 5.78 (1H, s), 7.15-7.36 (15H, m).

Reference Example 11D

tert-butyl 2-((2,2-bis(4-chlorophenyl)-5-hydroxypentyl)amino)-
2-oxoethylcarbamate




5 In the same manner as in Reference Example 2D, the compound was synthesized from 5-amino-4,4-bis(4-chlorophenyl)-1-pentanol.

amorphous powder.

¹H-NMR(CDCl₃)δ: 1.23-1.35 (2H, m), 1.41 (9H, s), 1.85 (1H, t, J=5.6 Hz), 2.05-2.14 (2H, m), 3.55 (2H, q, J=5.6 Hz), 3.68 (2H, d, J=6.0 Hz), 3.96 (2H, d, J=6.2 Hz), 4.98 (1H, br), 5.75 (1H, br), 7.13 (4H, d, J=8.7 Hz), 7.29 (4H, d, J=8.7Hz).

Reference Example 12D

tert-butyl 2-((2,2-bis(4-chlorophenyl)-5-(4-methylbenzenesulfonyloxy)pentyl)amino)-2-oxoethylcarbamate




In the same manner as in Reference Example 6D, the

compound was synthesized from the compound obtained in
Reference Example 11D.
amorphous powder.

¹H-NMR(CDCl₃)δ: 1.30-1.38 (2H, m), 1.40 (9H, s), 2.01-2.09 (2H,
5 m), 2.45 (3H, s), 3.68 (2H, d, J=6.1 Hz), 3.88-3.96 (4H, m),
4.94 (1H, br), 5.65 (1H, br), 7.02-7.12 (5H, m), 7.24-7.35 (5H,
m), 7.74 (2H, d, J=8.3 Hz).

Reference Example 13D

tert-butyl 2-((2,2-bis(4-chlorophenyl)-5-iodopentyl)amino)-2-
10 oxoethylcarbamate




In the same manner as in Reference Example 7D, the
compound was synthesized from the compound obtained in
Reference Example 12D.

15 amorphous powder.

¹H-NMR(CDCl₃)δ: 1.41 (9H, s), 1.47-1.56 (2H, m), 2.04-2.17 (2H,
m), 3.10 (2H, t, J=6.4 Hz), 3.68 (2H, d, J=6.2 Hz), 3.94 (2H, d,
J=6.2 Hz), 4.89 (1H, br), 5.68 (1H, br), 7.06-7.13 (4H, m),
7.25-7.33 (4H, m).

20 Reference Example 14D

bis(4-fluorophenyl)acetonitrile




Thionyl chloride (50 ml) was added to bis(4-fluorophenyl)methanol (24.2 g) at 0°C and after stirring for 30 min, the mixture was poured into 2N hydrochloric acid (500 ml).
 5 The mixture was extracted with ethyl acetate and the organic layer was dried over calcium chloride and concentrated under reduced pressure. The resulting residue was dissolved in dichloromethane (200 ml) and after addition of trimethylsilylcyanoide (16.4 ml), titanium tetrachloride (13.4
 10 ml) was added dropwise at 0°C. The mixture was stirred for 50 min. Methanol (5 ml) was added to the reaction mixture and the mixture was poured into saturated aqueous sodium hydrogen carbonate. The mixture was extracted with ethyl acetate and washed with saturated brine. The organic layer was dried over
 15 anhydrous magnesium sulfate and concentrated under reduced pressure to give the title compound (21.8 g).
 oil.

¹H-NMR(CDCl₃)δ: 5.11 (1H, s), 7.03-7.11 (4H, m), 7.26-7.33 (4H, m).

20 Reference Example 15D

ethyl 4,4-bis(4-fluorophenyl)-4-cyanobutyrates




A mixture of bis(4-fluorophenyl)acetonitrile (20.4 g) obtained in Reference Example 14D, ethanol (150 ml), 1,8-diazabicyclo[5.4.0]-7-undecene (2.63 ml) and ethyl acrylate
 5 (12.5 ml) was heated under reflux for 18 h. After cooling, the reaction mixture was concentrated under reduced pressure. 2N Hydrochloric acid (200 ml) was added and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, dried over anhydrous magnesium sulfate
 10 and concentrated under reduced pressure. The resulting residue was subject to silica gel column chromatography with elution of hexane-ethyl acetate (5:1) to give the title compound (28.8 g). oil.

¹H-NMR(CDCl₃) δ: 1.25 (3H, t, J=7.2 Hz), 2.42-2.47 (2H, m),
 15 2.70-2.75 (2H, m), 4.13 (2H, q, J=7.1 Hz), 7.06-7.13 (4H, m), 7.34-7.39 (4H, m).

Reference Example 16D

2,2-bis(4-fluorophenyl)-5-hydroxypentylamine




20 To a mixture of ethyl 4,4-bis(4-fluorophenyl)-4-cyanobutyrates (34 g) obtained in Reference Example 15D and THF

7.17 (4H, m).

Reference Example 18D

tert-butyl 2-((2,2-bis(4-fluorophenyl)-5-(4-methylbenzenesulfonyloxy)pentyl)amino)-2-oxoethylcarbamate




5

In the same manner as in Reference Example 6D, the compound was synthesized from the compound obtained in Reference Example 17D.
oil.

¹H-NMR(CDCl₃)δ: 1.25-1.40 (11H, m), 2.03-2.08 (2H, m), 2.45 (3H, s), 3.67 (2H, d, J= 6.1Hz), 3.89-3.95 (4H, m), 4.94 (1H, s), 5.61-5.65 (1H, m), 6.96-7.11 (8H, m), 7.33 (2H, d, J= 8.2Hz), 7.74 (2H, d, J= 8.3Hz).

Reference Example 19D

tert-butyl 2-((2,2-bis(4-fluorophenyl)-5-iodopentyl)amino)-2-oxoethylcarbamate




In the same manner as in Reference Example 7D, the compound was synthesized from the compound obtained in Reference Example 18D.

oil.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.41 (9H, s), 1.47-1.57 (2H, m), 2.08-2.14 (2H, m), 2.39-3.23 (1H, m), 3.82-3.92 (7H, m), 5.77-5.81 (1H, m), 6.84-7.06 (6H, m), 7.19-7.27 (6H, m).

Example 1

benzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate hydrochloride




To a solution of triphenylphosphine (520 mg) in acetonitrile (10 ml) was added bromine (320 mg) under ice-cooling. A solution of benzyl 2-((5-hydroxy-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate (0.88 g) in acetonitrile (10 ml) was then added dropwise to the reaction mixture. The mixture was stirred at room temperature for 1 h

and concentrated. The residue was dissolved in ethyl acetate and the mixture was washed with water, dried and concentrated. The residue was purified by silica gel column chromatography (eluent; IPE: ethyl acetate=1:1) to give a bromide. To a
 5 solution of the bromide in acetonitrile (20 ml) were added 4-phenylpiperidine (320 mg) and potassium carbonate (300 mg). The reaction mixture was stirred at 40°C overnight, poured into water and extracted with ethyl acetate. The organic layer was washed with water, dried and concentrated. The residue was
 10 purified by alumina column chromatography (eluent: ethyl acetate) and converted to hydrochloride. Recrystallization from ethyl acetate/ethanol gave the title compound (0.56g). melting point: 167-168°C.

Example 2


15 tert-butyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethylcarbamate



In the same manner as in Example 1, the compound was synthesized from tert-butyl 2-((5-hydroxy-2,2-diphenylpentyl)-
 20 amino)-2-oxoethylcarbamate. melting point: 145-146°C

Example 3


4,4-diphenyl-7-(4-phenylpiperidino)heptylamine dihydrochloride



To a solution of 7-oxo-4,4-diphenyl-7-(4-phenylpiperidino)heptanenitrile (2.2 g) in THF (20 ml) was added a suspension of lithium aluminum hydride (760 mg) in THF (40 ml) under ice-cooling and the mixture was stirred at 60°C for 14 h. After completion of the reaction, 1N aqueous sodium hydroxide solution was added dropwise slowly and the precipitated crystals were filtered off. The filtrate was concentrated. The resulting residue was dissolved in ethyl acetate and the organic layer was washed with saturated brine, dried and concentrated. The residue was converted to hydrochloride and recrystallized from dichloromethane-IPE to give the title compound (2.0 g).
melting point: 155-159°C.

Example 4

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-4-methylbenzenesulfonamide hydrochloride



HCl


To a solution of 4,4-diphenyl-7-(4-phenylpiperidino)-heptylamine (500 mg) in dichloromethane (15 ml) were added triethylamine (3 ml), p-tosyl chloride (209 mg, 1.1 mmol) and DMAP (catalytic amount) under ice-cooling and the mixture was stirred at room temperature for 1 h. After completion of the reaction, the solvent was evaporated off under reduced pressure and the resulting residue was subject to silica gel column chromatography. The resultant was purified with hexane-ethyl acetate (1:1), made to be a hydrochloride and recrystallized from chloroform-IPE to give the title compound (420 mg).

heptylamine (426 mg), benzaldehyde (106 mg) and p-tosylic acid hydrate (catalytic amount) in benzene (5 ml) was added anhydrous magnesium sulfate (1 g) and the mixture was stirred at 50°C for 1 h. The precipitate was filtered off and the filtrate was concentrated under reduced pressure. The resulting residue was dissolved in methanol (5 ml) and sodium borohydride (38 mg) was added. The mixture was stirred at room temperature for 5 min. After completion of the reaction, the mixture was concentrated under reduced pressure and the resulting residue was subject to silica gel column chromatography. The residue was purified by elution with ethyl acetate-methanol (1:0-20:1), made to be a hydrochloride and recrystallized from chloroform-IPE to give the title compound (350 mg).

melting point: 223-226°C

Example 7

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(3-methoxybenzyl)amine dihydrochloride




The compound was synthesized in the same manner as in Example 6.

Recrystallization solvent: chloroform-IPE.

melting point: 215-217°C.

Example 8

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-methoxybenzyl)amine dihydrochloride




The compound was synthesized in the same manner as in Example 6.

Recrystallization solvent: chloroform-IPE.

5 melting point: 100-108°C.

Example 9

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-fluorobenzyl)amine dihydrochloride




10 The compound was synthesized in the same manner as in Example 6.

Recrystallization solvent: chloroform-IPE

melting point: 198-200°C.

Example 10


15 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-thiophenecarboxamide hydrochloride



To a solution of 4,4-diphenyl-7-(4-phenylpiperidino)-
heptylamine (426 mg) in ethyl acetate (10 ml) was added a
saturated aqueous sodium carbonate solution (10 ml) and 2-
5 thiophenecarbonyl chloride (146 mg) was added while vigorously
stirring the mixture. After 30 min, the organic layer was
separated, washed with saturated brine, dried and concentrated.
The residue was applied to silica gel column chromatography,
eluted with hexane-ethyl acetate (1:1), converted to
10 hydrochloride and recrystallized from chloroform-IPE to give
the title compound (0.5 g).
melting point: 125-130°C.

Example 11

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-phenylacetamide
15 hydrochloride




The compound was synthesized in the same manner as in
Example 10.

Recrystallization solvent: chloroform-IPE.

20 melting point: 103-110°C

Example 12


N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-thienylmethyl)amine dihydrochloride



5 To a solution of a free form (300 mg) of N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-thiophenecarboxamide in THF (5 ml) was added a suspension of lithium aluminum hydride (114 mg) in THF (5 ml) under ice-cooling, and the mixture was heated under reflux for 12 h. After the completion of the reaction, a
10 1N aqueous sodium hydroxide solution was dropwise added gradually. The precipitated crystals were filtered off and the filtrate was concentrated. The residue was dissolved in ethyl acetate, and the organic layer was washed with saturated brine, dried and concentrated to give the title compound (300 mg) as
15 an amorphous. A part of the obtained compound was converted to hydrochloride, and recrystallized from chloroform-IPE.
melting point: 120-125°C.

Example 13


N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-methylamine dihydrochloride
20



To a solution of N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)amine (175 mg) in acetonitrile (5 ml)-37% formalin (0.3 ml) were added sodium cyanoborohydride (31 mg) and acetic acid (0.5 ml) and the mixture was stirred at room temperature for 1 h. After the completion of the reaction, the mixture was concentrated under reduced pressure, and the resulting residue was applied to silica gel column chromatography, purified by elution with ethyl acetate, converted to hydrochloride and recrystallized from chloroform-IPE to give the title compound (130 mg).
melting point: 115-120°C.

Example 14

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-6-phenylhexanamide
15 phenylhexanamide




To a solution of 6-phenylhexanoic acid (80 mg) in THF (5

ml) were added oxalyl chloride (63 mg) and one drop of DMF under ice-cooling, and the mixture was stirred at room temperature for 2 h and concentrated under reduced pressure. This residue was added to a solution of 2,2-diphenyl-5-(4-phenylpiperidino)pentanamine (110 mg) synthesized in Reference Example 5D and triethylamine (56 mg) in THF (10 ml) under ice-cooling, and the mixture was stirred for 1 h. To the reaction mixture was added saturated aqueous sodium hydrogen carbonate (100 ml), and the mixture was extracted with ethyl acetate (100 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate. The residue was concentrated under reduced pressure and the resulting residue was applied to silica gel column chromatography and eluted with ethyl acetate-hexane (2:1)-ethyl acetate to give the title compound (110 mg).
 15 amorphous powder.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.25-1.38 (5H, m), 1.45-1.68 (5H, m), 1.72-1.78 (2H, m), 1.82-2.03 (6H, m), 2.28 (2H, t, $J=7.3$ Hz), 2.25-2.45 (1H, m), 2.52 (2H, t, $J=7.6$ Hz), 2.88 (2H, d, $J=11.4$ Hz), 3.99 (2H, d, $J=5.9$ Hz), 4.96 (1H, t, $J=5.6$ Hz), 7.12-7.34 (20H, m).

20 **Example 15**

4-benzyloxy-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-butylamide




The compound was synthesized in the same manner as in Example 14.

amorphous powder.

¹H-NMR (CDCl₃)δ: 1.27-1.38 (2H, m), 1.68-2.44 (15H, m), 2.87-
 5 2.92 (2H, m), 3.41 (2H, t, J=6.0Hz), 3.99 (2H, d, J=5.9Hz),
 4.39 (2H, s), 5.14 (1H, t, J=5.7Hz), 7.15-7.31 (20H, m).

Example 16

4-benzoyl-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-
 butylamide



10


The compound was synthesized in the same manner as in Example 14.

Recrystallization solvent: ethyl acetate-hexane.

melting point: 101-102°C.

15 **Example 17**


N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-4-phenylbenzamide



To a solution of 2,2-diphenyl-5-(4-phenylpiperidino)-
 pentanamine (150 mg) synthesized in Reference Example 5D and
 triethylamine (57 mg) in THF (5 ml) was added 4-
 5 biphenylcarbonyl chloride (98 mg) under ice-cooling, and the
 mixture was stirred for 1.5 h. To the reaction mixture was
 added saturated aqueous sodium hydrogen carbonate (100 ml).
 The mixture was extracted with ethyl acetate (100 ml), and
 after washing with saturated brine, dried over anhydrous sodium
 10 sulfate and concentrated under reduced pressure. The resulting
 residue was applied to silica gel column chromatography and
 eluted with ethyl acetate-ethyl acetate-methanol (20:1) to give
 the title compound (110 mg).
 Recrystallization solvent: ether-hexane.
 15 melting point: 155-156°C

Example 18

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-3-
 ethoxycarbonylpropanamide




The compound was synthesized in the same manner as in Example 17.

amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.19-1.31 (5H, m), 1.68-2.60 (15H, m), 2.85-2.91 (2H, m), 3.99-4.02 (2H, m), 4.09 (2H, q, $J=7.1\text{Hz}$), 7.17-7.33 (15H, m).

Example 19

2-acetoxy-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-
10 acetamide




The compound was synthesized in the same manner as in Example 17.

Recrystallization solvent: diethyl ether-hexane.

- 15 melting point: 106-107°C.

Example 20

benzyl (1S)-2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-1-methyl-2-oxoethylcarbamate




To a solution of 2,2-diphenyl-5-(4-phenylpiperidino)-
 pentanamine (110 mg) synthesized in Reference Example 5D, Z-L-
 alanine (68 mg) and HOBt (41 mg) in acetonitrile (10 ml) was
 5 added WSC (58 mg) at -20°C , and the mixture was stirred at room
 temperature for 18 h. Water (100 ml) was added to the reaction
 mixture. The mixture was extracted with ethyl acetate (100 ml),
 and after washing with saturated brine, dried over anhydrous
 sodium sulfate and concentrated under reduced pressure. The
 10 resulting residue was applied to silica gel column
 chromatography and eluted with ethyl acetate-methanol (100:1-
 25:1) to give the title compound (120 mg).
 amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.39 (3H, d, $J=7.3\text{Hz}$), 1.59 (4H, s), 1.96 (2H,
 15 br), 2.58-2.92 (7H, m), 3.53-3.59 (1H, m), 3.71-3.82 (2H, m),
 4.11-4.25 (2H, m), 5.07 (2H, s), 5.89 (1H, br), 6.30 (1H, br),
 7.12-7.40 (20H, m).

Example 21

benzyl (1S)-1-(((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-
 20 amino)carbonyl)-3-methylbutylcarbamate




The compound was synthesized in the same manner as in Example 20.

amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 0.84 (6H, d, $J=6.2\text{Hz}$), 1.21-2.09 (12H, m), 2.22-2.45 (4H, m), 2.84-2.90 (2H, m), 3.87-4.03 (2H, m), 4.08-4.17 (1H, m), 5.05 (2H, s), 5.12 (1H, d, $J=8.4\text{Hz}$), 5.56 (1H, brs), 7.14-7.35 (20H, m).

Example 22

- 10 benzyl (1S)-1-benzyl-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethylcarbamate



The compound was synthesized in the same manner as in


Example 20.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.21-1.28 (3H, m), 1.81-2.03 (6H, m), 2.34-2.47 (3H, m), 2.99-3.02 (4H, m), 3.90-3.92 (2H, m), 4.06-4.30
 5 (2H, m), 5.00 (2H, s), 5.42 (1H, br), 5.53 (1H, br), 7.04-7.32 (25H, m).

Example 23

benzyl (1S)-2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-1-hydroxymethyl-2-oxoethylcarbamate



10


The compound was synthesized in the same manner as in Example 20.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.21-1.46 (2H, m), 1.82-2.30 (9H, m), 2.50-2.56 (3H, m), 3.15-3.32 (2H, m), 3.66 (1H, dd, $J=11.1\text{Hz}$, 4.7Hz),
 15 3.84 (1H, dd, $J=13.0\text{Hz}$, 5.0Hz), 4.06-4.17 (3H, m), 5.02 (2H, s), 5.98 (1H, br), 6.02 (1H, br), 7.14-7.32 (20H, m).

Example 24

benzyl (1R)-2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-1-methyl-2-oxoethylcarbamate
 20




The compound was synthesized in the same manner as in Example 20.

amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.21-1.32 (6H, m), 1.71-2.09 (6H, m), 2.24-2.52 (3H, m), 2.87-2.93 (2H, m), 3.83-4.16 (4H, m), 5.05 (2H, s), 5.30 (1H, d, $J=6.2\text{Hz}$), 5.55 (1H, t, $J=5.1\text{Hz}$), 7.14-7.34 (20H, m).

Example 25

- 10 (2S)-1-benzyloxycarbonyl-2-(((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)carbonyl)pyrrolidine




The compound was synthesized in the same manner as in Example 20.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.21-1.28 (2H, m), 1.61-2.47 (13H, m), 2.85-2.90 (2H, m), 3.18-3.34 (2H, m), 3.86-4.25 (3H, m), 5.07 (2H, brs), 5.54, 6.07 (1H, 2br), 7.16-7.34 (20H, m).

Example 26


N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-2-thiophenecarboxamide



To a solution of 2,2-diphenyl-5-(4-phenylpiperidino)-
 10 pentanamine (510 mg) synthesized in Reference Example 5D and 2-thiophenecarboxylic acid (176 mg) in acetonitrile (20 ml) was added WSC (265 mg) under ice-cooling, and the mixture was stirred at room temperature for 16 h. Water (200 ml) was added to the reaction mixture. The mixture was extracted with ethyl
 15 acetate (200 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography and eluted with ethyl acetate-ethyl acetate-methanol (20:1) to give the title compound (510 mg).
 20 Recrystallization solvent: diethyl ether.
 melting point: 80-81°C.

Example 27

4-benzyloxy-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-benzamide




The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: ethyl acetate-hexane.

5 melting point: 148-149°C.

Example 28

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-4-phenoxybenzamide hydrochloride




10 The compound was synthesized in the same manner as in Example 26 and converted to hydrochloride.

Recrystallization solvent: ethyl acetate-diethyl ether.

melting point: 199-204°C.

Example 29

15 5-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-2-furancarboxamide




The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: diethyl ether-hexane.

5 melting point: 129-130°C.

Example 30

5-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-2-thiophenecarboxamide




10 The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: ethyl acetate-hexane.

melting point: 147-148°C.

Example 31

15 N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-2-pyrrolicarboxamide




The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: diethyl ether-hexane.

5 melting point: 155-156°C.

Example 32

3-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)benzamide




The compound was synthesized in the same manner as in
10 Example 26.

Recrystallization solvent: diethyl ether-hexane.

melting point: 103-104°C.

Example 33

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-4-
15 biphenylacetamide




The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: ethyl acetate-hexane.

5 melting point: 86-87°C.

Example 34

3-benzyloxy-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-benzamide




10 The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: ethyl acetate-diethyl ether.

melting point: 122-123°C.

Example 35

15 2-benzoylamino-N-(2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)thiazole-4-carboxamide




The compound was synthesized in the same manner as in Example 26.

Recrystallization solvent: ethyl acetate-hexane.

5 melting point: 181-182°C.

Example 36

4-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)benzamide




The compound was synthesized in the same manner as in
10 Example 17.

Recrystallization solvent: ethyl acetate-hexane.

melting point: 140-141°C.

Example 37

4-benzoylamino-N-(2,2-diphenyl-5-(4-phenylpiperidino) -
15 pentyl)benzamide




The compound was synthesized in the same manner as in Example 26.

amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.26–1.43 (3H, m), 1.63–1.92 (5H, m), 2.13–2.41 (5H, m), 2.84–2.91 (2H, m), 4.18 (2H, d, $J=5.8\text{Hz}$), 5.67 (1H, brs), 7.14–7.39 (15H, m), 7.46–7.58 (5H, m), 7.67 (2H, d, $J=8.8\text{Hz}$), 7.84–7.89 (3H, m).

Example 38

- 10 N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-4-(phenylaminocarbonyl)benzamide




The compound was synthesized in the same manner as in Example 26.

- 15 Recrystallization solvent: ethyl acetate-diethyl ether.
melting point: 134–135°C.

Example 39

benzyl 3-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-3-oxopropylcarbamate




5 The compound was synthesized in the same manner as in Example 26.

amorphous powder.

¹H-NMR (CDCl₃) δ: 1.21-1.33 (2H, m), 1.67-2.09 (8H, m), 2.20-2.61 (5H, m), 2.89-2.94 (2H, m), 3.42 (2H, q, J=5.9Hz), 4.00
10 (2H, d, J=6.1Hz), 5.08 (1H, br), 5.11 (2H, s), 6.05 (1H, br), 7.13-7.37 (20H, m).

Example 40

benzyl 4-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-4-oxobutylcarbamate



15

The compound was synthesized in the same manner as in


Example 26.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.21-1.38 (2H, m), 1.67-2.12 (12H, m), 2.24-2.52 (3H, m), 2.86-2.91 (2H, m), 3.10 (2H, q, $J=6.4\text{Hz}$), 4.00
5 (2H, d, $J=5.9\text{Hz}$), 5.05 (2H, s), 5.14 (1H, br), 5.30 (1H, br), 7.17-7.32 (20H, m).

Example 41

benzyl N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-N-methylcarbamate hydrochloride



10


The compound was synthesized in the same manner as in Example 26.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.53-1.68 (3H, m), 2.17-2.28 (2H, m), 2.65 (4H,
15 brs), 2.89 (5H, br s), 3.49-3.55 (2H, m), 3.88-4.00 (4H, m), 5.08 (2H, s), 5.66 (1H, t, $J=5.7\text{Hz}$), 7.15-7.35 (20H, m), 12.02 (1H, br).

Example 42


1-benzyl-3-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)urea



A solution of 2,2-diphenyl-5-(4-phenylpiperidino)-
 pentanamine (110 mg) synthesized in Reference Example 5D and
 benzyl isocyanate (41 mg) in pyridine (5 ml) was stirred at room
 5 temperature for 3 h and concentrated under reduced pressure.
 This residue was partitioned between water (100 ml) and ethyl
 acetate (100 ml). The organic layer was washed with saturated
 brine, dried over anhydrous sodium sulfate and concentrated
 under reduced pressure. The resulting residue was
 10 recrystallized from ethyl acetate-hexane to give the title
 compound (114 mg).
 melting point: 156-157°C.

Example 43

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-
 15 oxoethyl)pyrrole-2-carboxamide




To a solution of tert-butyl 2-((2,2-diphenyl-5-(4-
 phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate (4.0 g)

synthesized in Example 2 in ethyl acetate (60 ml) was added 4N hydrogen chloride-ethyl acetate (150 ml) and the mixture was stirred at room temperature for 1 h. The reaction mixture was concentrated and reprecipitated with methanol-ethyl acetate to give diamine as hydrochloride (4.3 g). To a solution of amine hydrochloride (200 mg), pyrrole-2-carboxylic acid (46 mg) and triethylamine (77 mg) in acetonitrile (15 ml) was added WSC (80 mg) under ice-cooling, and the mixture was stirred at room temperature for 19 h. Water (200 ml) was added to the reaction mixture and the mixture extracted with ethyl acetate (200 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography, eluted with ethyl acetate-ethyl acetate-methanol (5:1), and recrystallized from ethyl acetate-hexane to give the title compound (97 mg).
melting point: 158-159°C.

Example 44

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide



The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: ethyl acetate-hexane.

melting point: 184-185°C.

$$\frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} f(\tau) d\tau = {}^{RL}\mathcal{I}_0^\alpha f(t), \quad {}^{RI}\mathcal{I}_0^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_t^T (T-\tau)^{\alpha-1} f(\tau) d\tau$$
c1ccc(cc1)C2(Cc3ccccc3)CCN(CCN(C(=O)Cc4c[nH]c5ccccc45)CCN6CCCCC6Cc7ccccc7)CC2

amorphous powder.

10 6.93 (1H, s), 7.11–7.41 (19H, m), 7.63–7.67 (1H, m).

Example 46

Chemical structure of compound 10: A complex molecule featuring a central piperidine ring. One side of the piperidine is connected via a methylene chain to a quaternary carbon atom bonded to two phenyl rings and a methylene group. This methylene group is part of an amide chain: -CH₂-C(=O)-NH-CH₂-C(=O)-NH-. The other end of this chain is connected to a benzofuran moiety at its 2-position. The other side of the piperidine ring is connected via a methylene chain to a phenyl ring.


amorphous powder.

2.99 (2H, m), 4.01–4.07 (4H, m), 5.61 (1H, brs), 7.06–7.55 (20H,

m), 7.68-7.72 (1H, m).

Example 47

5-chloro-N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)indole-2-carboxamide



5

The compound was synthesized in the same manner as in Example 43.


amorphous powder.

¹H-NMR (CDCl₃) δ: 1.22-1.29 (2H, m), 1.73-2.35 (13H, m), 2.94-2.98 (2H, m), 3.98-4.17 (4H, m), 5.80 (1H, brs), 6.92 (1H, s), 7.06-7.63 (19H, m), 9.92 (1H, s).

10

Example 48

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)naphthalene-2-carboxamide



15

The compound was synthesized in the same manner as in Example 43.


amorphous powder.

$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix}$

¹H-NMR (CDCl₃) δ: 1.25–1.33 (2H, m), 1.75–2.19 (9H, m), 2.30–2.38 (3H, m), 2.94–2.99 (2H, m), 3.98–4.05 (4H, m), 5.92 (1H, brs), 7.09–7.26 (15H, m), 7.53–7.59 (3H, m), 7.86–7.88 (4H, m), 8.32 (1H, s).

5 Example 49

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-1-hydroxynaphthalene-2-carboxamide



The compound was synthesized in the same manner as in


10 Example 43.

amorphous powder.

¹H-NMR (CDCl₃)δ: 1.25–1.33 (2H, m), 1.75–2.11 (10H, m), 2.38–2.46 (3H, m), 3.11–3.15 (2H, m), 4.00–4.05 (4H, m), 5.89 (1H, brs), 7.08–7.24 (18H, m), 7.45–7.72 (3H, m), 8.39–8.43 (1H, m).

15 **Example 50**

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-3-hydroxynaphthalene-2-carboxamide




The compound was synthesized in the same manner as in Example 43.

amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.26-1.34 (2H, m), 1.85-2.21 (10H, m), 2.42-2.49 (3H, m), 3.12-3.19 (2H, m), 4.01-4.06 (4H, m), 5.80 (1H, brs), 7.08-7.35 (18H, m), 7.48-7.53 (1H, m), 7.68 (1H, d, $J=8.1\text{Hz}$), 7.84 (1H, d, $J=8.4\text{Hz}$).

Example 51

- 10 N-(2-((2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)amino)-2-oxoethyl)-2,2,2-trifluoro-N-phenylacetamide




The compound was synthesized in the same manner as in Example 43.

- 15 Recrystallization solvent: ethyl acetate-hexane.
melting point: 145-146°C.

Example 52

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-3-carboxamide




5 The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: ethyl acetate-hexane.

melting point: 153-154°C.

Example 53

10 N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-4-biphenylcarboxamide



In the same manner as in Example 43, diamine hydrochloride was synthesized. To a solution of diamine hydrochloride (120 mg) and triethylamine (80 mg) in THF (10 ml) was added 4-biphenylcarbonyl chloride (54 mg) under ice-cooling, and the mixture was stirred for 2 h. Saturated aqueous sodium hydrogen carbonate (100 ml) was added and the reaction mixture was extracted with ethyl acetate (100 ml), and after washing


with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography and eluted with ethyl acetate-ethyl acetate-methanol (10:1) to give the title
 5 compound (100 mg).

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.22-1.48 (2H, m), 1.81-2.36 (9H, m), 2.42-2.68 (3H, m), 3.18-3.22 (2H, m), 4.01-4.13 (4H, m), 5.81 (1H, brs), 7.10-7.29 (20H, m), 7.40-7.51 (1H, m), 7.62-7.70 (2H, m),
 10 7.93 (1H, d, $J=8.3\text{Hz}$).

Example 54

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)benzamide




15 The compound was synthesized in the same manner as in Example 53.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.24-1.55 (2H, m), 1.82-1.89 (2H, m), 2.02-2.31 (9H, m), 2.54-2.60 (3H, m), 3.25-3.30 (2H, m), 3.98-4.09
 20 (2H, m), 5.89-5.92 (1H, m), 7.07-7.54 (18H, m), 7.79-7.83 (1H, m), 8.04-8.18 (1H, m).

Example 55

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-3-phenylpropanamide




The compound was synthesized in the same manner as in Example 53.

amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.21-1.38 (2H, m), 1.78-2.17 (9H, m), 2.39-2.53 (4H, m), 2.83-3.03 (4H, m), 3.73 (2H, d, $J=5.3\text{Hz}$), 3.99 (2H, d, $J=5.9\text{Hz}$), 5.67-5.73 (1H, m), 6.69-6.74 (1H, m), 7.13-7.31 (20H, m).

Example 56

- 10 4-nitrobenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in Example 53.

- 15 amorphous powder.

¹H-NMR (CDCl₃)δ: 1.21-1.33 (2H, m), 1.78-2.15 (8H, m), 2.29-2.49 (3H, m), 2.92-2.97 (2H, m), 3.73 (2H, d, J=5.5Hz), 4.01 (2H, d, J=5.9Hz), 5.14 (2H, s), 5.53 (1H, s), 5.79 (1H, s), 7.13-7.32 (15H, m), 7.45 (2H, d, J=8.5Hz), 8.18 (2H, d, J=8.7Hz).

Example 57

4-chlorobenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethylcarbamate




In the same manner as in Example 43, diamine hydrochloride was synthesized. To a solution of diamine hydrochloride (150 mg) and triethylamine (115 mg) in THF (6 ml) was added O-4-chlorobenzyl-O'-4-nitrophenylcarbonate (175 mg) and the mixture was stirred at room temperature for 18 h. Saturated aqueous sodium hydrogen carbonate (100 ml) was added to the reaction mixture, and the mixture was extracted with ethyl acetate (100 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography, eluted with ethyl acetate-ethyl acetate-methanol (10:1) and recrystallized from ethyl acetate-hexane to give the title compound (124 mg).

melting point: 130-131°C.

Example 58

3-chlorobenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethylcarbamate



5


The compound was synthesized in the same manner as in Example 57.

Recrystallization solvent: diethyl ether.

melting point: 125-126°C.

Example 59

2-anilino-N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)acetamide




To a solution of N-(2-((2-((2,2-diphenyl-5-(4-

phenylpiperidino)pentyl)amino)-2-oxoethyl)amino)-2-oxoethyl)-
2,2,2-trifluoro-N-phenylacetamide (150 mg) synthesized in
Example 51 in THF (3 ml) was added 5N aqueous sodium hydroxide
solution (3 ml) and the mixture was stirred at room temperature
for 3 h. After the completion of the reaction, water was added
to the reaction mixture, and the mixture was extracted with
ethyl acetate and washed with saturated brine. The organic
layer was dried over anhydrous magnesium sulfate and
concentrated under reduced pressure. The resulting residue was
applied to silica gel column chromatography and eluted with
ethyl acetate-methanol-aqueous ammonia (50:10:1) to give the
title compound (100 mg).
amorphous powder.

¹H-NMR (CDCl₃)δ: 1.22-1.30 (2H, m), 1.71-2.13 (8H, m), 2.29 (2H, t, J=7.2Hz), 2.35-2.45 (1H, m), 2.91 (2H, t, J=10.9Hz), 3.70 (2H, d, J=5.7Hz), 3.77 (2H, d, J=5.6Hz), 3.90 (2H, d, J=5.7Hz), 4.23 (1H, t, J=5.7Hz), 5.53 (1H, t, J=5.7Hz), 6.50 (2H, d, J=7.6Hz), 6.78 (1H, t, J=7.4Hz), 7.16-7.35 (18H, m).

Example 60

20 2-chlorobenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-
pentyl)amino)-2-oxoethylcarbamate



The compound was synthesized in the same manner as in


Example 57.

Recrystallization solvent: diethyl ether.

melting point: 83-84°C.

Example 61

- 5 4-methoxybenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-
pentyl)amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in
Example 57.

- 10 Recrystallization solvent: ethyl acetate-diethyl ether-hexane.
melting point: 115-116°C.

Example 62

3-methoxybenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-
amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in Example 57.

Recrystallization solvent: ethyl acetate-diethyl ether-hexane.

5 melting point: 96-97°C.

Example 63

2-methoxybenzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethylcarbamate



10 The compound was synthesized in the same manner as in Example 57.


amorphous powder.

¹H-NMR (CDCl₃) δ: 1.21-1.38 (2H, m), 1.70-2.11 (8H, m), 2.26-

2.52 (3H, m), 2.85-2.94 (2H, m), 3.71 (2H, d, J=5.8Hz). 3.83 (3H, m), 3.97 (2H, d, J=5.9Hz), 5.14 (2H, s), 5.41 (1H, brs), 5.62 (1H, brs), 6.87-6.98 (2H, m), 7.14-7.35 (17H, m).

Example 64

- 5 benzyl (1S)-1-benzyl-2-((2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)amino)-2-oxoethylcarbamate




- In the same manner as in Example 43, diamine
 10 hydrochloride was synthesized. To a solution of diamine hydrochloride (200 mg), Z-L-phenylalanine (125 mg) and HOBt (79 mg) in acetonitrile (15 ml) was added WSC (80 mg) at -20°C, and the mixture was stirred at room temperature for 3 days. Water (100 ml) was added to the reaction mixture. The mixture was
 15 extracted with ethyl acetate (100 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography and eluted with ethyl acetate-ethyl acetate-methanol (10:1) to give the title
 20 compound (123 mg).
 amorphous powder.

¹H-NMR (CDCl₃)δ: 1.28-1.45 (2H, m), 1.81-2.41 (8H, m), 2.42-2.69 (2H, m), 2.83-3.31 (4H, m), 3.60-4.12 (5H, m), 4.40-4.59 (1H, m), 5.02 (2H, s), 5.64 (1H, d, J=8.0Hz), 5.84 (1H, brs),

7.07-7.28 (25H, m), 7.58 (1H, brs).

Example 65

2-(((benzylamino) carbonyl) amino)-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl) acetamide



5

In the same manner as in Example 43, diamine hydrochloride was synthesized. A solution of diamine hydrochloride (150 mg) and benzyl isocyanate (42 mg) in pyridine (5 ml) was stirred at room temperature for 3 h. This reaction mixture was partitioned between saturated aqueous sodium hydrogen carbonate (100 ml) and ethyl acetate (100 ml). The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was recrystallized from ethyl acetate-ethyl acetate-diethyl ether to give the title compound (130 mg).

melting point: 149-152°C.


Example 66

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-5-hydroxy-5-phenylpentanamide

20

15 ¹H-NMR (CDCl₃)δ: 1.25–1.36 (2H, m), 1.61–1.77 (7H, m), 1.93–2.12 (6H, m), 2.31–2.51 (3H, m), 2.93–2.99 (2H, m), 3.56 (2H, s), 3.99 (2H, d, J=5.2Hz), 4.59–4.67 (1H, m), 5.16 (1H, s), 7.14–7.32 (20H, m).

20 N-benzyl-N'-(2,2-diphenyl-5-(4-phenylpiperidino)-
pentyl)succinamide




To a solution of N-(2,2-diphenyl-5-(4-phenylpiperidino)-
 pentyl)-3-ethoxycarbonylpropionamide (100 mg) synthesized in
 Example 18 in THF (1 ml) was added 2N aqueous sodium hydroxide
 5 solution (1 ml) and the mixture was stirred at room temperature
 for 1 h. The reaction mixture was diluted with water (100 ml),
 neutralized with 2N hydrochloric acid (1 ml) and extracted with
 ethyl acetate. The organic layer was washed with saturated
 brine, dried over anhydrous sodium sulfate and concentrated
 10 under reduced pressure to give carboxylic acid. To a solution
 of the obtained carboxylic acid and benzylamine (23 mg) and
 HOBt (30 mg) in acetonitrile (5 ml) was added WSC (42 mg) at -
 20°C and the mixture was stirred for 16 h at room temperature.
 To the reaction mixture was added saturated aqueous ammonium
 15 chloride solution (100 ml). The mixture was extracted with
 ethyl acetate (100 ml), and after washing with saturated brine,
 dried over anhydrous sodium sulfate and concentrated under
 reduced pressure. The resulting residue was applied to silica
 gel column chromatography and eluted with ethyl acetate-ethyl
 20 acetate-methanol (10:1-20:3) to give the title compound (100
 mg).

amorphous powder.

¹H-NMR (CDCl₃) δ: 1.21-1.35 (2H, m), 1.72-2.15 (9H, m), 2.27-
 2.50 (6H, m), 2.89-2.94 (2H, m), 3.96 (2H, d, J=6.1Hz), 4.37
 25 (2H, d, J=5.9Hz), 5.41-5.47 (1H, m), 6.51-6.58 (1H, m), 7.14-

[illegible]

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-3-
((phenylacetyl)amino)propanamide




5

melting point: 149-150°C.

Example 69

N-(4-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-4-oxobutyl)benzamide



5


The compound was synthesized in the same manner as in Example 67 from benzyl 4-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-4-oxobutylcarbamate synthesized in Example 40.

Recrystallization solvent: ethyl acetate-hexane

10 melting point: 144-145°C.

Example 70

benzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbonate



15


To a solution of 2-acetoxy-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)acetamide (130 mg) synthesized in Example 19 in THF (5 ml) was added 2N aqueous sodium hydroxide solution (5 ml) and the mixture was stirred at room temperature

for 1 h. The reaction mixture was diluted with water (100 ml) and extracted with ethyl acetate. The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. To a solution of this
 5 residue, triethylamine (272 mg) and DMAP (2 mg) in THF (5 ml) was added carbobenzoxy chloride (448 mg) under ice-cooling, and the mixture was stirred at room temperature for 20 h. To the reaction mixture was added saturated aqueous sodium hydrogen carbonate (100 ml). The mixture was extracted with ethyl
 10 acetate (100 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography and eluted with ethyl acetate-ethyl acetate-methanol (20:1-10:1) to give the title compound (70 mg).
 15 amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.19-1.37 (2H, m), 1.66-2.08 (6H, m), 2.23-2.43 (3H, m), 2.84-2.89 (2H, m), 3.99 (2H, d, $J=5.9\text{Hz}$), 4.52 (2H, s), 5.13 (2H, s), 5.77 (1H, brs), 7.14-7.39 (20H, m).

Example 71

20 N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-5-phenyl-2-thiophenecarboxamide




To a solution of 5-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-2-thiophenecarboxamide (140 mg) synthesized in Example 30, phenyl borate (33 mg) and 2N aqueous
 25

sodium hydrogencarbonate solution (2 ml) in dimethoxyethane (10 ml) was added tetrakis(triphenylphosphine)palladium(0) (28 mg) and the mixture was heated under reflux under a nitrogen atmosphere for 6 h. After cooling, the reaction mixture was
 5 diluted with water (200 ml) and extracted with ethyl acetate (200 ml). The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was applied to silica gel column chromatography, eluted with ethyl acetate-hexane (2:1) and
 10 recrystallized to give the title compound (66 mg).
 melting point: 137-138°C.

Example 72

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-5-phenyl-2-furancarboxamide




15

The compound was synthesized in the same manner as in Example 70 from 5-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)-2-furancarboxamide synthesized in Example 29. Recrystallization solvent: diethyl ether-hexane
 20 melting point: 131-132°C.

Example 73

N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-3-phenylbenzamide




The compound was synthesized in the same manner as in Example 70 from 3-bromo-N-(2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)benzamide synthesized in Example 32.

- 5 Recrystallization solvent: diethyl ether-hexane
melting point: 140-141°C.

Example 74

(2S)-2-amino-N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-3-phenylpropanamide




10

To a solution of benzyl (1S)-1-benzyl-2-((2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl) amino)-2-oxoethyl) amino)-2-oxoethylcarbamate (110 mg) synthesized in Example 64 in ethanol (5 ml) was added 10% palladium carbon (11 mg) and the
15 mixture was stirred at room temperature under a hydrogen atmosphere for 20 h. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure. The

¹H-NMR (CDCl₃) δ: 1.21–1.42 (2H, m), 1.70–2.19 (9H, m), 2.25–2.58 (3H, m), 2.60–3.29 (4H, m), 3.45–4.37 (6H, m), 5.58–5.81 (2H, m), 7.02–7.27 (20H, m), 7.87 (1H, brs).

10 benzyl 2-(N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)-N-methylamino)-2-oxoethylcarbamate




190

methanol-saturated aqueous ammonia (50:10:1). To a solution of this and Z-glycine (84 mg) in acetonitrile (10 ml) was added WSC (76 mg) under ice-cooling, and the mixture was stirred at room temperature for 16 h. Water (200 ml) was added to the
 5 reaction mixture. The mixture was extracted with ethyl acetate (200 ml), and after washing with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was applied to silica gel column chromatography and eluted with ethyl acetate-ethyl
 10 acetate-methanol (20:1) to give the title compound (130 mg). amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.22-1.39 (2H, m), 1.65-2.05 (9H, m), 2.23-2.51 (3H, m), 2.75-2.96 (2H, m), 3.88-3.90 (2H, m), 4.06-4.18 (2H, m), 5.11 (2H, s), 5.85 (1H, s), 7.20-7.35 (20H, m).

15 **Example 76**

benzyl 2-((5-(4-(3-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




To a solution (10 ml) of the compound (0.31 g) obtained
 20 in Reference Example 7D in tetrahydrofuran was added 4-(3-fluorophenyl)piperidine (178 mg). The reaction mixture was stirred overnight at room temperature and the reaction mixture was concentrated under reduced pressure. The residue was dissolved in ethyl acetate, washed with water, dried and
 25 concentrated. The resulting residue was purified by silica gel column chromatography using methanol-ethyl acetate (0:100-5:95). To a solution (10 ml) of the obtained free amine in ethanol was

added 1M ethereal hydrogen chloride (2.0 ml) under ice-cooling and the mixture was stirred at the same temperature for 15 min. The reaction mixture was concentrated under reduced pressure, and the residue was crystallized from ethyl acetate-hexane to
 5 give the title compound (0.29 g).
 amorphous powder.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.23-1.33 (2H, m), 1.43-1.47 (2H, m), 1.97-2.01 (2H, m), 2.57-2.90 (7H, m), 3.84-4.13 (6H, m), 5.02 (2H, s), 5.86 (1H, s), 6.73 (1H, s), 6.90-7.61 (19H, m), 11.37 (1H, brs).

10 **Example 77**

benzyl 2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in
 15 Example 76 from the compound obtained in Reference Example 7D.
 amorphous powder.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.25 (2H, brs), 1.47 (2H, s), 1.96-2.00 (2H, m), 2.53-3.11 (7H, m), 3.81-3.98 (6H, m), 5.01 (2H, s), 5.86 (1H, s), 6.85 (1H, s), 6.98-7.43 (19H, m), 11.30 (1H, brs).

20 **Example 78**

benzyl 2-((5-(4-(4-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

5 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.46 (2H, brs), 1.69 (2H, brs), 1.93-1.97 (2H, m), 2.54-2.89 (7H, m), 3.78-3.98 (6H, m), 5.02 (2H, s), 5.86 (1H, s), 6.83 (1H, s), 6.96-7.38 (19H, m), 11.31 (1H, brs).

Example 79

benzyl 2-((5-(4-(3-methoxyphenyl)piperidino)-2,2-
10 diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

15 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.50 (2H, brs), 1.64 (2H, s), 1.98-2.02 (2H, m), 2.57-2.85 (5H, m), 2.91 (2H, brs), 5.03 (2H, s), 5.89 (1H, s), 6.74-6.92 (4H, m), 7.18-7.37 (18H, m), 11.33 (1H, brs).

Example 80

benzyl 2-((5-(4-(2-methoxyphenyl)piperidino)-2,2-
20 diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.26 (2H, brs), 1.62 (2H, s), 1.97-1.99 (2H, brs), 2.57-2.65 (4H, m), 2.89-2.90 (2H, m), 3.11-3.16 (1H, m), 3.81 (3H, s), 3.89-3.98 (4H, m), 5.01 (2H, s), 5.91 (1H, s), 6.83-6.97 (3H, m), 7.16-7.38 (19H, m), 11.23 (1H, brs).

Example 81

benzyl 2-((5-(4-(4-methoxyphenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride



The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D.

Recrystallization solvent: ethyl acetate


melting point: 119-120°C

¹H-NMR(CDCl₃)δ: 1.49 (2H, brs), 1.67 (2H, s), 1.99-2.06 (2H, m), 2.58-2.80 (5H, m), 2.90 (2H, brs), 3.80 (3H, s), 3.91-4.03 (4H, m), 5.03 (2H, s), 5.91 (1H, s), 6.85-6.90 (3H, m), 7.12-7.63 (19H, m), 11.31 (1H, brs).

•

Example 84

benzyl 2-((2,2-diphenyl-5-piperidinopentyl)amino)-2-oxoethylcarbamate hydrochloride




5 The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.33-1.44 (2H, m), 1.67 (3H, s), 1.79-1.89 (2H, m), 2.33-2.54 (5H, m), 2.85-2.87 (2H, m), 3.65-3.68 (2H, m),
 10 3.92-3.99 (4H, m), 5.05 (2H, s), 5.91 (1H, s), 6.91-6.93 (1H, s), 7.18-7.41 (15H, m), 10.99 (1H, brs).

Example 85

benzyl 2-((5-(4-benzylpiperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride



15


The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.44 (2H, s), 1.71-1.83 (3H, m), 2.15-2.28 (2H, m), 2.35-2.64 (6H, m), 2.83-2.84 (2H, m), 3.69-3.72 (2H, m),
 20 3.89-3.98 (4H, m), 5.06 (2H, s), 5.90 (1H, s), 6.90 (1H, s),

7.09-7.63 (21H, m), 11.10 (1H, brs).

Example 86

benzyl 2-((5-(4-(3-fluorobenzyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride



5


The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.42 (2H, s), 1.63-1.80 (4H, m), 2.15-2.27 (2H, m), 2.38-2.62 (6H, m), 2.81 (2H, s), 3.69-3.72 (2H, m), 3.91-3.96 (4H, m), 5.04 (2H, s), 5.86 (1H, s), 6.78-6.93 (4H, m), 7.16-7.61 (17H, m), 11.10 (1H, brs).

Example 87

benzyl 2-((5-(4-(2-fluorobenzyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride

15




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.41 (2H, s), 1.76-1.80 (4H, m), 2.14-2.27 (2H, m), 2.36-2.66 (4H, m), 2.81 (2H, s), 3.66-3.70 (2H, m), 3.91-

3.96 (4H, m), 5.04 (2H, s), 5.89 (1H, s), 7.00-7.35 (23H, m), 11.07 (1H, brs).

Example 88

benzyl 2-((5-(4-(4-fluorobenzyl)piperidino)-2,2-
5 diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

10 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.42 (2H, s), 1.63-1.79 (4H, m), 2.13-2.25 (2H, m), 2.33-2.59 (6H, m), 2.81 (2H, s), 3.68-3.90 (2H, m), 3.90-3.96 (4H, m), 5.04 (2H, s), 5.87 (1H, s), 6.93-7.35 (19H, m), 11.10 (1H, brs).

Example 89

15 benzyl 2-((5-(4-(2,4-difluorobenzyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

20 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.42-1.44 (2H, m), 1.69-1.80 (4H, m), 2.16-2.29

(2H, m), 2.37-2.64 (6H, m), 2.84 (2H, s), 3.69-3.72 (2H, m), 3.92-4.01 (4H, m), 5.06 (2H, s), 5.89 (1H, s), 6.75-6.87 (4H, m), 7.05-7.37 (14H, m), 11.11 (1H, brs).

Example 90

5 benzyl 2-((5-(4-(4-cyanobenzyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D.
10 amorphous powder.

¹H-NMR(CDCl₃)δ: 1.44 (2H, s), 1.66-1.79 (4H, m), 2.21-2.53 (6H, m), 2.67-2.70 (2H, m), 2.84 (2H, s), 3.72-3.75 (2H, m), 3.92-3.98 (4H, m), 5.06 (2H, s), 5.85 (1H, s), 6.86 (1H, s), 7.18-7.37 (17H, m), 7.58-7.63 (2H, m), 11.16 (1H, brs).

15 **Example 91**

benzyl 2-((5-(4-anilinopiperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D.
20 Recrystallization solvent: ethyl acetate

melting point: 134-136°C

¹H-NMR(CDCl₃)δ: 1.40 (2H, s), 2.26-3.04 (10H, m), 3.53-3.94 (8H, m), 4.99 (2H, s), 5.88 (1H, s), 6.63 (1H, s), 6.91 (1H, s), 7.15-7.44 (19H, m), 10.30 (1H, brs), 10.87 (1H, brs).

5 Example 92

benzyl 2-((2,2-diphenyl-5-(1,2,3,4-tetrahydroisoquinolin-2-yl)pentyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in
10 Example 76 from the compound obtained in Reference Example 7D.
amorphous powder.

¹H-NMR(CDCl₃)δ: 1.26-1.68 (4H, m), 2.52-2.65 (2H, m), 2.99 (3H, s), 3.36 (2H, brs), 3.66 (1H, brs), 3.87-4.00 (4H, m), 4.67-4.76 (1H, m), 5.00 (2H, s), 5.90 (1H, s), 6.84 (1H, s), 7.13-
15 7.37 (19H, m), 11.97 (1H, brs).

Example 93

benzyl 2-((5-(isoindolin-2-yl)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride



20 The compound was synthesized in the same manner as in
Example 76 from the compound obtained in Reference Example 7D.


Recrystallization solvent: ethyl acetate-hexane

melting point: 121-123°C

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.53-1.62 (6H, m), 2.58-2.64 (2H, m), 3.12-3.13 (2H, m), 3.92-4.01 (4H, m), 4.16-4.22 (2H, m), 5.00 (2H, s),
 5 5.11-5.18 (2H, m), 5.86 (1H, s), 6.77 (1H, s), 7.17-7.35 (15H, m), 12.39 (1H, brs).

Example 94

benzyl 2-((5-(N-methyl-N-phenethylamino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride




10

The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.36 (1H, brs), 1.68 (2H, s), 2.16-2.20 (1H, m),
 15 2.81-2.84 (4H, m), 3.06-3.33 (5H, m), 3.75-4.04 (3H, m), 4.21-4.28 (1H, m), 4.98-5.08 (2H, m), 5.88 (1H, s), 6.81 (1H, s), 7.20-7.63 (20H, m), 11.52 (1H, brs).

Example 95

benzyl 2-((5-(N-benzyl-N-methylamino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate hydrochloride
 20




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.29-1.36 (1H, m), 1.59-1.70 (1H, m), 2.05-2.13 (1H, m), 2.68-2.89 (6H, m), 3.74-4.06 (4H, m), 4.21-4.27 (1H, m), 4.39-4.44 (1H, m), 5.04 (2H, s), 5.94 (1H, s), 6.88 (1H, s), 7.18-7.55 (20H, m), 11.55 (1H, brs).

Example 96

benzyl 2-((2,2-diphenyl-4-(4-phenylpiperidino)butyl)amino)-2-oxoethylcarbamate hydrochloride




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR(CDCl₃)δ: 1.61 (1H, s), 1.94-2.04 (2H, m), 2.61 (4H, s), 2.81-2.89 (4H, m), 3.69 (2H, s), 3.93-4.05 (4H, m), 5.08 (2H, s), 6.12 (1H, s), 6.43 (1H, s), 7.13-7.35 (20H, m), 11.87 (1H, brs).

Example 97

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)benzothiophene-2-carboxamide




The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: diethyl ether-hexane.

5 melting point: 110-113°C.

Example 98

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-4-carboxamide




10 The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: ethyl acetate-diethyl ether.

melting point: 215-218°C.

Example 99

15 N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-5-carboxamide




The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: ethyl acetate-diethyl ether.

5 melting point: 156-160°C.

Example 100

N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-6-carboxamide




10 The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: ethyl acetate-diethyl ether.

melting point: 149-152°C.

Example 101

15 N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-7-carboxamide




The compound was synthesized in the same manner as in Example 43.

Recrystallization solvent: ethyl acetate-diethyl ether.

5 melting point: 166-168°C.

Example 102

tert-butyl 2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate



10 The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 13D. amorphous powder.


¹H-NMR(CDCl₃)δ: 1.21-1.28 (2H, m), 1.40 (9H, s), 1.68-2.02 (8H, m), 2.24-2.31 (2H, m), 2.37-2.53 (1H, m), 2.86-2.91 (2H, m),
 15 3.63 (2H, d, J=5.9Hz), 3.93 (2H, d, J=5.9Hz), 5.25 (1H, t, J=5.5Hz), 5.89 (1H, br), 7.08-7.29 (13H, m).

Recrystallization solvent: ethyl acetate-diethyl ether.

melting point: 143-145°C.

Example 105

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-
5 amino)-2-oxoethyl)-5-chloroindole-2-carboxamide




The compound was synthesized in the same manner as in
Example 43 from the compound obtained in Example 102.

Recrystallization solvent: ethyl acetate-diethyl ether.

melting point: 159-161°C.

Example 106

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-
pentyl)amino)-2-oxoethyl)-1-methylindole-2-carboxamide



The compound was synthesized in the same manner as in


Example 43 from the compound obtained in Example 102.

Recrystallization solvent: diethyl ether-hexane.

melting point: 107-110°C.

Example 107

- 5 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-
pentyl)amino)-2-oxoethyl)-5-fluoroindole-2-carboxamide



The compound was synthesized in the same manner as in
Example 43 from the compound obtained in Example 102.

- 10 Recrystallization solvent: ethyl acetate-diethyl ether-hexane.
melting point: 133-135°C.

Example 108

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-
amino)-2-oxoethyl)-5-methoxyindole-2-carboxamide

Recrystallization solvent: ethyl acetate-hexane.

Example 109


Oc1ccc2c(c1)c(c[nH]2)C(=O)NCC(=O)NCC(Cc3ccc(Cl)cc3)(Cc4ccc(Cl)cc4)CCCN5CCCCC5c6ccccc6

Recrystallization solvent: ethyl acetate-hexane.

melting point: 154-157°C.

Example 110

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-2-phenoxyacetamide



5


The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 102.

Recrystallization solvent: diethyl ether-hexane.

melting point: 143-145°C.

10 Example 111

benzyl 2-((5-(4-phenoxy-piperidino)-2,2-diphenylpentyl)amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in

Example 76 from the compound obtained in Reference Example 7D.
amorphous powder.

¹H-NMR (CDCl₃) δ: 1.24 (1H, m), 1.45 (2H, brs), 1.78 (4H, s), 2.07-2.25 (2H, m), 2.53-2.91 (7H, m), 3.56-3.59 (1H, m), 3.82-3.97 (4H, m), 4.59-4.61 (1H, m), 5.03 (2H, s), 5.83-5.89 (1H, m), 6.83-7.01 (4H, m), 7.17-7.34 (16H, m).

Example 112

benzyl 2-((2,2-diphenyl-5-(4-piperidinopiperidino)pentyl)-
amino)-2-oxoethylcarbamate




10

The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

¹H-NMR (CDCl₃) δ: 1.28-1.48 (2H, m), 1.81-1.87 (4H, m), 2.04-2.41 (10H, m), 2.62-2.94 (5H, m), 3.11-3.18 (2H, m), 3.40-3.49 (2H, m), 3.77-3.96 (6H, m), 5.04 (2H, m), 5.78 (1H, s), 6.80 (1H, s), 7.18-7.34 (15H, m).

Example 113

benzyl 2-((5-(4-(2-fluorophenyl)-1,2,5,6-tetrahydropyridin-1-
20 yl)-2,2-diphenylpentyl) amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 7D. amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.63 (4H, s), 2.53–2.81 (4H, m), 3.06–3.23 (4H, m), 3.82–3.98 (4H, m), 4.23–4.29 (1H, m), 5.06 (2H, s), 5.89 (2H, s), 6.84–7.71 (19H, m).

Example 114

tert-butyl 2-((5-(4-(2-fluorophenyl)piperidino)-2,2-
10 diphenylpentyl)amino)-2-oxoethylcarbamate



The compound was synthesized in the same manner as in Example 1 from tert-butyl 2-((5-hydroxy-2,2-diphenylpentyl)-


amino)-2-oxoethylcarbamate.

amorphous powder.

¹H-NMR (CDCl₃) δ: 1.39 (9H, s), 1.66-1.76 (5H, m), 1.90-2.30 (7H, m), 2.68-2.89 (3H, m), 3.63-3.67 (2H, m), 3.97-4.01 (2H, m),
5 4.92 (1H, s), 5.59 (1H, s), 7.06-7.42 (14H, m).

Example 115

N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)-amino)-2-oxoethyl)indole-2-carboxamide




10 The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 114.

amorphous powder.

¹H-NMR (CDCl₃) δ: 1.45-1.49 (2H, m), 1.65 (5H, s), 2.05-2.10 (1H, m), 2.57-2.96 (6H, m), 3.91-4.02 (4H, m), 4.16-4.19 (2H, m),
15 4.00-4.02 (1H, m), 5.91 (1H, s), 6.95-7.46 (17H, m), 7.60-7.62 (1H, m), 9.04 (1H, s), 7.79-7.82 (1H, m), 9.04 (1H, s).

Example 116

N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)-amino)-2-oxoethyl)-1-methylindole-2-carboxamide




The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 114.

amorphous powder.

¹H-NMR (CDCl₃)δ: 1.25-1.81 (5H, m), 2.03-2.07 (1H, m), 2.34-3.33 (7H, m), 3.90-4.00 (7H, m), 4.16-4.19 (2H, m), 5.89 (1H, s), 6.91-7.47 (17H, m), 7.58-7.66 (1H, m), 7.77-7.82 (1H, m).

Example 117

5-chloro-N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethyl)-1-methylindole-2-carboxamide



The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 114.


amorphous powder.

¹H-NMR (CDCl₃)δ: 1.27-1.83 (7H, m), 2.06-2.10 (1H, m), 2.39-

3.17 (7H, m), 3.73-4.19 (7H, m), 5.86 (1H, s), 6.94-7.78 (19H, m).

Example 118

5-chloro-N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)amino)-2-oxoethyl)indole-2-carboxamide




The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 114.
amorphous powder.

¹H-NMR (CDCl₃)δ: 1.33-1.61 (7H, m), 2.07-2.10 (1H, m), 2.38-3.15 (6H, m), 3.83-4.15 (5H, m), 5.81 (1H, m), 6.91-7.73 (19H, m), 9.12 (1H, s).

Example 119

tert-butyl 2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 13D. amorphous powder.

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.26–1.39 (11H, m), 1.85–2.56 (10H, m), 2.91–3.30 (3H, m), 3.73–3.94 (4H, m), 5.82 (1H, s), 6.97–7.43 (13H, m).

Example 120

N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-
10 piperidino)pentyl)amino)-2-oxoethyl)-5-fluoroindole-2-carboxamide




The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 119.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.14-1.84 (6H, m), 1.96-3.19 (8H, m), 3.65-4.15 (5H, m), 5.81 (1H, s), 6.90-7.65 (17H, m), 9.17 (1H, s).

Example 121

5 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-methoxyindole-2-carboxamide




The compound was synthesized in the same manner as in
10 Example 43 from the compound obtained in Example 119.
amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.23-1.93 (6H, m), 2.07-3.14 (8H, m), 3.63-4.23 (8H, m), 5.84 (1H, s), 6.89-7.72 (17H, m), 9.04 (1H, s).

Example 122

15 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide




The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 119.

amorphous powder

¹H-NMR (CDCl₃)δ: 1.32-1.93 (6H, m), 2.04-3.10 (8H, m), 3.63-4.23 (8H, m), 5.84 (1H, s), 6.89-7.72 (17H, m), 9.04 (1H, s).

Example 123

tert-butyl 2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate



10


The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 19D.
amorphous powder.

¹H-NMR (CDCl₃)δ: 1.19-1.34 (2H, m), 1.40 (9H, s), 1.66-1.80 (4H,

m), 1.89-2.05 (4H, m), 2.25-2.30 (2H, m), 2.39-2.50 (1H, m), 2.86-2.90 (2H, m), 3.66 (2H, d, J =6.0Hz), 3.94 (2H, d, J =6.0Hz), 4.91 (1H, s), 5.67-5.71 (1H, s), 6.97-7.02 (3H, m), 7.12-7.31 (10H, m).

5 Example 124

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)indole-2-carboxamide




The compound was synthesized in the same manner as in
 10 Example 43 from the compound obtained in Example 123.
 amorphous powder.

¹H-NMR (CDCl₃)δ: 1.24-1.27 (2H, m), 1.77-2.09 (8H, m), 2.29-
 2.48 (3H, m), 2.93-2.96 (2H, m), 3.96-4.02 (4H, m), 5.85-5.88
 (1H, m), 6.82-6.91 (4H, m), 6.96 (1H, s), 7.07-7.35 (12H, m),
 15 7.45 (1H, d, J=8.1Hz), 7.68 (1H, d, J=8.0Hz), 9.40 (1H, s).

Example 125

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)-5-chloroindole-2-carboxamide




The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 123.
amorphous powder.

- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.25–1.33 (2H, m), 1.81–2.07 (8H, m), 2.19–2.46 (3H, m), 2.98–3.01 (2H, m), 3.96–4.02 (4H, m), 6.83–6.91 (5H, m), 7.03–7.31 (11H, m), 7.38 (1H, d, $J=8.8\text{Hz}$), 7.65 (1H, s), 9.44 (1H, s).

Example 126

- 10 tert-butyl 2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethylcarbamate




The compound was synthesized in the same manner as in Example 76 from the compound obtained in Reference Example 19D.

amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.35 (9H, s), 1.45-1.47 (2H, m), 2.00-2.04 (2H, m), 2.39-3.23 (1H, m), 3.82-3.92 (7H, m), 5.77-5.81 (1H, m), 6.84-7.06 (6H, m), 7.19-7.27 (6H, m).

5 **Example 127**

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide




The compound was synthesized in the same manner as in
10 Example 43 from the compound obtained in Example 126.
amorphous powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.23-1.28 (2H, m), 1.76-2.36 (10H, m), 2.89-2.99 (3H, m), 3.79 (3H, s), 3.96-4.02 (4H, m), 5.89-5.92 (1H, m), 6.81-7.35 (16H, m), 7.45 (1H, d, $J=8.3\text{Hz}$), 7.68 (1H, d, $J=8.0\text{Hz}$), 9.46 (1H, s).
15

Example 128

N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-chloroindole-2-carboxamide




The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 126.
amorphous powder.


- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.26-1.30 (2H, m), 1.78-1.87 (3H, m), 2.02-2.14 (5H, m), 2.35-2.71 (3H, m), 2.88-3.03 (3H, m), 3.81 (3H, s), 3.98-4.03 (3H, m), 5.92-5.96 (1H, m), 6.83-6.95 (7H, m), 7.06-7.21 (6H, m), 7.38 (2H, d, $J = 8.8\text{Hz}$), 7.52 (1H, s), 7.63-7.66 (1H, s), 9.77 (1H, s).

10 **Example 129**

tert-butyl 2-((2,2-bis(4-fluorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethylcarbamate



The compound was synthesized in the same manner as in



The compound was synthesized in the same manner as in Example 43 from the compound obtained in Example 129.
amorphous powder.

5 $^1\text{H-NMR}$ (CDCl_3) δ : 1.23-1.33 (2H, m), 1.67-2.37 (10H, m), 2.80-2.87 (1H, m), 3.02-3.05 (2H, m), 3.93-4.02 (4H, m), 5.84-5.88 (1H, m), 6.73-6.94 (4H, m), 6.97-7.47 (12H, m), 7.64 (1H, s), 9.69 (1H, s).

The following compounds can be synthesized in completely
10 the same manner.

N-(2-((2,2-bis(4-methylphenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)indole-2-carboxamide.

N-(2-((2,2-bis(4-methylphenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)-5-chloroindole-2-carboxamide.

15 N-(2-((2,2-bis(4-methylphenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide.

N-(2-((2,2-bis(4-methylphenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-chloroindole-2-carboxamide.

20 N-(2-((2,2-bis(4-methylphenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide.

N-(2-((2,2-bis(4-methylphenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl)amino)-2-oxoethyl)-5-chloroindole-2-carboxamide.

Formulation Example 1A

(1) compound of Reference Example IIA-45	10.0 g
(2) lactose	60.0 g
(3) cornstarch	35.0 g
5 (4) gelatin	3.0 g
(5) magnesium stearate	2.0 g

Using 10 wt% aqueous gelatin solution (30 ml) (3.0 g as gelatin), a mixture of the compound (10.0 g) obtained Reference Example IIA-45, lactose (60.0 g) and cornstarch (35.0 g) was
 10 granulated by passing through a 1 mm mesh sieve. The granules were dried at 40°C and passed through the sieve again. The obtained granules were mixed with magnesium stearate (2.0 g) and compressed. The obtained core tablets were coated with a sugar coating of aqueous suspension containing sucrose,
 15 titanium dioxide, talc and gum arabic. The coated tablets were glazed with bee wax to give 1000 coated tablets.

Formulation Example 2A

(1) compound of Reference Example IIA-45	10.0 g
(2) lactose	70.0 g
20 (3) cornstarch	50.0 g
(4) soluble starch	7.0 g
(5) magnesium stearate	3.0 g

Using an aqueous solution (70 ml) of soluble starch (7.0 g as soluble starch), the compound (10.0 g) obtained in
 25 Reference Example IIA-45 and magnesium stearate (3.0 g) were granulated, dried and mixed with lactose (70.0 g) and cornstarch (50.0 g). The mixture was compressed to give 1000 tablets.

Formulation Example 1B

30 (1) compound of Reference Example 4B-2	10.0 g
(2) lactose	60.0 g
(3) cornstarch	35.0 g
(4) gelatin	3.0 g

(5) magnesium stearate 2.0 g

Using a 10 wt% aqueous gelatin solution (30 ml) (3.0 g as gelatin), a mixture of the compound (10.0 g) obtained Reference Example 4B-2, lactose (60.0 g) and cornstarch (35.0 g) was
 5 granulated by passing through a 1 mm mesh sieve. The granules were dried at 40°C and passed through the sieve again. The obtained granules were mixed with magnesium stearate (2.0 g) and compressed. The obtained core tablets were coated with a sugar coating of aqueous suspension containing sucrose,
 10 titanium dioxide, talc and gum arabic. The coated tablets were glazed with bee wax to give 1000 coated tablets.

Formulation Example 2B

(1) compound of Reference Example 4B-2	10.0 g
(2) lactose	70.0 g
15 (3) cornstarch	50.0 g
(4) soluble starch	7.0 g
(5) magnesium stearate	2.0 g

Using an aqueous solution (70 ml) of soluble starch (7.0 g as soluble starch), the compound (10.0 g) obtained in
 20 Reference Example 4B-2 and magnesium stearate (3.0 g) were granulated, dried and mixed with lactose (70.0 g) and cornstarch (50.0 g). The mixture was compressed to give 1000 tablets.

Formulation Example 1C

25 (1) compound of Reference Example 5C-3	10.0 g
(2) lactose	60.0 g
(3) cornstarch	35.0 g
(4) gelatin	3.0 g
(5) magnesium stearate	2.0 g

30 Using a 10 wt% aqueous gelatin solution (30 ml) (3.0 g as gelatin), a mixture of the compound (10.0 g) obtained Reference Example 5C-3, lactose (60.0 g) and cornstarch (35.0 g) was granulated by passing through a 1 mm mesh sieve. The granules

give 1000 coated tablets.

Formulation Example 2D

(1) compound of Example 1	10.0 g
(2) lactose	70.0 g
5 (3) cornstarch	50.0 g
(4) soluble starch	7.0 g
(5) magnesium stearate	2.0 g

Using an aqueous solution (70 ml) of soluble starch (7.0 g as soluble starch), the compound (10.0 g) obtained in Example 10 1 and magnesium stearate (3.0 g) were granulated, dried and mixed with lactose (70.0 g) and cornstarch (50.0 g). The mixture was compressed to give 1000 tablets.

Reference Example 1E Amplification of rat SLC-1 receptor cDNA by PCR using cDNA derived from rat brain

15 Using poly (A)⁺ RNA derived from rat brain (Clontech) as a template and a random primer, reverse-transcription reaction was carried out. For the reverse-transcription reaction, a reagent of TaKaRa RNA PCR ver. 2 kit was used. Using this reverse-transcription product as a template and synthetic DNA 20 primers of SEQ Nos:1 and 2, amplification was performed by the PCR method. The synthetic DNA primers were constructed such that the gene in the region to be translated into the receptor protein could be amplified, during which restriction enzyme recognizing sequences of restriction enzyme Sal I and 25 restriction enzyme Spe I were added to the 5' side and 3' side, respectively, so that a base sequence recognized by the restriction enzyme Sal I would be added to the 5' side of the gene and a base sequence recognized by the restriction enzyme Spe I would be added to the 3' side of the gene. The 30 composition of the reaction mixture was cDNA template 5 μ l, each synthetic DNA primer 0.4 μ M, 0.25 mM dNTPs, pfu (Stratagene) DNA polymerase 0.5 μ l and buffer annexed to the enzyme, with the total reaction volume of 50 μ l. For

amplification cycle, Thermal Cycler (Perkins Elmer) was used. After heating at 94°C for 60 seconds, a cycle of heating at 94°C for 60 seconds, at 60°C for 30 seconds, and at 72°C for 150 seconds was repeated 35 times, and the mixture was finally
5 reacted at 72°C for 10 minutes. The amplified product was confirmed by ethidium bromide staining after 0.8% agarose gel electrophoresis.

Reference Example 2E Subcloning of PCR product to plasmid vector and confirmation of amplified cDNA sequence by decoding
base sequence of insert cDNA

The reaction product after PCR conducted in Reference Example 1E was separated using 0.8% low melting point agarose gel and the band was excised with a razor, and subjected to minimization, phenol extraction, phenol-chloroform extraction and ethanol precipitation to recover DNA. According to the direction of PCR-Script™ Amp SK(+) cloning kit (Stratagene), the recovered DNA was subcloned to plasmid vector pCR-Script Amp SK(+). This was introduced into Escherichia coli XL-1 Blue (Stratagene) to allow transformation, after which clones containing cDNA insert fragment were selected in an LB agar medium containing ampicillin and X-gal, separated using a sterile toothpick for white clones to give transformant E. coli XL-1 Blue/rat SLC-1. The respective clones were cultured overnight in an LB medium containing ampicillin, and using QIA prep8 mini prep (QIAGEN), plasmid DNAs were prepared. A part of the prepared DNAs was cleaved with restriction enzymes Sal I and Spe I to confirm the size of the inserted receptor cDNA fragment. The reaction for determination of the base sequence was carried out using DyeDeoxy Terminator Cycle Sequence Kit (Perkins Elmer), and decoded using a fluorescence automatic DNA sequencer. The sequences of the obtained three clones were analyzed and confirmed to be identical with the gene sequence consisting of a cDNA sequence (Lakaye, B. et al. Biochim.

Biophys. Acta, Vol. 1401, pp. 216-220 (1998), accession No. AF08650) encoding rat SLC-1 protein (SEQ No:3) whose full length sequence had been reported, a Sal I recognizing sequence added on the 5' side and a Spe I recognizing sequence added on
 5 the 3' side (SEQ No:4).

Reference Example 3E Preparation of rat SLC-1 expression CHO cell

From a clone of E. coli transformed with a plasmid incorporating a gene encoding a full length amino acid sequence
 10 of rat brain derived SLC-1 and having a Sal I recognizing sequence added on the 5' side and a Spe I recognizing sequence added on the 3' side, whose sequence was confirmed in Reference Example 2E, a plasmid was prepared using Plasmid Midi Kit (QIAGEN) and cleaved with restriction enzymes Sal I and Spe I
 15 to excise an insert. The insert DNA was recovered by excising, after electrophoresis, from agarose gel with a razor and applying minimization, phenol extraction, phenol-chloroform extraction and ethanol precipitation. This insert DNA was added to animal cell expression vector plasmid pAKKO-111H
 20 (vector plasmid identical with pAKKO1.11H described in Hinuma, S. et al. Biochim. Biophys. Acta, Vol. 1219, pp. 251-259 (1994)) cleaved with Sal I and Spe I and ligated using T4 ligase (Takara Shuzo Co.) to construct protein expression plasmid pAKKO-SLC-1.

25 E. coli DH5 (TOYOBO) transformed with pAKKO-SLC-1 was cultured and plasmid DNA of pAKKO-SLC-1 was prepared using Plasmid Midi Kit (QIAGEN). This was introduced into CHO dhfr⁻ cell using CellPfect Transfection Kit (Amersham Pharmacia Biotech) and in accordance with the attached protocol. DNA (10
 30 µg) was prepared into a coprecipitation suspension with calcium phosphate and added into a 10 cm dish inoculated with 5 x 10⁵ or 1 x 10⁶ CHO dhfr⁻ cells 24 hours before. The cells were cultured in an MEM α medium containing 10% fetal bovine serum

for one day, passaged and cultured in a nucleic acid-free MEM α medium (selection medium) containing 10% dialyzed fetal bovine serum. 56 clones of transformed cell colonies, which were SLC-1 expression CHO cells grown in the selection medium, were
 5 selected.

Reference Example 4E Selection of CHO/SLC-1 cell line with high expression amount of full length rat SLC-1 receptor protein mRNA

The expression amount of full length rat SLC-1-receptor
 10 protein mRNA by 56 clones of CHO/SLC-1 cell line established in Reference Example 3E was measured as in the following using Cytostar T Plate (Amersham Pharmacia Biotech) and in accordance with the attached protocol. Each clone of the CHO/SLC-1 cell
 15 line was inoculated to each well of Cytostar T Plate at 2.5×10^4 , cultured for 24 hours and fixed with 10% formalin. After 0.25% Triton X-100 was added to each well to enhance permeability of the cells, riboprobe of ^{35}S labeled SEQ No:5 was added for hybridization. RNase A (20 mg/ml) was added to each well to digest free riboprobe. The plate was washed thoroughly
 20 and the radioactivity of the hybridized riboprobe was measured on Topcounter. The cell strain having high radioactivity showed higher expression amount of mRNA. Of the three clones showing high mRNA expression amount, particularly clone No. 44 was used mainly.

25 **Reference Example 5E** Isolation of plasmid containing human SLC-1 cDNA

According to the manual attached to Genetrappor cDNA positive selection system (GIBCOBRL) and using phage F1 endonuclease, nick was inserted into cDNA derived from human
 30 fetal brain library (SUPERSRIPTTM cDNA Library; GIBCOBRL) and digested with Escherichia coli exonuclease III to prepare a single strand cDNA derived from human fetal brain library.

Using Terminal Deoxynucleotidyl Transferase, biotin-14-

dCTP was added to the 3' terminal of the synthetic oligonucleotide (corresponding to 1434-1451 of accession No. U71092) of SEQ No:6 prepared based on the report of Kolakowski Jr. et al. (Kolakowski Jr., et al (1996) FEBS Lett. Vol. 398, pp. 253-258), whereby biotinylated oligonucleotide was prepared. The composition of the reaction mixture and reaction time followed the manual.

Single strand cDNA library derived from human fetal brain 4 µg was kept at 95°C for 1 min and rapidly cooled on ice. Biotinylated oligonucleotide (20 ng) was added and the mixture was hybridized in the accompanying hybridization buffer at 37°C for 1 hr. Streptavidin beads were added and single strand cDNA derived from human fetal brain hybridized to biotinylated oligonucleotide was isolated using MAGNA-SEP Magnetic Particle Separator (GIBCOBRL). Using synthetic oligonucleotide (50 ng, corresponding to 1011-1028 of accession No. U71092) of SEQ No:7 prepared according to the report of Kolakowski Jr. et al. (Kolakowski Jr., et al (1996) FEBS Lett. Vol. 398, pp. 253-258) as a primer, a complementary chain was synthesized according to the manual to give a double strand plasmid.

Reference Example 6E Determination of base sequence of plasmid containing isolated human SLC-1 cDNA.

The plasmid obtained in Reference Example 5E was introduced into ELECTROMAX™ DH10B™ Cells by electroporation method to allow transformation, after which clones containing cDNA insert fragment were selected in an LB agar medium containing ampicillin and X-gal and separated using a sterile toothpick for white clones to give transformant E. coli DH10B/hSLC-1. The respective clones were cultured overnight in an LB medium containing ampicillin, and using QIA prep8 mini prep (QIAGEN), the plasmid DNA was purified. The reaction for determination of the base sequence was carried out using DyeDeoxy Terminator Cycle Sequence Kit (Perkins Elmer), and

decoded using a fluorescence automatic DNA sequencer. As a result, the sequence depicted in SEQ No:8 was obtained. The amino acid sequence (SEQ No:9) encoded by the obtained base sequence was different from the human SLC-1 amino acid sequence as the sequence deduced from rat SLC-1 based on the human chromosomal DNA sequence (accession number:Z86090) containing the sequence of human SLC-1, in a report by Lakaye et al. (Lakaye, B. et al. (1998) Biochem. Biophys. Acta, vol. 1401, pp. 216-220) in that the presence of the initiating codon ATG on mRNA was indicated at 69 and 64 amino acids further upstream of the deduced sequence. A transformant Escherichia coli DH10B/phSLC1L8 obtained using a plasmid containing DNA encoding this sequence was deposited at IFO and NIBH.

Reference Example 7E Amplification of human SLC-1 cDNA by PCR using cDNA derived from human fetal brain

Using, as a template, the plasmid containing human SLC-1 DNA sequence cloned by the gene-trap method, synthetic DNA primers of SEQ Nos:10 and 11 and synthetic DNA primers of SEQ Nos:12 and 13, amplification was conducted by the PCR method. The amplified DNA of the former was named human SLC-1(S) and the amplified DNA of the latter was named human SLC-1(L). The synthetic DNA primers were constructed such that the gene of the region to be translated into the receptor protein was amplified, during which restriction enzyme recognizing sequences of restriction enzyme Sal I and restriction enzyme Spe I were added to the 5' side and 3' side, respectively, so that a base sequence recognized by the restriction enzyme Sal I would be added to the 5' side of the gene and a base sequence recognized by the restriction enzyme Spe I would be added to the 3' side of the gene. The composition of the reaction mixture for human SLC-1(S) amplification was plasmid template containing human SLC-1 DNA sequence 5 μ l, each synthetic DNA primer 0.4 μ M, 0.2 mM dNTPs, pfuDNA polymerase 0.5 μ l and

buffer annexed to the enzyme, with the total reaction volume of 50 μ l. For amplification cycle, Thermal Cycler (Perkins Elmer) was used. After heating at 94°C for 60 seconds, a cycle of heating at 94°C for 60 seconds, at 57°C for 60 seconds, and at 72°C for 150 seconds was repeated 25 times, and the mixture was finally incubated at 72°C for 10 minutes. The composition of the reaction mixture for human SLC-1(L) amplification was plasmid template containing human SLC-1 DNA sequence 5 μ l, each synthetic DNA primer 0.4 μ M, 0.2 mM dNTPs, pfuDNA polymerase 0.5 μ l and buffer annexed to the enzyme, with the total reaction volume of 50 μ l. For amplification cycle, Thermal Cycler (Perkins Elmer) was used. After heating at 94°C for 60 seconds, a cycle of heating at 94°C for 60 seconds, at 60°C for 60 seconds, and at 72°C for 3 min was repeated 25 times, and the mixture was finally incubated at 72°C for 10 minutes. The amplified product was confirmed by ethidium bromide staining after 0.8% agarose gel electrophoresis.

Reference Example 8E Subcloning of PCR product to plasmid vector and confirmation of amplified cDNA sequence by decoding base sequence of insert cDNA

The reaction product after PCR conducted in Reference Example 7E was separated using 0.8% low melting point agarose gel and the band region was excised with a razor, and subjected to minimization, phenol extraction, phenol-chloroform extraction and ethanol precipitation to recover DNA. According to the direction of PCR-Script™ Amp SK(+) cloning kit (Stratagene), the recovered DNA was subcloned to plasmid vector pCR-Script Amp SK(+). This was introduced into Escherichia coli DH5 α competent cell (TOYOBO) to allow transformation, after which clones containing cDNA insert fragment were selected in an LB agar medium containing ampicillin and X-gal, separated using a sterile toothpick for white clones to give transformant E. coli DH5 α /hSLC-1(S) of human SLC-1(S) and

transformant E. coli DH5 α /hSLC-1(L) of human SLC-1(L). The respective clones were cultured overnight in an LB medium containing ampicillin, and using QIA prep8 mini prep (QIAGEN), the plasmid DNA was prepared. A part of the prepared DNA was
5 cleaved with restriction enzymes Sal I and Spe I to confirm the size of the inserted receptor cDNA fragment. The reaction for determination of the base sequence was carried out using DyeDeoxy Terminator Cycle Sequence Kit (Perkins Elmer), and decoded using a fluorescence automatic DNA sequencer. The
10 sequences of the obtained clones were respectively identical with the DNA sequence (SEQ No:14) to be amplified using synthetic DNA primers of SEQ Nos:10 and 11 and DNA sequence (SEQ No:15) to be amplified using synthetic DNA primers of SEQ Nos: 12 and 13, with human SLC-1 gene as a template.

15 **Reference Example 9E** Preparation of human SLC-1(S) expression
CHO cell and human SLC-1(L) expression CHO cell

From a clone of *E. coli* transformed with a plasmid incorporating human SLC-1(S) and human SLC-1(L), whose sequences were confirmed in Reference Example 8E, a plasmid was prepared using Plasmid Midi Kit (QIAGEN) and cleaved with restriction enzymes Sal I and Spe I to excise an insert. The insert DNA was recovered by cutting out, after electrophoresis, from agarose gel with a razor and applying minimization, phenol extraction, phenol-chloroform extraction and ethanol precipitation. This insert DNA was added to animal cell expression vector plasmid pAKKO-111H (vector plasmid identical with pAKKO1.11H described in Hinuma, S. et al. *Biochim. Biophys. Acta*, Vol. 1219, pp. 251-259 (1994)) cleaved with Sal I and Spe I and ligated using T4 ligase (Takara Shuzo Co.) to respectively construct protein expression plasmids pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L).

E. coli DH5 α (TOYOBO) transformed with pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) was cultured and, using Plasmid Midi Kit

(QIAGEN), plasmid DNAs of pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) were prepared. These were introduced into CHO dhfr⁻ cells using CellPfect Transfection Kit (Amersham Pharmacia Biotech) in accordance with the attached protocol. DNA (10 µg) was
5 prepared into a coprecipitation suspension with calcium phosphate and added into a 10 cm dish inoculated with 5×10^5 or 1×10^6 CHO dhfr⁻ cells 24 hours before. The cells were cultured in an MEM α medium containing 10% fetal bovine serum for one day, passaged and cultured in a nucleic acid-free MEM α
10 medium (selection medium) containing 10% dialyzed fetal bovine serum. 56 clones of transformed cell colonies, which were human SLC-1(S) gene introduced CHO cells, and 61 clones of transformed cell colonies, which were human SLC-1(L) gene introduced CHO cells, grew in the selection medium and were
15 selected.

Reference Example 10E Selection of gene introduced cell line with high expression amount of human SLC-1(S) mRNA and human SLC-1(L) mRNA

The expression amount of mRNA of 56 clone of CHO/hSLC-
20 1(S) cell line and 61 clones of CHO/hSLC-1(L) cell line established in Reference Example 9E was measured as in the following using Cytostar T Plate (Amersham Pharmacia Biotech) and in accordance with the attached protocol.

Each clone of the CHO/hSLC-1(S) cell line and CHO/hSLC-
25 1(L) cell line was inoculated to each well of Cytostar T Plate at 2.5×10^4 , cultured for 24 hours and fixed with 10% formalin. After adding 0.25% Triton X-100 to each well to enhance permeability of the cells, riboprobe of ³⁵S labeled SEQ No:16 was added for hybridization. RNase A (20 mg/ml) was added to
30 each well to digest free riboprobe. The plate was washed thoroughly and the radioactivity of the hybridized riboprobe was measured on Topcounter. The cell strain having high radioactivity showed higher expression amount of mRNA. Of the

Experimental Example 1 Determination of antagonistic activity using GTP γ S binding assay of test compound

5 Using human SLC-1 expression CHO cell clone 57 obtained
in Reference Example 10E and rat SLC-1 expression CHO cell
clone 44 obtained in Reference Example 4E, membrane fractions
were prepared by the following method. In phosphate buffered
10 saline (pH 7.4) supplemented with 5 mM EDTA (ethylenediamine
teraacetic acid) were suspended human and rat SLC-1 expression
CHO cells (1×10^8) and centrifuged. Homogenate buffer (10 ml,
10 mM NaHCO_3 , 5 mM EDTA, pH 7.5) was added to the pellets of
the cells and, using Polytron Homogeniser, the mixture was
homogenated. The supernatant obtained after centrifugation at
15 $400 \times g$ for 15 min was further centrifuged at $100,000 \times g$ for 1 hr
to give precipitate of the membrane fraction. The precipitate
was suspended in 2 ml of an assay buffer [50 mM Tris-HCl (pH
7.5), 1 mM EDTA, 0.1% BSA (bovine serum albumin), 10 mM MgCl_2 ,
100 mM NaCl, 1 mM GDP (guanosine 5'-diphosphate), 0.25 mM PMSF
20 (phenylmethylsulfonylfluoride), 1 mg/ml pepstatin, 20 mg/ml
leupeptin, 10 mg/ml phosphoramidon] and centrifuged at
 $100,000 \times g$ for 1 hr. The membrane fraction recovered as
precipitate was suspended again in 20 ml of an assay buffer,
and after dispensing, preserved at -80°C and used upon thawing
25 each time when in use.

The antagonistic activity of the test compound was determined as follows. The SLC-1 expression CHO cell membrane fraction (171 μ l) diluted with an assay buffer was dispensed to a polypropylene 96 well plate and 3×10^{-10} M MCH (2 ml) diluted with DMSO solution, test compound solution (2 ml) diluted to various concentrations and [35 S]-Guanosine-5'-(γ -thio) triphosphate (25 ml, Daiichi Pure Chemicals Co., Ltd.) were respectively added (cell membrane final concentration: 20 mg/ml,

[³⁵S]-Guanosine 5'-(γ -thio)triphosphate final concentration: 0.33 nM). The reaction mixture was reacted at 25°C for 1 hr with stirring, suction filtered with a glass filter (GF-C) and washed 3 times with a wash solution (300 ml, 50 mM Tris-HCl buffer, pH 7.5). Liquid Scintillator (50 ml) was added to the glass filter and the residual radioactivity was determined by a liquid scintillation counter.

Binding inhibition (%) = (radioactivity upon addition of compound and MCH - radioactivity upon addition of DMSO solution) / (radioactivity upon addition of MCH - radioactivity upon addition of DMSO solution) x 100

From the binding inhibition (%), IC₅₀ of the compound was calculated. The results are shown in Table 1.

Table 1


Compound	Inhibitory activity (IC ₅₀ :nM)
Reference Example IIA-5	100
Example 1	5

Industrial Applicability

The compound (I) and a salt thereof or a prodrug thereof of the present invention have a superior MCH antagonistic action and are useful as an agent for the prophylaxis and treatment of the diseases (e.g., obesity and the like) caused by melanin-concentrating hormone.

WHAT IS CLAIMED IS

1. A melanin-concentrating hormone antagonist containing a compound of the formula



5 wherein

Ar¹ and Ar² are each an aromatic group optionally having substituents,

P and Q are each a divalent aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which optionally

10

R¹ and R³ are each (i) a hydrogen atom, (ii) an acyl group or (iii) a hydrocarbon group optionally having substituents,

15 R² and R⁴ are each (i) a hydrogen atom, (ii) an alkyl group optionally having substituents or (iii) an alkylcarbonyl group optionally having substituents,

R¹ and R² or R³ and R⁴

20 optionally form, together with the adjacent nitrogen atom, a monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents, and

j is 0 or 1,

25 or a salt thereof or a prodrug thereof.

2. The antagonist of claim 1, wherein Ar¹ and Ar² are each (i) a C₆₋₁₄ aryl group or (ii) a 5 to 14-membered monocyclic or

(a) a halogen atom,
(b) a C₁₋₃ alkylenedioxy group,
(c) a nitro group,
(d) a cyano group,

(i) a hydroxy group,

(k) a mono-C₁₋₆ alkylamino group,

(l) a di-C₁₋₆ alkylamino group,

halogen atom or C₁₋₆ alkoxy-carbonyl group,

(g) a carboxyl group,

²⁵ (s) a carbamoyl group,

C₁₋₆ alkoxy-carbonyl group,

C₁₋₆ alkoxy-carbonyl group,

(w) a C₁₋₆ alkylsulfonyl group,

(y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s)

selected from above-mentioned (a) to (x),

(z) a C₆₋₁₀ aryloxy group optionally having 1 to 4

substituent(s) selected from the above-mentioned (a) to (x),

(aa) an optionally halogenated C₆₋₁₀ aryl-carbonyl group,

⁵ (ab) an optionally halogenated 5 or 6-membered heterocyclic ring-carbonyl group,

(ac) a C_{1-6} alkoxy-carbonylamino group,

(ad) a C₆₋₁₀ aryl-carbonylamino group and

(ae) a C₇₋₁₆ aralkyloxy-carbonyl group,

¹⁰ P and Q are each a divalent C₁₋₆ aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which is optionally substituted by oxo group or thioxo group;

R¹ and R³ are each (i) hydrogen atom, (ii) acyl group

¹⁵ represented by $-\text{CO}-\text{R}^a$, $-\text{CONR}^a\text{R}^b$, $-\text{SO}-\text{R}^a$, $-\text{SO}_2-\text{R}^a$, $-\text{CONR}^a\text{R}^b$, $-\text{COO}-\text{R}^a$, $-(\text{C}=\text{S})\text{O}-\text{R}^a$, $-(\text{C}=\text{S})\text{NR}^a\text{R}^b$, $-\text{SONR}^a\text{R}^b$, $-\text{SO}_2\text{NR}^a\text{R}^b$, $-\text{SO}-\text{O}-\text{R}^a$ or $-\text{SO}_2-\text{O}-\text{R}^a$, wherein R^a is (A) hydrogen atom; (B) carboxyl group;

(C) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋

20 16 aralkyl group, which optionally has 1 to 5 substituent(s)
selected from the group (group Ba) consisting of

(a) a halogen atom,

(b) a C₁₋₃ alkylenedioxy group,

(c) a nitro group,

²⁵ (d) a cyano group,

(e) an optionally halogenated C₁₋₆ alkyl group,

(f) an optionally halogenated C₃₋₆ cycloalkyl group,

(g) an optionally halogenated C₁₋₆ alkoxy group,

(h) an optionally halogenated C₁₋₆ alkylthio group,

³⁰ (i) a hydroxy group,

(j) an amino group,


(k) a mono-C₁₋₆ alkylamino group,

(1) a di-C₁₋₆ alkylamino group,

- (m) a C₁₋₆ alkyl-carbonylamino group,
- (n) a formyl group,
- (o) a C₁₋₆ alkyl-carbonyl group,
- (p) a C₁₋₆ alkyl-carbonyloxy group,
- 5 (q) a carboxyl group,
- (r) a C₁₋₆ alkoxy-carbonyl group,
- (s) a carbamoyl group,
- (t) a mono-C₁₋₆ alkyl-carbamoyl group,
- (u) a di-C₁₋₆ alkyl-carbamoyl group,
- 10 (v) a sulfo group,
- (w) a C₁₋₆ alkylsulfonyl group,
- (x) a C₁₋₆ alkylsulfinyl group,
- (y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s) selected from the aforementioned (a) to (x),
- 15 (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4 substituent(s) selected from the aforementioned (a) to (x),
- (zz) a 5 to 7-membered heterocyclic group optionally having 1 to 4 substituent(s) selected from the aforementioned (a) to (x),
- (aa) a di-C₁₋₆ alkyl-carbonylamino group,
- 20 (ab) a sulfamoyl group,
- (ac) a C₁₋₆ alkoxy-carbonylamino group,
- (ad) a C₇₋₁₆ aralkyloxy-carbonylamino group,
- (ae) a C₇₋₁₆ aralkyloxy group,
- (af) a C₆₋₁₀ aryl-carbonyl group,
- 25 (ag) a C₁₋₆ alkyl-carbonyloxy group,
- (ah) a C₆₋₁₀ aryl-carbonylamino group,
- (ai) a C₆₋₁₀ aryl-carbamoyl group,
- (aj) a C₇₋₁₆ aralkylaminocarbonyl group,
- (ak) a C₇₋₁₆ aralkylcarbonylamino group and
- 30 (al) a C₇₋₁₆ aralkyloxy-carbonyloxy group;
- (D) a 5 to 10-membered heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, which


substituent(s) selected from group Aa, W^a is hydrogen atom, hydroxy group or C_{1-6} alkyl group),

(ii) the formula



5 wherein ring B is monocyclic or bicyclic 4 to 12-membered ring optionally substituted by 1 or 2 oxo group(s) or 1 to 5 C_{1-6} alkyl group(s), ring D is a 4 to 12-membered aromatic ring optionally having 1 to 5 substituent(s) selected from group Aa or

10 (iii) the formula



wherein ring E is a 4 to 12-membered aromatic ring optionally having 1 to 5 substituent(s) selected from group Aa;

X is $-CH_2-$, $-CO-$ or $-CH(OH)-$;

15 Y is $-CH_2-$, $-O-$ or $-NW^b-$ (W^b is (a) hydrogen atom or (b) C_{1-6} alkyl group optionally having substituents selected from group Ba);

k and m are each an integer of 0 to 4, and $k+m$ is an integer of 1 to 4;

20 n is an integer of 1 to 3.

3. The antagonist of claim 1, wherein Ar^1 and Ar^2 are each (i) a C_{6-14} aryl group or (ii) a 5 to 14-membered monocyclic or fused aromatic heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from the group consisting of nitrogen atom, sulfur atom and oxygen atom, which optionally has 1 to 5 substituent(s) selected from the group (group A)

selected from the aforementioned (a) to (x),
 (z) a C₆₋₁₀ aryloxy group optionally having 1 to 4
 substituent(s) selected from the aforementioned (a) to (x) and
 (zz) a 5 to 7-membered heterocyclic group optionally having 1
 5 to 4 substituent(s) selected from the aforementioned (a) to (x),
 or
 (d) a 5 to 10-membered heterocyclic group containing, besides
 carbon atom, 1 to 4 heteroatom(s) selected from the group
 consisting of nitrogen atom, oxygen atom and sulfur atom, which
 10 optionally has 1 to 5 substituent(s) selected from the group
 (group C) consisting of
 (a) a halogen atom,
 (b) a C₁₋₃ alkylendioxy group,
 (c) a nitro group,
 15 (d) a cyano group,
 (e) a C₁₋₆ alkyl group optionally having substituents selected
 from the group consisting of (aa) a halogen atom, (bb) C₁₋₃
 alkylendioxy group, (cc) nitro group, (dd) cyano group, (ee)
 an optionally halogenated C₁₋₆ alkyl group, (ff) an optionally
 20 halogenated C₃₋₆ cycloalkyl group, (gg) an optionally
 halogenated C₁₋₆ alkoxy group, (hh) an optionally halogenated
 C₁₋₆ alkylthio group, (ii) a hydroxy group, (jj) amino group,
 (kk) a mono-C₁₋₆ alkylamino group, (ll) a di-C₁₋₆ alkylamino
 group, (mm) C₁₋₆ alkyl-carbonylamino group, (nn) a formyl group,
 25 (oo) C₁₋₆ alkyl-carbonyl group, (pp) C₁₋₆ alkyl-carboxyloxy group,
 (qq) carboxyl group, (rr) C₁₋₆ alkoxy-carbonyl group, (ss)
 carbamoyl group, (tt) a mono-C₁₋₆ alkylcarbamoyl group, (uu) a
 di-C₁₋₆ alkylcarbamoyl group, (vv) a sulfo group, (ww) C₁₋₆
 alkylsulfonyl group, (xx) C₁₋₆ alkylsulfinyl group, (yy) C₆₋₁₀
 30 aryl group optionally having 1 to 4 substituent(s) selected
 from the aforementioned (aa) to (xx), (zz) C₆₋₁₀ aryloxy group
 optionally having 1 to 4 substituent(s) selected from the
 aforementioned (aa) to (xx) and (zzz) 5 to 7-membered

heterocyclic group optionally having 1 to 4 substituent(s)
selected from the aforementioned (aa) to (xx),

(f) an optionally halogenated C₃₋₆ cycloalkyl group,

(g) an optionally halogenated C₁₋₆ alkoxy group,

⁵ (h) an optionally halogenated C₁₋₆ alkylthio group,

(i) a hydroxy group,

(j) an amino group,

(k) a mono-C₁₋₆ alkylamino group,

(1) a di-C₁₋₆ alkylamino group,

¹⁰ (m) an optionally halogenated C₁₋₆ alkyl-carbonylamino group,

(n) a formyl group,

(o) a C₁₋₆ alkyl-carbonyl group,

(p) a C₁₋₆ alkyl-carbonyloxy group,

(g) a carboxyl group,

¹⁵ (r) a C₁₋₆ alkoxy-carbonyl group,

(s) a carbamoyl group,

(t) a mono-C₁₋₆ alkylcarbamoyl group,

(u) a di-C₁₋₆ alkylcarbamoyl group,

(v) a sulfo group,

²⁰ (w) a C₁₋₆ alkylsulfonyl group,

(x) a C₁₋₆ alkylsulfinyl group,

(y) a C₆₋₁₀ aryl group optionally having 1 to 4 substituent(s)

selected from the aforementioned (a) to (x) and

(z) a C₆₋₁₀ aryloxy group optionally having 1 to 4
 25 substituent(s) selected from the aforementioned (a) to (x), and
 R^b is a hydrogen atom or a C₁₋₆ alkyl group) or


(iii) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋₁₆ aralkyl group, which optionally has 1 to 5

³⁰ substituent(s) selected from group B.


R² and R⁴ are each (i) hydrogen atom, (ii) C₁₋₆ alkyl optionally having substituents selected from group B or (iii) C₁₋₆ alkyl-carbonyl group optionally having substituents selected from

group B,


R¹ and R² or R³ and R⁴ form, together with the adjacent nitrogen atom, a nitrogen-containing heterocyclic group represented by (i) the formula



wherein ring A is a 4 to 8-membered ring optionally substituted by hydroxy or oxo, V is a group represented by the formula >O, >C=O, >C-(W)W^a or >N-W (W is (a) hydrogen atom, (b) (1) C₁₋₆ alkyl group, (2) C₂₋₆ alkenyl group, (3) C₂₋₆ alkynyl group, (4) C₃₋₆ cycloalkyl group, (5) C₆₋₁₄ aryl group or (6) C₇₋₁₆ aralkyl group, which optionally has 1 to 5 substituent(s) selected from group B, or (c) 5 to 10-membered heterocyclic group containing, besides carbon atom, 1 to 4 heteroatom(s) selected from nitrogen, oxygen and sulfur, which optionally has 1 to 5 substituent(s) selected from group A, W^a is hydrogen atom or hydroxy group),
(ii) the formula



wherein ring B is monocyclic or bicyclic 4 to 12-membered ring optionally substituted by oxo group or 1 to 5 C₁₋₆ alkyl group(s), ring D is a 4 to 12-membered aromatic ring optionally having 1 to 5 substituent(s) selected from group A or
(iii) the formula




wherein ring E is a 5 to 10-membered aromatic ring optionally


wherein ring A¹ is a 4 to 8-membered ring optionally substituted by hydroxy or oxo, V¹ is a group represented by the formula >O, >C(W¹)-W^{a1} or >N-W¹ (W¹ is (a) hydrogen atom, (b) C₆₋₁₄ aryl group optionally having 1 or 2 substituent(s)

5 selected from the group consisting of a halogen atom, optionally halogenated C₁₋₆ alkyl group and optionally halogenated C₁₋₆ alkoxy group, (c) C₁₋₆ alkyl group optionally having 1 or 2 C₆₋₁₀ aryl group(s) or (d) pyridyl group, W^{a1} is hydrogen atom, hydroxy group or C₁₋₆ alkyl group),

¹⁰ (ii) the formula



wherein ring B¹ is a monocyclic or bicyclic 5 to 10-membered ring optionally substituted by oxo group or 1 or 2 C₁₋₆ alkyl group(s), ring D¹ is a benzene ring optionally having 1 or 2 substituent(s) selected from the group consisting of C₁₋₆ alkyl group, C₁₋₆ alkoxy group and C₁₋₆ alkyl-carbonyl group or (iii) the formula




wherein ring E¹ is a benzene ring optionally having 1 to 3
substituent(s) selected from the group consisting of C₁₋₃
alkylenedioxy group, nitro group, C₁₋₆ alkoxy group, amino group,
C₁₋₆ alkyl-carbonylamino group and C₁₋₆ alkoxy-carbonyl group,
X¹ is -CH₂- or -CO-, and
Y¹ is -CH₂- or -O-,
R³ is (i) hydrogen atom,
(ii) a group represented by the formula -CO-R⁵ (R⁵ is (a)
hydrogen atom, (b) carboxyl group, (c) C₁₋₆ alkyl group, (d) C₅₋₆
cycloalkyl group optionally having C₁₋₆ alkoxy, and which is

alkoxy-carbonyl, morpholino and mono- or di-C₁₋₆ alkylamino, (c) C₁₋₆ alkoxy-carbonyl group, (d) a group represented by the formula -CO-R^d (R^d is C₆₋₁₀ aryl group optionally having halogen atom or 5 or 6-membered heterocyclic group containing, besides
 5 carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom), (e) a group represented by the formula -CO-(CH₂)^{r¹}-R^e (r¹ is an integer of 1 to 3, R^e is C₁₋₆ alkoxy-carbonyl group or 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected
 10 from nitrogen atom, oxygen atom and sulfur atom) or (f) a group represented by -CONH-R^f (R^f is C₁₋₆ alkyl group or C₆₋₁₄ aryl group),

(vi) a group represented by the formula

-COOR⁹ (R⁹ is optionally halogenated C₁₋₆ alkyl group),

15 (vii) a group represented by the formula




wherein R¹⁰ is hydrogen atom, C₁₋₆ alkoxy-carbonyl group, mono- or di-C₁₋₆ alkyl-carbamoyl group, optionally halogenated nicotinoyl group or optionally halogenated isonicotinoyl group,

20 (viii) a group represented by the formula

-CONR¹¹-R¹² (R¹¹ is hydrogen atom or C₁₋₆ alkyl group, R¹² is C₁₋₆ alkyl group optionally having substituents selected from the group consisting of (a) hydroxy, (b) amino, (c) a mono- or di-C₁₋₆ alkyl-amino, (d) C₁₋₆ alkyl-carbonyl, (e) C₁₋₆ alkoxy-

25 carbonyl, (f) C₁₋₆ alkyl-carbonyloxy, (g) sulfamoyl and (h) 5 to 7-membered heterocyclic group optionally substituted by oxo, and (i) C₆₋₁₄ aryl group),


(ix) a group represented by the formula



30 wherein R¹³ is (a) hydrogen atom, (b) C₁₋₆ alkyl group optionally having substituents selected from the group


(xv) C₁₋₆ alkyl group substituted by 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom,

5 (xvi) a group represented by the formula




wherein R²⁴ is hydrogen atom or C₇₋₁₆ aralkyloxy-carbonyl group;

(xvii) a group represented by the formula




10 wherein R²⁵ is hydrogen atom, C₆₋₁₀ aryl group, C₇₋₁₆ aralkyloxy group, C₆₋₁₀ aryloxy group, halogen atom, C₆₋₁₀ aryl-carbonylamino group or C₆₋₁₀ aryl-carbamoyl group;

(xviii) a group represented by the formula



15 [Alk is C₁₋₆ alkylene group optionally having substituents; R²⁷ is hydrogen atom or C₁₋₆ alkyl group; Alk₂ and Alk₃ are the same or different and each is a bond or C₁₋₆ alkylene group optionally having substituents; R²⁸ is C₆₋₁₀ aryl group optionally having substituents or hydrogen atom];


20 (xix) a group represented by the formula



[Alk₂, Alk₃ and R²⁷ are as defined above; R²⁹ is (1) C₆₋₁₀ aryl group optionally having substituent or (2) 5 to 10-membered aromatic heterocyclic group optionally having substituent,

25 which contains, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom];


(xx) a group represented by the formula



[Alk, R^{27} , Alk_2 , Alk_3 are as defined above; R^{30} is hydrogen atom, C_{1-6} alkyl group or optionally halogenated C_{1-6} alkyl-carbonyl group; and R^{31} is C_{6-10} aryl group optionally having


5 substituents];

(xxi) a group represented by the formula




[Alk, R^{27} , Alk_2 , Alk_3 and R^{31} are as defined above; and R^{32} is the same as the aforementioned R^{27}];

10 (xxii) a group represented by the formula



[Alk, R^{27} , Alk_2 and R^{31} are as defined above]; or


(xxiii) a group represented by the formula



15 [Alk, Alk_2 and R^{31} are as defined above];

R^4 is hydrogen atom or C_{1-6} alkyl group;

or R^3 and R^4 may form, together with the adjacent nitrogen atom, a group represented by the formula



20 wherein R^{18} is halogen atom, oxo group, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group.

8. The antagonist of claim 1, wherein R^1 is (i) C_{1-6} alkyl group optionally having 5 or 6-membered nitrogen-containing

25 heterocyclic group, (ii) C_{7-16} aralkyl group optionally having nitro, amino or C_{1-6} alkoxy-carbonyl or (iii) cyclohexyl group fused with benzene ring optionally having C_{1-6} alkoxy,

R^2 is (i) hydrogen atom, (ii) C_{1-6} alkyl group or (iii) C_{7-16} aralkyl group, or R^1 and R^2 may form, together with the


30 adjacent nitrogen atom, a nitrogen-containing heterocyclic group represented by

(i) the formula




wherein ring A¹ is a 4 to 8-membered ring optionally substituted by hydroxy or oxo, V¹ is a group represented by the
 5 formula >O, >C-(W¹)W^{a1} or >N-W¹ (W¹ is (a) hydrogen atom, (b) C₆₋₁₄ aryl group optionally having 1 or 2 substituent(s) selected from the group consisting of halogen atom, optionally halogenated C₁₋₆ alkyl group and C₁₋₆ alkoxy group or (c) C₁₋₆ alkyl group optionally having 1 or 2 C₆₋₁₀ aryl group(s), W^{a1} is
 10 hydrogen atom or hydroxy group),

(ii) the formula




wherein ring B¹ is a monocyclic or bicyclic 5 to 10-membered ring optionally substituted by oxo group or 1 or 2 C₁₋₆ alkyl
 15 group(s), ring D¹ is a benzene ring optionally having 1 or 2 substituent(s) selected from the group consisting of C₁₋₆ alkyl group, C₁₋₆ alkoxy group and C₁₋₆ alkyl-carbonyl group or


(iii) the formula



20 wherein ring E¹ is a benzene ring optionally having 1 or 2 substituent(s) selected from the group consisting of C₁₋₃ alkylenedioxy group, nitro group, C₁₋₆ alkoxy group, amino group, C₁₋₆ alkyl-carbonylamino group and C₁₋₆ alkoxy-carbonyl group
 X¹ is -CH₂- or -CO-,
 25 Y¹ is -CH₂- or -O-,
 R³ is (i) hydrogen atom, (ii) a group represented by the

20 ---CO---Q---  N---R^7

(v) a group represented by the formula



258


5 or 6-membered heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom),

(e) a group represented by the formula $-\text{CO}-(\text{CH}_2)_{r^1}-\text{R}^e$ (r^1 is an integer of 1 to 3, R^e is C_{1-6} alkoxy-carbonyl group or 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 or 2 heteroatom(s) selected from nitrogen atom, oxygen atom and sulfur atom) or (f) a group represented by $-\text{CONH}-\text{R}^f$ (R^f is C_{1-6} alkyl group or C_{6-14} aryl group)),

(vi) a group represented by the formula

$-\text{COOR}^9$ (R^9 is optionally halogenated C_{1-6} alkyl group),


(vii) a group represented by the formula



wherein R^{10} is hydrogen atom, C_{1-6} alkoxy-carbonyl group, mono or di- C_{1-6} alkyl-carbamoyl group, optionally halogenated nicotinoyl group or optionally halogenated isonicotinoyl group,

(viii) a group represented by the formula $-\text{CONR}^{11}-\text{R}^{12}$ (R^{11} is hydrogen atom or C_{1-6} alkyl group, R^{12} is C_{1-6} alkyl optionally having substituents selected from the group consisting of (a) hydroxy, (b) amino, (c) a mono- or di- C_{1-6} alkyl-amino, (d) C_{1-6} alkyl-carbonyl, (e) C_{1-6} alkoxy-carbonyl, (f) C_{1-6} alkyl-carbonyloxy, (g) sulfamoyl and (f) 5 or 6-membered cyclic amine optionally substituted by oxo),


(ix) a group represented by the formula



wherein R^{13} is (a) a hydrogen atom, (b) C_{1-6} alkyl group optionally having substituents selected from the group consisting of a hydroxy and C_{1-6} alkoxy-carbonyl, (c) C_{7-16} aralkyl group, (d) C_{1-6} alkyl-carbonyl group optionally having substituents selected from the group consisting of a halogen and C_{1-6} alkoxy-carbonyl or (e) C_{1-6} alkyl-carbamoyl group


optionally having C₁₋₆ alkoxy-carbonyl,

(x) a group represented by the formula



wherein R¹⁴ is C₁₋₆ alkyl group or C₇₋₁₆ aralkyl group,


5 (xi) a group represented by the formula



wherein ring F is 5 to 7-membered cyclic amino group optionally fused with benzene ring, R¹⁵ is hydrogen atom, C₁₋₆ alkoxy-carbonylamino group or optionally halogenated C₁₋₆ alkyl-

10 carbonylamino group,

(xii) a group represented by the formula



wherein R¹⁶ is (a) C₁₋₆ alkyl group optionally having substituents selected from the group consisting of a hydroxy and C₁₋₆ alkoxy-carbonyl, (b) a formyl group, (c) C₁₋₆ alkoxy-carbonyl group or (d) a 5 or 6-membered heterocyclic ring-carbonyl group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom,

20 (xiii) a group represented by the formula


---SO₂---R¹⁷ (R¹⁷ is (i) C₁₋₆ alkyl group optionally having 5 or 6-membered nitrogen-containing ring group, (ii) C₂₋₆ alkenyl group or (iii) C₆₋₁₄ aryl group optionally having C₁₋₆ alkyl),

(xiv) C₇₋₁₆ aralkyl group optionally having 1 to 3 halogen

25 atom(s), or


(xv) C₁₋₆ alkyl group substituted by 5 or 6-membered heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom,

R^4 is hydrogen atom or C_{1-6} alkyl group,
or R^3 and R^4 may form, together with the adjacent nitrogen atom,
a group of the formula



5 wherein R^{18} is halogen atom, oxo group, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group.

9. The antagonist of claim 1, wherein R^1 and R^2 form, together with the adjacent nitrogen atom, a nitrogen-containing
10 heterocyclic group represented by
(i) the formula




wherein q^1 is hydrogen atom or halogen atom, q^2 is hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or
15 C_{1-6} alkoxy group, q^3 is hydrogen atom or halogen atom, q^4 is hydrogen atom, halogen atom or C_{1-6} alkoxy group, q^5 is hydrogen atom or C_{1-6} alkyl group optionally having 1 or 2 C_{6-10} aryl group(s),


(ii) the formula



wherein ring B² is represented by the formula




wherein ring D¹ is benzene ring optionally having 1 or 2
substituent(s) selected from the group consisting of C₁₋₆ alkyl
5 group, C₁₋₆ alkoxy group and C₁₋₆ alkyl-carbonyl group or
(iii) the formula



wherein ring E¹ is benzene ring optionally having 1 or 2
substituent(s) selected from the group consisting of C₁₋₃
10 alkylenedioxy group, nitro group, C₁₋₆ alkoxy group, amino group,
C₁₋₆ alkyl-carbonylamino group and C₁₋₆ alkoxy-carbonyl group, X¹
is -CH₂- or -CO-, and Y¹ is -CH₂- or -O-.


10. The antagonist of claim 1, wherein the compound is
15 represented by the formula



wherein R¹⁹ is (i) hydrogen atom, (ii) carboxyl, (iii) C₁₋₆
alkoxy-carbonyl group, (iv) C₁₋₆ alkyl group optionally having
substituents selected from the group consisting of carboxyl,
20 C₁₋₆ alkyl-carbonyl, C₁₋₆ alkoxy-carbonyl, C₁₋₆ alkoxy-
carbonylamino and C₇₋₁₆ aralkyloxy-carbonylamino, (v) a mono- or
di-C₁₋₆ alkylamino group or (iv) C₆₋₁₄ aryloxy group; P¹ is C₁₋₃
alkylene group; Q¹ is C₁₋₃ alkylene group; X² is CH, C-OH or N;


Y^2 is hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or C_{1-6} alkoxy group; and Z is CO, SO or SO_2 .

11. The antagonist of claim 1, wherein the compound is
5 represented by the formula




wherein R^{20} is (i) hydrogen atom or (ii) C_{1-6} alkyl group optionally having substituents selected from the group consisting of C_{1-6} alkoxy-carbonylamino and C_{7-16} aralkyloxy-
10 carbonylamino; P^2 is C_{1-3} alkylene group; X^3 is CH, C-OH or N; Y^3 is hydrogen atom, halogen atom or C_{1-6} alkoxy group.


12. The antagonist of claim 1, wherein the compound is represented by the formula



wherein R^{21} is a nitrogen-containing heterocyclic group represented by (i) the formula



wherein X^4 is CH or N, Y^4 is hydrogen atom, halogen atom or C_{1-6}
20 alkoxy group or (ii) the formula



wherein ring E² is benzene ring optionally having 1 to 3 C₁₋₆ alkoxy,

R²² is (i) hydrogen atom, (ii) C₇₋₁₆ aralkyl group, (iii) formyl group, (iv) C₁₋₆ alkyl-carbonyl group, (v) C₆₋₁₄ aryl-carbonyl group optionally having C₁₋₆ alkyl or (vi) C₆₋₁₄ aryl-sulfonyl group optionally having 1 to 4 C₁₋₆ alkyl; P³ is C₁₋₃ alkylene group; and Q³ is C₁₋₃ alkylene group.

13. The antagonist of claim 1, wherein the compound is
- 10 1-(5-amino-4,4-diphenylpentyl)-4-phenylpiperidine or a salt thereof,
- 3,4-dihydro-6-methoxy-1'-(5-amino-4,4-diphenylpentyl)-spiro[naphthalene-2(1H),2'-piperidine] or a salt thereof,
- 1-[5-amino-4-(4-methoxyphenyl)-4-phenylpentyl]-4-phenylpiperidine or a salt thereof,
- 15 1-[5-amino-4,4-bis(4-chlorophenyl)pentyl]-4-(4-fluorophenyl)-piperazine or a salt thereof,
- 3,4-dihydro-6-methoxy-1'-(6-amino-4,4-diphenylhexyl)-spiro[naphthalene-2(1H),2'-piperidine] or a salt thereof,
- 20 3,4-dihydro-6,7-dimethoxy-1'-(7-amino-4,4-diphenylheptyl)-spiro[naphthalene-2(1H),2'-piperidine] or a salt thereof,
- 4,4-diphenyl-5-formylamino-1-(4-phenylpiperidino)pentane or a salt thereof,
- 1-[4-(4-fluorophenyl)piperazin-1-yl]-5-formylamino-4,4-diphenylpentane or a salt thereof,
- 25 4,4-diphenyl-1-(4-phenylpiperazin-1-yl)-5-(tosylamino)pentane or a salt thereof,
- 4,4-diphenyl-1-[4-(2-methoxyphenyl)piperazin-1-yl]-5-(tosylamino)pentane or a salt thereof,
- 30 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-phenylpiperidino)pentane or a salt thereof,
- 4-(4-chlorophenyl)-5-formylamino-4-phenyl-1-(4-phenylpiperazin-1-yl)pentane or a salt thereof,

4-(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-formylamino-4-phenylpentane or a salt thereof,

4-(4-chlorophenyl)-1-[4-(diphenylmethyl)piperazin-1-yl]-5-formylamino-4-phenylpentane or a salt thereof,

5 5-formylamino-4-(4-methoxyphenyl)-4-phenyl-1-(4-phenylpiperidino)pentane or a salt thereof,

4,4-bis(4-chlorophenyl)-1-[4-(4-fluorophenyl)piperazin-1-yl]-5-(formylamino)pentane or a salt thereof,

1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-5,5-
10 diphenylhexane or a salt thereof,

1-[4-(4-fluorophenyl)piperazin-1-yl]-6-formylamino-4,4-diphenylhexane or a salt thereof,

4,4-diphenyl-1-(4-phenylpiperidino]-6-(tosylamino)hexane or a salt thereof,

15 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof,

5-[4-(4-fluorophenyl)piperazin-1-yl]-1-formylamino-2,2-diphenylpentane or a salt thereof,

1-formylamino-5-(4-hydroxy-4-phenylpiperidino)-2,2-
20 diphenylpentane or a salt thereof,

5-[4-(4-trifluoromethylphenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof,

5-[4-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof,

25 5-[4-(3,5-dichlorophenyl)-4-hydroxypiperidino]-1-formylamino-
2,2-diphenylpentane or a salt thereof,

5-[4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridin-1-yl]-1-formylamino-2,2-diphenylpentane or a salt thereof,

1-formylamino-2,2-diphenyl-5-(4-phenylpiperidino)pentane or a
³⁰ salt thereof,

5-[4-(4-chlorophenyl)piperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof,

7-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylamino-4,4-

- diphenylheptane or a salt thereof,
 5-[4-(4-fluorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane or a salt thereof,
 1-formylamino-5-[4-hydroxy-4-(4-methoxyphenyl)piperidino]-2,2-diphenylpentane or a salt thereof,
 1-formylamino-5-[4-hydroxy-4-(2-pyridyl)piperidino]-2,2-diphenylpentane or a salt thereof,
 1-acetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane or a salt thereof,
 1-acetoacetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane or a salt thereof,
 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamate or a salt thereof,
 N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid or a salt thereof,
 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-ethylurea or a salt thereof,
 N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]methanesulfonamide or a salt thereof,
 phenyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]carbamate or a salt thereof,
 1-acetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentane or a salt thereof,
 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]oxamate or a salt thereof,
 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]malonamate or a salt thereof,
 ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glutaramate or a salt thereof,
 benzyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate or a salt thereof,
 tert-butyl 2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethylcarbamate or a salt thereof,

4,4-diphenyl-7-(4-phenylpiperidino)heptylamine or a salt thereof,

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-4-methylbenzenesulfonamide or a salt thereof,

5 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)acetamide or a salt thereof,

N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)amine or a salt thereof,

10 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(3-methoxybenzyl)amine or a salt thereof,

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-methoxybenzyl)amine or a salt thereof,

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-fluorobenzyl)amine or a salt thereof,

15 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-thiophenecarboxamide or a salt thereof,

N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-2-phenylacetamide or a salt thereof,

20 N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-(2-thienylmethyl)amine or a salt thereof, or

N-benzyl-N-(4,4-diphenyl-7-(4-phenylpiperidino)heptyl)-N-methylamine or a salt thereof.

14. The antagonist of claim 1, which is an agent for the
25 prophylaxis or therapy of a disease caused by melanin-concentrating hormone.


15. The antagonist of claim 1, which is an agent for the
prophylaxis or therapy of obesity.

30

16. The antagonist of claim 1, which is an agent for suppressing food intake.

phenylpiperidino)pentyl) amino)-2-oxoethylcarbamate or a salt thereof.


20. A production method of a compound of claim 17, which
5 comprises reacting a compound represented by the formula



wherein each symbol is as defined in claim 17 or a salt thereof with a reactive derivative of an organic acid of the formula
 $R^{23}-COOH$

- 10 wherein R^{23} is as defined in claim 17.

21. A production method of a compound of claim 17, which
comprises reacting a compound represented by the formula



- 15 wherein each symbol is as defined in claim 17, or a salt thereof with a reactive derivative of the formula




wherein R^{32} is C_{7-16} aralkyloxy-carbonyl group, and X is a leaving group.

20

22. A pharmaceutical composition containing a compound of claim 17.

23. A compound represented by the formula



wherein R^{26} and R^{27} are the same or different and each is hydrogen atom or C_{1-6} alkyl group; Alk_2 and Alk_3 are the same or different and each is a bond or C_{1-6} alkylene group optionally having substituents; R^{29} is (1) C_{6-10} aryl group optionally having substituents or (2) 5 to 10-membered aromatic heterocyclic group optionally having substituents, which contains, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom; X^2 is CH, C-OH or N; P^5 and Q^5 are the same or different and each is C_{1-6} alkylene group; Y^6 , Y^7 and Y^8 are the same or different and each is hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group, or a salt thereof or a prodrug thereof.

24. The compound of claim 23, wherein Alk_2 and Alk_3 are the same or different and each is a bond, or C_{1-6} alkylene group optionally having substituents selected from the group consisting of halogen atom, hydroxy, amino and C_{6-10} aryl; R^{29} is (1) C_{6-10} aryl group or (2) 5 to 10-membered aromatic heterocyclic group containing, besides carbon atom, 1 to 3 heteroatom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, which optionally has substituents selected from the group consisting of nitro, halogen atom, C_{1-6} alkyl, hydroxy, C_{1-6} alkoxy and C_{6-10} aryl.

25. The compound of claim 23 or 24, wherein R^{29} is indol-2-yl optionally having substituents.


26. The compound of claim 23 or 24, wherein R²⁹ is indol-2-yl optionally having substituents selected from halogen atom, C₁₋₆ alkyl, C₁₋₆ alkoxy and hydroxy.

- 5 27. N-(2-((2,2-Diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)indole-2-carboxamide or a salt thereof,
N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)pentyl)amino)-2-oxoethyl)-1-methylindole-2-carboxamide or a salt thereof,
5-chloro-N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)-
10 pentyl)amino)-2-oxoethyl)indole-2-carboxamide or a salt thereof,
N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)indole-2-carboxamide or a salt thereof,
N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)-5-chloroindole-2-carboxamide or a
15 salt thereof,
N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)-1-methylindole-2-carboxamide or a salt thereof,
N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-
20 pentyl)amino)-2-oxoethyl)-5-fluoroindole-2-carboxamide or a salt thereof,
N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)-5-methoxyindole-2-carboxamide or a salt thereof,
25 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)-5-hydroxyindole-2-carboxamide or a salt thereof,
N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)-amino)-2-oxoethyl)indole-2-carboxamide or a salt thereof,
30 N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl)-amino)-2-oxoethyl)-1-methylindole-2-carboxamide or a salt thereof,
5-chloro-N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-


- diphenylpentyl) amino) -2-oxoethyl) -1-methylindole-2-carboxamide or a salt thereof,
- 5-chloro-N-(2-((5-(4-(2-fluorophenyl)piperidino)-2,2-diphenylpentyl) amino) -2-oxoethyl) indole-2-carboxamide or a salt thereof,
- 5 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl) amino) -2-oxoethyl) -5-fluoroindole-2-carboxamide or a salt thereof,
- N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl) amino) -2-oxoethyl) -5-methoxyindole-2-carboxamide or a salt thereof,
- 10 N-(2-((2,2-bis(4-chlorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl) amino) -2-oxoethyl) indole-2-carboxamide or a salt thereof,
- 15 N-(2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)-pentyl) amino) -2-oxoethyl) indole-2-carboxamide or a salt thereof,
- N-(2-((2,2-bis(4-fluorophenyl)-5-(4-phenylpiperidino)-pentyl) amino) -2-oxoethyl) -5-chloroindole-2-carboxamide or a salt thereof,
- 20 N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl) amino) -2-oxoethyl) indole-2-carboxamide or a salt thereof,
- N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-methoxyphenyl)-piperidino)pentyl) amino) -2-oxoethyl) -5-chloroindole-2-carboxamide or a salt thereof,
- 25 N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl) amino) -2-oxoethyl) indole-2-carboxamide or a salt thereof, or
- N-(2-((2,2-bis(4-fluorophenyl)-5-(4-(2-fluorophenyl)-piperidino)pentyl) amino) -2-oxoethyl) -5-chloroindole-2-carboxamide or a salt thereof.
- 30

28. A production method of a compound of claim 23, which

comprises reacting a compound represented by the formula




wherein each symbol is as defined in claim 23, or a salt thereof with a reactive derivative of an organic acid of the formula



wherein each symbol is as defined in claim 23.

29. A pharmaceutical composition containing a compound of claim 23.

30. A compound represented by the formula




wherein R^{26} and R^{27} are the same or different and each is hydrogen atom or C_{1-6} alkyl group; R^{30} is hydrogen atom, C_{1-6} alkyl group or optionally halogenated C_{1-6} alkyl-carbonyl group; Alk is C_{1-6} alkylene group optionally having substituents; Alk_2 and Alk_3 are the same or different and each is a bond or C_{1-6} alkylene group optionally having substituents; R^{31} is C_{6-10} aryl group optionally having substituents; X^2 is CH, C-OH or N; P^5 and Q^5 are the same or different and each is C_{1-6} alkylene group; Y^6 , Y^7 and Y^8 are the same or different and each is hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group, or a salt thereof or a prodrug thereof.

31. The compound of claim 30, wherein Alk is C₁₋₆ alkylene group optionally having substituents selected from the group consisting of halogen atom, hydroxy, amino and C₆₋₁₀ aryl; Alk₂ and Alk₃ are the same or different and each is a bond or C₁₋₆ alkylene group optionally having substituents selected from the group consisting of halogen atom, hydroxy, amino and C₆₋₁₀ aryl; R³¹ is C₆₋₁₀ aryl group optionally having substituents selected from the group consisting of halogen atom, C₁₋₆ alkyl, hydroxy, C₁₋₆ alkoxy and C₆₋₁₀ aryl.


32. N-(2-((2-((2,2-Diphenyl-5-(4-phenylpiperidino)pentyl)-amino)-2-oxoethyl)amino)-2-oxoethyl)-2,2,2-trifluoro-N-phenylacetamide or a salt thereof,
2-anilino-N-(2-((2,2-diphenyl-5-(4-phenylpiperidino)-pentyl)amino)-2-oxoethyl)acetamide or a salt thereof, or 2-(((benzylamino)carbonyl)amino)-N-(2,2-diphenyl-5-(4-phenylpiperidino)pentyl)acetamide or a salt thereof.

33. A production method of a compound of claim 30, which comprises reacting a compound represented by the formula



wherein each symbol is as defined in claim 30, or a salt thereof, with,

(1) when Alk₂ is C₁₋₆ alkylene group optionally having substituents, a reactive derivative of an organic acid compound of the formula



wherein each symbol is as defined in claim 30,


(2) when Alk_2 is a bond, a reactive derivative of the formula $\text{R}^{31}\text{-Alk}_3\text{-NR}^{30}\text{-CO-X}$ or $\text{R}^{31}\text{-Alk}_3\text{-NCO}$

wherein X is leaving group, and other symbols are as defined in claim 30.

5

34. A pharmaceutical composition containing a compound of claim 30.

35. A method for antagonizing melanin-concentrating hormone,
10 comprising administering, to a mammal, an effective amount of a compound of the formula




wherein Ar^1 and Ar^2 are each an aromatic group optionally having substituents,

15 P and Q are each a divalent aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which optionally has substituents,

R^1 and R^3 are each (i) hydrogen atom, (ii) acyl group or (iii) a hydrocarbon group optionally having substituents,

20 R^2 and R^4 are each (i) hydrogen atom, (ii) an alkyl group optionally having substituents or (iii) an alkylcarbonyl group optionally having substituents, R^1 and R^2 or R^3 and R^4 optionally form, together with the adjacent nitrogen atom, a monocyclic or fused nitrogen-containing heterocyclic group
25 optionally having substituents, and j is 0 or 1, or a salt thereof or a prodrug thereof.

36. Use of a compound of the formula




wherein Ar^1 and Ar^2 are each an aromatic group optionally having substituents,

P and Q are each a divalent aliphatic hydrocarbon group which optionally contains ether oxygen or ether sulfur in a carbon chain and which optionally has substituents,

R^1 and R^3 are each (i) hydrogen atom, (ii) acyl group or (iii) hydrocarbon group optionally having substituents,


R^2 and R^4 are each (i) hydrogen atom, (ii) an alkyl group optionally having substituents or (iii) an alkylcarbonyl group optionally having substituents, R^1 and R^2 or R^3 and R^4 optionally form, together with the adjacent nitrogen atom, a monocyclic or fused nitrogen-containing heterocyclic group optionally having substituents, and j is 0 or 1, or a salt thereof or a prodrug thereof, for production of a melanin-concentrating hormone antagonist.

37. A compound represented by the formula



wherein R^{27} is hydrogen atom, halogen atom, optionally halogenated C_{1-6} alkyl group or optionally halogenated C_{1-6} alkoxy group, or a salt thereof.

20



10/31/02 10/31/02

PTO/SB/106 (5-00)
Approved for use through 10/31/02. OMB 0651-0032
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

Declaration and Power of Attorney for Patent Application

特許出願宣言書及び委任状

Japanese Language Declaration

日本語宣言書

私は、以下に記名された発明者として、ここに下記の通り宣言する：

As a below named inventor, I hereby declare that

私の住所、郵便の宛先そして国籍は、私の氏名の後に記載された通りである。

My residence, post office address and citizenship are as stated next to my name

下記の名称の発明について、特許請求範囲に記載され、且つ特許が求められている発明主題に関して、私は、最初、最先且つ唯一の発明者である（唯一の氏名が記載されている場合）か、或いは最初、最先且つ共同発明者である（複数の氏名が記載されている場合）と信じている。

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

MCH ANTAGONISTS

上記発明の明細書はここに添付されているが、下記の欄がチェックされている場合は、この限りでない：

the specification of which is attached hereto unless the following box is checked:

☐ _____ の日に出願され、
この出願の米国出願番号またはPCT国際出願番号は、
_____ であり、且つ
_____ の日に補正された出願（該当する場合）

☒ was filed on September 19, 2000
as United States Application Number or
PCT International Application Number
PCT/JP00/06376 and was amended on
_____ (if applicable)

私は、上記の補正書によって補正された、特許請求範囲を含む上記明細書を検討し、且つ内容を理解していることをここに表明する。

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above

私は、連邦規則法典第37編規則1.56に記載されている、特許性について重要な情報を開示する義務があることを認める。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56

Burden Hour Statement This form is estimated to take 0.4 hours to complete. Time will vary depending upon the need of the individual case. Any comments on the amount of time you are required to complete this form should be sent to Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO Commissioner of Patents and Trademarks, Washington, DC 20231

Japanese Language Declaration (日本語宣言書)

私は、ここに、以下に記載した外国での特許出願または発明者証の出願、或いは米国以外の少なくとも一國を指定している米国法典第35編第365条(a)によるPCT国際出願について、同第119条(a)-(d)項又は第365条(b)項に基づいて優先権を主張するとともに、優先権を主張する本出願の出願日より前の出願日を有する外国での特許出願または発明者証の出願、或いはPCT国際出願については、いかなる出願も、下記の枠内をチェックすることにより示した。

I hereby claim foreign priority under Title 35, United States Code, Section 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application for which priority is claimed

Prior Foreign Application(s)

外国での先行出願

266278/1999

Japan

20/9/1999

(Number)
(番号)

(Country)
(国名)

(Day/Month/Year Filed)
(出願日/月/年)

221055/2000

Japan

17/7/2000

(Number)
(番号)

(Country)
(国名)

(Day/Month/Year Filed)
(出願日/月/年)

Priority Claimed

優先権主張

☒ ☐
Yes No
はい いいえ

☒ ☐
Yes No
はい いいえ

私は、ここに、下記のいかなる米国仮特許出願についても、その米国法典第35編第119条(e)項の利益を主張する。

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Application No.)
(出願番号)

(Filing Date)
(出願日)

私は、ここに、下記のいかなる米国出願についても、その米国法典第35編第120条に基づく利益を主張し、又米国を指定するいかなるPCT国際出願についても、その同第365条(c)に基づく利益を主張する。また、本出願の各特許請求の範囲の主題が、米国法典第35編第112条第1段に規定された態様で、先行する米国出願又はPCT国際出願に開示されていない場合においては、その先行出願の出願日と本国内出願日またはPCT国際出願日との間の期間中に入手された情報で、連邦規則法典第37編規則1.56に定義された特許性に関わる重要な情報について開示義務があることを承認する。

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of application

PCT/JP00/06376

September 19, 2000

pending

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可、係居中、放棄)

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可、係居中、放棄)

私は、ここに表明された私自身の知識に係わる陳述が真実であり、且つ情報と信ずることに基づく陳述が、真実であると信じられることを宣言し、さらに、故意に虚偽の陳述などを行った場合は、米国法典第18編第1001条に基づき、罰金または拘禁、若しくはその両方により処罰され、またそのような故意による虚偽の陳述は、本出願またはそれに対して発行されるいかなる特許も、その有効性に問題が生ずることを理解した上で陳述が行われたことを、ここに宣言する。

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon

Japanese Language Declaration (日本語宣言書)

委任状: 私は本出願を審査する手続を行い、且つ米国特許商標庁との全ての業務を遂行するために、記名された発明者として、下記の弁護士及び/または弁理士を任命する。(氏名及び登録番号を記載すること)

POWER OF ATTORNEY As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith (list name and registration number)

Mark Chao, Reg. No. 37293; Elaine M. Ramesh, Reg. No. 43032

書類送付先

Send Correspondence to:

Mark Chao, PhD JD

Intellectual Property Department

Takeda Pharmaceuticals North America, Inc

Suite 500-475 Half Day Road

Lincolnshire, IL 60069 USA

直通電話連絡先: (氏名及び電話番号)

Direct Telephone Calls to: (name and telephone number)

Mark Chao, PhD JD

Voice: (847)383-3391 Fax: (847)383-3481

Elaine M. Ramesh, PhD JD

Voice: (847)383-3391 Fax: (847)383-3481

唯一または第一発明者氏名

Full name of sole or first inventor

Kaneyoshi KATO

発明者の署名

日付

Inventor's signature

Date

Kaneyoshi Kato March 26, 2002

住所

Residence

2-40, Maruyamadai 2-chome, Kawanishi-shi, Hyogo 666-0152 Japan

国籍

Citizenship

Japan

郵便の宛先

Post Office Address

same as above

第二共同発明者がある場合、その氏名

Full name of second joint inventor, if any

Masaaki MORI

第二共同発明者の署名

日付

Second inventor's signature

Date

Masaaki Mori March 27, 2002

住所

Residence

8-5, Kasuga 3-chome, Tsukuba-shi, Ibaraki 305-0821 Japan

国籍

Citizenship

Japan

郵便の宛先

Post Office Address

same as above

(第三以下の共同発明者についても同様に記載し、署名をすること)

(Supply similar information and signature for third and subsequent joint inventors.)

Japanese Language Declaration (日本語宣言書)

第三共同発明者がある場合、その氏名		Full name of third joint inventor, if any <u>Nobuhiro SUZUKI</u>	
第三共同発明者の署名	日付	Third inventor's signature <u>Nobuhiro Suzuki</u>	Date March 26, 2002
住所		Residence 6-51, Fushiharacho, Nishinomiya-shi, Hyogo 663-8031 Japan	
国籍		Citizenship Japan	
郵便の宛先		Post Office Address same as above	
第四共同発明者がある場合、その氏名		Full name of fourth joint inventor, if any <u>Yukio SHIMOMURA</u>	
第四共同発明者の署名	日付	Fourth inventor's signature <u>Yukio Shimomura</u>	Date March 27, 2002
住所		Residence Takeda Yakuhin Matsushiro Residence 410, 12-1, Matsushiro 3-chome, Tsukuba-shi, Ibaraki 305-0035 Japan	
国籍		Citizenship Japan	
郵便の宛先		Post Office Address same as above	
第五共同発明者がある場合、その氏名		Full name of fifth joint inventor, if any <u>Shiro TAKEKAWA</u>	
第五共同発明者の署名	日付	Fifth inventor's signature <u>Shiro Takekawa</u>	Date March 26, 2002
住所		Residence 12-8-508, Miyanishicho, Nishinomiya-shi, Hyogo 662-0976 Japan	
国籍		Citizenship Japan	
郵便の宛先		Post Office Address same as above	

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Japanese Language Declaration
(日本語宣言書)

Page 5 of 5